LOCAL UNCERTAINTY INEQUALITIES FOR FOURIER SERIES

JOHN F. PRICE AND PAUL C. RACKI

ABSTRACT. Necessary and sufficient conditions are given on α, β and t for there to exist a constant K such that

$$\left(\sum_{n \in E} |\hat{f}(n)|^2 \right)^{1/2} \leq K |E|^\alpha \|f|x|^\beta\|_t,$$

for all $f \in L^1(\mathbf{T}^d)$ and finite $E \subset \mathbf{Z}^d$.

1. Introduction. The classical uncertainty principle inequality [2] states that

$$(1.1) \quad \|x|f\|_2 \|y|F\|_2 \geq (d/4\pi) \|f\|_2^2$$

for all functions $f \in L^2(\mathbf{R}^d)$, where the Fourier transform F is defined by

$$F(y) = \int_{\mathbf{R}^d} f(x) \exp(-2\pi i x \cdot y) dx \quad \text{for } y = (y_1, \dots, y_d) \in \mathbf{R}^d.$$

The natural generalization of this to a product $\mathbf{T}^d = (\mathbf{R}/2\pi\mathbf{Z})^d$ of circle groups fails since $\sum_{n \in \mathbf{Z}^d} |n|^2 |\hat{f}(n)|^2 = 0$ for all constant functions $f \in L^2(\mathbf{T}^d)$. (Here and below,

$$\hat{f}(n) = (2\pi)^{-d} \int_{\mathbf{T}^d} f(x) \exp(-in \cdot x) dx \quad \text{for } n = (n_1, \dots, n_d) \in \mathbf{Z}^d.)$$

Recently, local versions of (1.1) have been developed and applied in quantum physics [3, 4]. Roughly speaking, they assert that if a function is condensed, then not only is its Fourier transform broad, but it cannot be “too” localized. Some of these inequalities involve general L^p -norms and powers of $|x|$. For example, in [4] it is shown that, given $t \in [1, \infty]$ and $\beta \in \mathbf{R}$, there exists a constant K such that

$$\left(\int_E |F(y)|^2 dy \right)^{1/2} \leq Km(E)^{\beta - d/t^*} \|f|x|^\beta\|_t,$$

for all $f \in L^2(\mathbf{R}^d)$ and measurable $E \subset \mathbf{R}^d$, with $m(E) < \infty$ if and only if $d/t^* < \beta < d/t'$ or $(t, \beta) = (1, 0)$ or $(2, 0)$, where $t' = t/(t-1)$ and $t^* = 2t/(t-2)$. Furthermore, no other power of $m(E)$ apart from $\beta - d/t^*$ is possible. (Global extensions of (1.1) are given in [1].)

Here we establish the following analogous result for $\mathbf{T}^d$.

Received by the editors December 12, 1983 and, in revised form, February 27, 1984.
1980 *Mathematics Subject Classification*. Primary 42B05; Secondary 26D05, 42A05.

THEOREM. Given $t \in [1, \infty]$ and $\alpha, \beta \in \mathbf{R}$, there exists constant K such that for all functions $f \in L^1(\mathbf{T}^d)$,

$$(1.2) \quad \left(\sum_{n \in E} |\hat{f}(n)|^2 \right)^{1/2} \leq K |E|^\alpha \|f|x|^\beta\|_t$$

for all finite $E \subset \mathbf{Z}^d$ if and only if α, β, t satisfy the following conditions (see Figure 1):

- (i) $\beta < d/t'$ if $1 < t \leq \infty$, otherwise $\beta \leq d/t'$ if $t = 1$;
- (ii) $\alpha \geq \max\{0, -1/t^*\}$ if $\beta \leq 0$ and $1 \leq t \leq 2$, or $\beta < d/t^*$ and $t < 2$;
- (iii) $d\alpha \geq \beta - d/t^*$ if $\max\{0, d/t^*\} < \beta < d/t'$.

FIGURE 1a

FIGURE 1b

The shaded area of Figure 1(a) is the region of validity of the inequality for $1 \leq t \leq 2$ with the boundary $\beta = d/t'$ ($\alpha \geq \frac{1}{2}$) being included for $t = 1$. When $2 < t \leq \infty$ the region of validity is the shaded area of Figure 1(b).

2. Proof of sufficiency. Assuming the conditions of the theorem hold, let $N \subset \mathbf{T}^d$, $N' = \mathbf{T}^d - N$ be its complement, and f be a function in $L^2(\mathbf{T}^d)$. We thus have $f = f\chi_N + f\chi_{N'}$, where χ_N is the characteristic function on N , so that $\hat{f} = (f\chi_N)^\wedge + (f\chi_{N'})^\wedge$. Hence,

$$(2.1) \quad \left(\sum_{n \in E} |\hat{f}(n)|^2 \right)^{1/2} \leq \left(\sum_{n \in E} |(f\chi_N)^\wedge(n)|^2 \right)^{1/2} + \left(\sum_{n \in E} |(f\chi_{N'})^\wedge(n)|^2 \right)^{1/2}$$

by Minkowski's inequality; we shall estimate the two quantities separately. First, using Hölder's inequality,

$$(2.2) \quad \left(\sum_{n \in E} |(f\chi_N)^\wedge(n)|^2 \right)^{1/2} \leq \max\{ |(f\chi_N)^\wedge(n)| : n \in E \} |E|^{1/2} \leq \|f\chi_N\|_1 |E|^{1/2} \\ = \|f\chi_N|x|^\beta|x|^{-\beta}\|_1 |E|^{1/2} \leq \|f\chi_N|x|^\beta\|_t \|\chi_N|x|^{-\beta}\|_{t'} |E|^{1/2}.$$

Considering the second term of (2.1), we have

$$\begin{aligned}
 (2.3) \quad \left(\sum_{n \in E} |(f\chi_{N'})^\wedge(n)|^2 \right)^{1/2} &\leq \left(\sum_{n \in \mathbb{Z}^d} |(f\chi_{N'})^\wedge(n)|^2 \right)^{1/2} = \|f\chi_{N'}\|_2 \\
 &= \|f\chi_{N'}|x|^\beta |x|^{-\beta}\|_2 \leq \|f\chi_{N'}|x|^\beta\|_{2r} \|\chi_{N'}|x|^{-\beta}\|_{2r'} \\
 &= \|f\chi_{N'}|x|^\beta\|_t \|\chi_{N'}|x|^{-\beta}\|_{t^*},
 \end{aligned}$$

where we used Hölder's inequality with $r = t/2$ and assumed $t > 2$.

If we assume that $t \in [1, 2]$ we estimate the second term as follows:

$$\begin{aligned}
 (2.4) \quad \left(\sum_{n \in E} |(f\chi_{N'})^\wedge(n)|^2 \right)^{1/2} &= \left(\sum_{n \in E} |(f\chi_{N'})^\wedge(n) \cdot 1|^2 \right)^{1/2} \\
 &\leq \left(\sum_{n \in E} |(f\chi_{N'})^\wedge(n)|^{2r} \right)^{1/2r} \left(\sum_{n \in E} 1^{2r'} \right)^{1/2r'} \\
 &= \left(\sum_{n \in E} |(f\chi_{N'})^\wedge(n)|^{t'} \right)^{1/t'} |E|^{-1/t^*} \\
 &\leq \left(\sum_{n \in \mathbb{Z}^d} |(f\chi_{N'})^\wedge(n)|^{t'} \right)^{1/t'} |E|^{-1/t^*} \\
 &\leq \|f\chi_{N'}\|_t |E|^{-1/t^*} = \|f\chi_{N'}|x|^\beta |x|^{-\beta}\|_t |E|^{-1/t^*} \\
 &\leq \|f\chi_{N'}|x|^\beta\|_t \|\chi_{N'}|x|^{-\beta}\|_\infty |E|^{-1/t^*},
 \end{aligned}$$

where we used Hölder's inequality with $r = t'/2$, followed by Hausdorff-Young and a final application of Hölder's inequality.

Substituting (2.2)–(2.4) into (2.1), we obtain

$$(2.5) \quad \left(\sum_{n \in E} |\hat{f}(n)|^2 \right)^{1/2} \leq K \|f|x|^\beta\|_t,$$

where $K = K_1 + K_2$, with

$$\begin{aligned}
 K_1 &= \|\chi_N |x|^{-\beta}\|_t |E|^{1/2}, \\
 K_2 &= \begin{cases} \|\chi_{N'} |x|^{-\beta}\|_\infty |E|^{-1/t^*} & \text{for } 1 \leq t \leq 2, \\ \|\chi_{N'} |x|^{-\beta}\|_{t^*} & \text{for } 2 < t \leq \infty. \end{cases}
 \end{aligned}$$

Letting $N = \{x \in \mathbb{T}^d : |x| < a\}$ for some a , we now evaluate the above norms. First,

$$\begin{aligned}
 (2.6) \quad \|\chi_N |x|^{-\beta}\|_{t'}^{t'} &= \int_{|x| \leq a} |x|^{-\beta t'} dx = W_d \int_0^a r^{-\beta t'} r^{d-1} dr \\
 &= (W_d / (d - \beta t')) a^{d - \beta t'},
 \end{aligned}$$

as $\beta < d/t'$, where $W_d = 2\pi^{d/2}/\Gamma(d/2)$. Second

$$(2.7) \quad \left\| \chi_{N'} |x|^{-\beta} \right\|_{t^*}^{t^*} = \int_{a \leq |x| \leq \pi} |x|^{-\beta t^*} dx = W_d \int_a^\pi r^{-\beta t^* + d-1} dr \\ = W_d (\beta t^* - d)^{-1} (a^{d-\beta t^*} - \pi^{d-\beta t^*}) \leq W_d (\beta t^* - d)^{-1} a^{d-\beta t^*},$$

as $\beta > \max\{0, d/t^*\}$. Thirdly if $\beta > 0$,

$$(2.8) \quad \left\| \chi_{N'} |x|^{-\beta} \right\|_\infty = \sup\{|x|^{-\beta} : a \leq |x| \leq \pi\} = a^{-\beta}.$$

Now, letting $a = |E|^{-1/d}$ and substituting (2.6)–(2.8) into (2.5), we obtain

$$\left(\sum_{n \in E} |\hat{f}(n)|^2 \right)^{1/2} \leq K |E|^{\beta/d-1/t^*} \|f|x|^\beta\|_t,$$

with $K = K_1 + K_2$, where $K_1 = (W_d/(d - \beta t'))^{1/t'}$ and

$$K_2 = \begin{cases} 1 & \text{for } 1 \leq t \leq 2, \\ (W_d/(\beta t^* - d))^{1/t^*} & \text{for } 2 < t \leq \infty, \end{cases}$$

provided $\max\{0, d/t^*\} < \beta < d/t'$.

Finally, if $\beta \leq 0$, $\|\chi_{N'} |x|^{-\beta}\|_\infty = \pi$; hence, $K_2 = \pi |E|^{-1/t^*}$ and

$$K_1 = (W_d/(d - \beta t'))^{1/t'} a^{d/t'-\beta} |E|^{1/2} = (W_d/(d - \beta t'))^{1/t'} |E|^{-1/t^*}$$

upon letting $a = |E|^{-1/(d-\beta t')}$ for $t \neq 1$. If $t = 1$,

$$K_1 = \left\| \chi_N |x|^{-\beta} \right\|_\infty |E|^{1/2} = a^{-\beta} |E|^{1/2} = |E|^{-1/t^*}$$

upon letting $a = 1$. Thus

$$\left(\sum_{n \in E} |\hat{f}(n)|^2 \right)^{1/2} \leq K |E|^{-1/t^*} \|f|x|^\beta\|_t,$$

for $\beta \leq 0$ and $1 \leq t \leq 2$ with $K = K_1 + K_2$, where $K_2 = \pi$ and

$$K_1 = \begin{cases} 1 & \text{for } t = 1, \\ (W_d/(d - \beta t'))^{1/t'} & \text{for } 1 < t \leq 2. \end{cases}$$

We thus have the inequality holding along the line segment in the (α, β) -plane given by $d\alpha = \beta - d/t^*$ for $\max\{0, d/t^*\} < \beta < d/t'$. When $t = 1$ we also have it holding at the endpoint $\beta = d/t' = 0$ since

$$\left(\sum_{n \in E} |\hat{f}(n)|^2 \right)^{1/2} \leq |E|^{1/2} \|\hat{f}\|_\infty \leq |E|^{1/2} \|f\|_1.$$

These results may be extended as follows: If

$$\left(\sum_{n \in E} |\hat{f}(n)|^2 \right)^{1/2} \leq K |E|^\alpha \|f|x|^\beta\|_t,$$

holds for given (α, β, t) , then it holds for (α', β', t') with K replaced by $K \| |x|^{\beta-\beta'} \|_\infty$, where $\alpha' \geq \alpha$, $\beta' \leq \beta$ and $t' \geq t$. To see this, notice that $|E|^{\alpha'} \geq |E|^\alpha$ and $\|f|x|^\beta\|_t \leq \| |x|^{\beta-\beta'} \|_\infty \|f|x|^{\beta'}\|_{t'}$.

This leaves only the case $\alpha = 0$ and $t \geq 2$; it can be established as follows:

$$\|\hat{f}\|_2 = \|f\|_2 = \|f|x|^\beta|x|^{-\beta}\|_2 \leq \|f|x|^\beta\|_t \| |x|^{-\beta} \|_{t^*}$$

with

$$\| |x|^{-\beta} \|_{t^*} = W_d \int_0^\pi r^{-\beta t^* + d-1} dr < \infty \quad \text{for } \beta < d/t^* (t > 2)$$

and

$$\| |x|^{-\beta} \|_{t^*} = \sup \{ |x|^{-\beta} : x \in \mathbf{T}^d \} < \infty \quad \text{for } \beta \leq d/t^* = 0 (t = 2).$$

We have thus covered the stated region, and the sufficiency of the conditions is established.

REMARK. In the interior of the region an alternate value of the constant is $K = \| |x|^{-\beta} \|_{2t/(t+2\alpha t-2)}$. To see this, notice that

$$\begin{aligned} \left(\sum_{n \in E} |\hat{f}(n)|^2 \right)^{1/2} &\leq |E|^{1/2r'} \left(\sum_{n \in \mathbb{Z}^d} |\hat{f}(n)|^{2r} \right)^{1/2r} \\ &\leq |E|^{1/2r'} \|f|x|^\beta|x|^{-\beta}\|_{(2r)'} \\ &\leq |E|^{1/2r'} \| |x|^{-\beta} \|_{(2r)'} \|f|x|^\beta\|_{(2r)'} \\ &= |E|^\alpha \| |x|^{-\beta} \|_{2t/(t+2\alpha t-2)} \|f|x|^\beta\|_t. \end{aligned}$$

3. Proof of necessity. The necessity of the conditions is obtained by ruling out the remaining regions beginning with $\alpha \leq 0$. Firstly, we can not have $\alpha < 0$, since then $|E|^\alpha \rightarrow 0$ as $|E| \rightarrow \infty$. Secondly, considering the case $\alpha = 0$, $\beta = d/t^*$ with $t > 2$, the function $f \in L^1(T^d)$ defined by

$$f(x) = \begin{cases} |x|^{-d/2} |\log|x||^{-1/2} & \text{for } |x| \leq 1/2, \\ 0 & \text{otherwise} \end{cases}$$

provides a counterexample, since

$$\|f\|_2^2 = W_d \int_0^{1/2} r^{-1} \log^{-1} r dr = \infty,$$

while

$$\|f|x|^{d/t^*}\|_t^t = W_d \int_0^{1/2} r^{-1} |\log r|^{-t/2} dr < \infty \quad \text{if } t > 2.$$

This also rules out the region $\alpha = 0$, $\beta > d/t^*$ for $t > 2$. Thirdly, consider the region given by $\alpha = 0$, $\beta < d/t^*$ with $1 \leq t < 2$. Here define $f(x) = |x|^{-1} |x|^{-1/2}$ for $1/2 \leq |x| \leq 3/2$ and 0 otherwise, so that

$$\|f\|_2^2 = W_d \int_{1/2}^{3/2} |r-1|^{-1} r^{d-1} dr = \infty,$$

while

$$\|f|x|^\beta\|_t^t = W_d \int_{1/2}^{3/2} |r-1|^{-t/2} r^{\beta t + d-1} dr < \infty \quad \text{for } t < 2.$$

Finally, the region given by $\alpha = 0$, $\beta > d/t^* = 0$ for $t = 2$ is ruled out by $f(x) = |x|^{-d/2}$ since

$$\|f\|_2^2 = W_d \int_0^\pi r^{-1} dr = \infty,$$

while

$$\|f|x|^\beta\|_2 = W_d \int_0^\pi r^{-1+2\beta} dr < \infty,$$

which completes the $\alpha \leq 0$ case.

Now consider the boundary $\beta \geq d/t'$; here the function

$$f_\varepsilon(x) = \begin{cases} |x|^{-d} |\log|x||^{-1} & \text{for } 0 < \varepsilon \leq |x| \leq \frac{1}{2}, \\ 0 & \text{otherwise} \end{cases}$$

provides the required counterexample as $\varepsilon \rightarrow 0$. For assume that $E = \{0\}$; then

$$\left(\sum_{n \in E} |\hat{f}_\varepsilon(n)|^2 \right)^{1/2} = |\hat{f}_\varepsilon(0)| = \left| W_d \int_\varepsilon^{1/2} r^{-1} \log^{-1} r dr \right| \rightarrow \infty \quad \text{as } \varepsilon \rightarrow 0,$$

while

$$\|f_\varepsilon|x|^\beta\|_t^t = W_d \int_\varepsilon^{1/2} r^{\beta t - dt + d - 1} |\log r|^{-t} r dr \leq \text{constant}$$

for all $\varepsilon > 0$ if $\beta t - dt + d - 1 \leq -1$ and $t > 1$ or $\beta t - dt + d - 1 > -1$ and $t = 1$.

Hence, there is a contradiction for all α if $\beta \geq d/t'$ and $t > 1$ or $\beta > d/t'$ and $t = 1$.

Consider now the region $d\alpha < \beta - d/t^*$ for $\max\{0, d/t^*\} \leq \beta < d/t'$. Define

$$f_N(x) = \begin{cases} 1 & \text{if } x \in \square_N, \\ 0 & \text{otherwise,} \end{cases}$$

where

$$\square_N = \{x = (x_1, \dots, x_d) \in T^d: |x_1| \leq 1/N, \dots, |x_d| \leq 1/N\}$$

and N is a positive integer. Now

$$\begin{aligned} \hat{f}_N(n) &= \hat{f}_N(n_1, \dots, n_d) = \int_{\square_N} e^{-in_1 x_1} \dots e^{-in_d x_d} dx_1 \dots dx_d \\ &= 2^d n_1^{-1} \sin n_1/N \dots n_d^{-1} \sin n_d/N. \end{aligned}$$

Thus, letting $E = \{n = (n_1, \dots, n_d) \in \mathbb{Z}^d: |n_1| \leq N, \dots, |n_d| \leq N\}$, we have

$$\begin{aligned} \left(\sum_{n \in E} |\hat{f}_N(n)|^2 \right)^{1/2} &= 2^d \left(\sum_{|n_1| \leq N} \frac{\sin^2 n_1/N}{n_1^2} \dots \sum_{|n_d| \leq N} \frac{\sin^2 n_d/N}{n_d^2} \right)^{1/2} \\ &\sim 1/N^{d/2} \quad \text{as } N \rightarrow \infty \end{aligned}$$

since

$$\lim_{N \rightarrow \infty} \sum_{|n| \leq N} \left(\frac{n}{N} \right)^{-2} \sin^2 \left(\frac{n}{N} \right) N^{-1} = \int_{-1}^1 x^{-2} \sin^2 x dx,$$

while

$$\begin{aligned} |E|^\alpha \|f_N |x|^\beta\|_t &= N^{d\alpha} \left(\int_{\square_N} |x|^{\beta t} dx \right)^{1/t} \\ &\leq N^{d\alpha} \left(\int_{|x| \leq d^{1/2}/N} |x|^{\beta t} dx \right)^{1/t} \\ &= (W_d/(\beta t + d))^{1/t} d^{(\beta/2)+(d/2t)} N^{d\alpha - \beta - d/t}, \end{aligned}$$

assuming $\beta > d/t^* > -d/t$. We thus have a contradiction if $-d/2 > d/\alpha - \beta - d/t$, that is, if $d\alpha < \beta - d/t^*$, as required.

Finally, define $g_N(x) = f_N(x - \mathbf{1})$, f_N as above, and $\mathbf{1} = (1, \dots, 1)$, for $N > 1$ being an integer. With E as above we have

$$\left(\sum_{n \in E} |\hat{g}_N(n)|^2 \right)^{1/2} \sim N^{-d/2} \quad \text{as } N \rightarrow \infty$$

since $|\hat{g}_N(n)| = |\hat{f}_N(n)|$. Therefore consider, for $\beta < 0$,

$$\begin{aligned} |E|^\alpha \|g_N |x|^\beta\|_t &= N^{d\alpha} \left(\int_{\square_{N-1}} |x|^{\beta t} dx \right)^{1/t} = N^{d\alpha} \left(\int_{\square_N} |y + 1|^{\beta t} dy \right)^{1/t} \\ &\leq \left(\frac{1}{2} \right)^\beta N^{d\alpha} \left(\int_{\square_N} dy \right)^{1/t} = 2^{d/t - \beta} N^{d\alpha} N^{-d/t}. \end{aligned}$$

We thus obtain a contradiction as $N \rightarrow \infty$ if $-d/2 > d\alpha - d/t$, that is, $\alpha < -1/t^*$ for $\beta < 0$.

All the required regions are now eliminated and necessity is established which completes the proof of the Theorem.

REFERENCES

1. M. G. Cowling and J. F. Price, *Bandwidth versus time concentration: the Heisenberg-Pauli-Weyl inequality*, SIAM J. Math. Anal. **15** (1984), 151–165.
2. D. Dym and H. P. McKean, *Fourier series and integrals*, Academic Press, New York, 1972.
3. W. G. Faris, *Inequalities and uncertainty principles*, J. Math. Phys. **19** (1978), 461–466.
4. J. F. Price, *Inequalities and local uncertainty principles*, J. Math. Phys. **27** (1983), 1711–1714.

SCHOOL OF MATHEMATICS, UNIVERSITY OF NEW SOUTH WALES, KENSINGTON, NEW SOUTH WALES 2033, AUSTRALIA