CHARACTERIZATIONS OF NOETHERIAN AND HEREDITARY RINGS

ZHENG - XU HE

ABSTRACT. We characterize the left Noetherian rings by the existence of decompositions of left modules into direct sums of an injective submodule and a submodule containing no injective submodule except 0. We also prove that a left Noetherian ring is left hereditary iff the suspension of each left ideal (see [7]) is injective or, equivalently, the above decomposition is unique.

0. Let A be an Abelian group. Then $A = D \oplus B$, where D is divisible and B is reduced in the sense of Kaplansky [8]; that is, B has no divisible submodule except 0. If this holds for left modules over a unitary ring R , with “injective” and “ i -reduced” replacing “divisible” and “reduced”, respectively, then R must be left Noetherian. The proof requires the theorem of Eilenberg-Bass (see, e.g., Theorem 20.1, p. 111, of [2]) which characterizes left Noetherian by the requirement that direct sums of injective left modules are injective. Furthermore, if the decomposition, which will be referred to as an i -decomposition, is unique in the sense that D is the unique injective submodule of A for which $A = D \oplus B$, with B i -reduced, then every sum of injective submodules is injective, and hence R must be left hereditary by a theorem of Matlis [9]. At the same time, it will be shown that a left Noetherian ring is left hereditary iff the suspension of each left ideal is injective.

1. Throughout this paper, R denotes a unitary ring.

THEOREM 1. *The following conditions are equivalent:*

- (i) R is left Noetherian.
- (ii) *The direct sum of any countably infinite family of injective left R -modules is injective.*
- (iii) *Any left R -module admits an i -decomposition.*
- (iv) *Every injective left R -module is a direct sum of indecomposable left R -modules.*

PROOF. The equivalence (i) $\Leftrightarrow$ (ii) is known, and due to Eilenberg-Bass. The implication (i) $\Rightarrow$ (iii) is immediate; and (i) $\Leftrightarrow$ (iv) is the theorem of Matlis [9] and Papp [10]. Let us prove (iii) $\Rightarrow$ (ii). Let $\{S_n; n = 1, 2, 3, \dots\}$ be a countably infinite

Received by the editors December 19, 1983 and, in revised form, May 23, 1984.

1980 *Mathematics Subject Classification.* Primary 13E05; Secondary 18G05.

Key words and phrases. Injective module, i -reduced module, i -decomposition, unique i -decomposition, Noetherian ring, left hereditary ring, FP-injective module, suspension of a module.

©1985 American Mathematical Society
 0002-9939/85 \$1.00 + \$.25 per page

family of injective left R -modules. Let D and B realize an i -decomposition of $S = \bigoplus_{n=1}^{\infty} S_n$. For each positive integer k , $T_k = \bigoplus_{n=1}^k S_n$ is an injective submodule of S . B being i -reduced, we must have $E(T_k \cap D) = T_k$ by which we deduce that $T_k \subseteq D$, where $E(T_k \cap D)$ denotes the injective hull (included in T_k) of $T_k \cap D$. In this way, $S = D$, which is injective. Q.E.D.

2. For convenience, we will “borrow” some notation from the homotopy theory of modules introduced by B. Eckmann and P. Hilton (for more details, see [7, Chapter 13; 6 and 5]). For any left R -module A , we define the “suspension” of A to be $S(A) = E(A)/A$, where $E(A)$ denotes the injective hull of A . Let $f: A \rightarrow B$ be a morphism of left R -modules; we denote by $f \simeq_i 0$ the fact that f can be extended to any module containing A . We write $\bar{\pi}(A, B) = 0$ if, for any $f: A \rightarrow B$, $f \simeq_i 0$. We say that a left R -module B is *FG-injective* if $\bar{\pi}(A, B) = 0$ for any finitely generated left R -module A . The following properties hold:

- (1) $f \simeq_i 0$ iff f can be extended to the injective hull $E(A)$ of A or, equivalently, to any injective extension E of A .
- (2) If $f = g + h$, then $g \simeq_i 0$ and $h \simeq_i 0$ imply $f \simeq_i 0$.
- (3) If $f = \alpha \circ \beta$, then either $\alpha \simeq_i 0$ or $\beta \simeq_i 0$ implies $f \simeq_i 0$.
- (4) For any injective extension E of A , $S(A)$ is injective (or FG-injective, or FP-injective) iff E/A is.
- (5) If $\bar{\pi}(I, B) = 0$ for each left ideal I of R , then B is injective.
- (6) B is FP-injective iff $\bar{\pi}(A, B) = 0$ for any finitely presented left R -module A .

LEMMA. Assume that R satisfies:

(*) For any left ideal I of R , $S(I)$ is FG-injective.

Then the sum of any FG-injective submodules of a left R -module is FG-injective and hence FP-injective.

PROOF. Let $\{V_i; i \in J\}$ be a family of FG-injective submodules of a left R -module; we need to show that $\sum_{i \in J} V_i$ is FG-injective. We may reduce the problem to the case when J is finite, and so it suffices to consider $J = \{1, 2\}$. For any positive integer m , let $\mathfrak{F}^m$ be the family of all R -modules which can be spanned by m elements. We will prove the following by induction on m :

(**) $\bar{\pi}(F, V_1 + V_2) = 0$ for any $F \in \mathfrak{F}^m$.

Firstly, for $m = 1$, let $F = Ra \in \mathfrak{F}^1$ and let $f: F \rightarrow V_1 + V_2$. $f(a) = v_1 + v_2$ for some $v_i \in V_i$, $i = 1, 2$. Since V_i is FG-injective, there exists $\alpha_i: E(R) \rightarrow V_i$ such that $\alpha_i(r) = rv_i$ for all $r \in R$. Let $I = \{r \in R; r(v_1 + v_2) = 0\}$; and set $S = E(R)/I$ which is FG-injective by (*). Define $\beta: F \rightarrow S$ and $\alpha: S \rightarrow V_1 + V_2$ by $\beta(ra) = [r] \in S = E(R)/I$ and $\alpha([r]) = \alpha_1(\bar{r}) + \alpha_2(\bar{r})$ for $r \in R$ and $\bar{r} \in E(R)$. We get $f = \alpha \circ \beta$. Now $\beta \simeq_i 0$, because S is FG-injective. So $f \simeq_i 0$, and thus (**) holds for $m = 1$.

Next, we prove (**) for $m > 1$, assuming that (**) is true if m is replaced by $m - 1$. Let $F = Ra_1 + Ra_2 + \cdots + Ra_m \in \mathfrak{F}^m$. We need show that $f \simeq_i 0$ for $f: F \rightarrow V_1 + V_2$. Consider $f_1 = f/Ra_1$. By the inductive hypothesis, $f_1 \simeq_i 0$, i.e. there is $f'_1: E(F) \rightarrow V_1 + V_2$ such that $f'_1/Ra_1 = f_1$. Let $g = f'_1/F$ and let $h = f - g$. Then

$h(a_1) = 0$, and so there exists $\alpha: H = F/(Ra_1) \rightarrow V_1 + V_2$ such that $h = \alpha \circ p$, where $p: F \rightarrow H = F/(Ra_1)$ is the canonical projection. Clearly, $H \in \mathfrak{F}^m$; again by the inductive hypothesis, $\alpha \simeq_i 0$, whence $h \simeq_i 0$. But obviously, $g \simeq_i 0$; thus $f = g + h \simeq_i 0$. Q.E.D.

As a consequence, if R is a left Noetherian ring verifying $(*)$, then $\sum_{i \in J} V_i$ is injective when V_i ($i \in J$) all are.

We also observe that if $\{V_i; i \in J\}$ is a family of injective submodules of a left module over a left Noetherian ring, and if $\sum_{i \in J} V_i$ admits a unique i -decomposition, then $\sum_{i \in J} V_i$ is injective. At this stage, using a theorem of Matlis [9] (see, e.g., Proposition 10 of [3]), we may conclude

THEOREM 2. *The following conditions are equivalent:*

- (i) *R is left Noetherian and left hereditary.*
- (ii) *R is left Noetherian and satisfies $(*)$.*
- (iii) *Any left R -module admits a unique i -decomposition.*
- (iv) *Every sum of injective submodules of a left R -module is injective.*

The equivalence (i) $\Leftrightarrow$ (ii) includes Corollary 14.B of [4].

ACKNOWLEDGEMENT. The author wishes to thank the referee for useful suggestions.

REFERENCES

1. C. Faith, *Algebra I: Rings, modules, and categories*, Springer-Verlag, Berlin, Heidelberg, and New York, 1973; corrected reprint, 1981.
2. ———, *Algebra I: Ring theory*, Springer-Verlag, Berlin and New York, 1976.
3. ———, *Lectures on injective modules and quotient rings*, Lecture Notes in Math., Vol. 49, Springer-Verlag, Berlin and New York, 1967.
4. ———, *On hereditary rings and Boyle's conjecture*, Arch. Math. **27** (1976), 113–119.
5. Zheng-Xu He, *Some results on homotopy theory of modules*, Rend. Accad. Naz. Lincei **74** (1983), 366–372.
6. ———, *On weak i -homotopy equivalences of modules*, Rend. Accad. Naz. Lincei **76** (1984).
7. P. Hilton, *Homotopy theory and duality*, Notes on Math. and its Appl., Gordon & Breach, New York, London and Paris, 1965.
8. I. Kaplansky, *Infinite abelian groups*, Univ. of Michigan Press, Ann Arbor, 1969.
9. E. Matlis, *Injective modules over Noetherian rings*, Pacific J. Math. **8** (1958), 511–528.
10. Z. Papp, *On algebraically closed rings*, Publ. Math. Debrecen **6** (1959), 311–327.

FACULTY OF MATHEMATICS, UNIVERSITY OF IASI, R - 6600, IASI, ROMANIA