WEAK CONVERGENCE THEOREMS FOR NONEXPANSIVE MAPPINGS AND SEMIGROUPS IN BANACH SPACES WITH OPIAL'S PROPERTY

T. KUCZUMOW

ABSTRACT. Let X be a Banach space with Opial's property, C a weakly compact subset of X , $x \in C$ and S a nonexpansive semigroup on C . Then $\{S(t)x\}_{t \geq 0}$ converges weakly to a common fixed point of S iff $S(t+h)x - S(t)x \rightarrow 0$ as $t \rightarrow \infty$ for all $h > 0$.

In this paper we consider a necessary and sufficient condition for the weak (weak-*) convergence of trajectories of nonexpansive mappings and one parameter semigroups of nonexpansive mappings in Banach spaces which satisfy Opial's condition. This problem was studied in [1, 2, 5, 6, 7, 9].

A Banach space X satisfies Opial's condition [4] for the weak topology if $x_n \rightarrow y$ in X implies that

$$\limsup_n \|x_n - y\| < \limsup_n \|x_n - z\|$$

for all $z \neq y$.

All spaces l^p ($1 < p < \infty$) have Opial's property but $L^p[0, 2\pi]$ with $1 < p \neq 2$ lacks it [4].

A Banach space X satisfies Opial's condition for the weak-* topology if X is a conjugate space to a separable Banach space and for $x_n \rightarrow *y$ in X we have

$$\limsup_n \|x_n - y\| < \limsup_n \|x_n - z\|$$

for $z \neq y$.

l^1 satisfies Opial's condition for the weak-* topology [3]. l^∞ does not have Opial's property.

Let C be a subset of a Banach space X . A mapping $T: C \rightarrow C$ is said to be nonexpansive if $\|Tx - Ty\| \leq \|x - y\|$ for all $x, y \in C$. A one parameter semigroup $S = \{S(t): t \geq 0\}$ of nonexpansive mappings on C is a family of nonexpansive mappings of C into itself satisfying the following conditions:

- (i) $S(s + t)x = S(s)S(t)x$ for $s, t \geq 0$ and $x \in C$,

Received by the editors July 1, 1983 and, in revised form, January 5, 1984.

1980 *Mathematics Subject Classification*. Primary 47H10.

Key words and phrases. Nonexpansive mappings, nonexpansive semigroups, fixed points, Opial's property.

©1985 American Mathematical Society
0002-9939/85 \$1.00 + \$.25 per page

- (ii) $\|S(t)x - S(t)y\| \leq \|x - y\|$ for $t \geq 0$ and $x, y \in C$.
 (iii) $S(0)x = x$ for $x \in C$.

LEMMA. Let C be a weakly (weakly-*) compact subset of a Banach space X with Opial's property for the weak (weak-*) topology, $z \in X$ and $\{x_n^m: m \in \mathbf{N}\}$ be a family of sequences which satisfies the following conditions:

- (i) $x_n^m \in C$ for $n, m \in \mathbf{N}$,
 (ii) every $\{x_n^m\}$ converges to y in the weak (weak-*) topology,
 (iii) $\limsup_n \|x_n^m - y\| \geq \limsup_n \|x_n^{m+1} - z\| \geq \limsup_n \|x_n^{m+1} - y\|$.
 Then $y = z$.

PROOF. Let $r_m = \limsup_n \|x_n^m - y\|$ for $m \in \mathbf{N}$. Because

$$C \cap \lim\left\{w \in X: \bigvee_{m,n} w = x_n^m \vee w = y \vee w = z\right\} \quad (C)$$

with the weak (weak-*) topology is metrized by some metric ρ [8], then for every $m \geq 2$ we may choose $x_{n_m}^m$ such that

$$\rho(x_{n_m}^m, y) \leq 1/m, \quad \|x_{n_m}^m - y\| \geq r_m - 1/m, \quad \|x_{n_m}^m - z\| \leq r_{m-1} + 1/m.$$

Hence we have

$$\begin{aligned} x_{n_m}^m &\rightharpoonup y \quad (x_{n_m}^m \rightharpoonup *y), \\ \limsup_m \|x_{n_m}^m - z\| &= \lim_m r_m = \limsup_m \|x_{n_m}^m - y\| \end{aligned}$$

and therefore $y = z$.

THEOREM 1. Let X and C be such as in the Lemma, $T: C \rightarrow C$ be a nonexpansive mapping and $x \in C$. $\{T^i x\}$ converges weakly (weakly-*) to a fixed point of T iff $T^{i+1}x - T^i x \rightarrow_i 0$ ($T^{i+1}x - T^i x \rightarrow_i *0$).

PROOF. There exists a subsequence $\{T^{i_n} x\}$ which converges weakly (weakly-*) to y . Then for $m \geq 1$ we have

$$T^{i_n+m} x \rightharpoonup y \quad (T^{i_n+m} x \rightharpoonup *y)$$

and

$$\begin{aligned} \limsup_n \|T^{i_n+m} x - y\| &\geq \limsup_n \|T^{i_n+m+1} x - Ty\| \\ &\geq \limsup_n \|T^{i_n+m+1} x - y\|. \end{aligned}$$

By the Lemma, $y = Ty$ and thus in view of monotonicity of $\{\|T^i x - v\|\}$ for any $v \in \text{Fix } T$, the sequence $\{T^i x\}$ tends weakly to y ($T^i x \rightarrow_i *y$).

THEOREM 2. Let X and C be such as in the Lemma, $S = \{S(t): t \geq 0\}$ be a semigroup of nonexpansive mappings on C and let $x \in C$. Then $\{S(t)x\}_{t \geq 0}$ converges weakly (weakly-*) to a common fixed point of S iff

$$\begin{aligned} S(t+h)x - S(t)x &\rightarrow 0 \quad \text{as } t \rightarrow \infty \\ (S(t+h)x - S(t)x) &\rightarrow *0 \quad \text{as } t \rightarrow \infty \end{aligned}$$

for all $h > 0$.

PROOF. We choose $\{t_n\}$ such that $t_n \rightarrow_n \infty$ and $S(t_n)x \rightarrow_n y$ ($S(t_n)x \rightarrow_n *y$). For $t > 0$ and $m \in \mathbb{N}$ we have

$$S(t_n + mt)x - S(t_n)x \rightarrow_n 0 \quad \left(S(t_n + mt)x - S(t_n)x \rightarrow_n *0 \right)$$

and hence

$$\begin{aligned} \limsup_n \|S(t_n + mt)x - y\| &\geq \limsup_n \|S(t_n + (m+1)t)x - S(t)y\| \\ &\geq \limsup_n \|S(t_n + (m+1)t)x - y\|. \end{aligned}$$

Therefore $y = S(t)y$ for all $t \geq 0$ and $S(t) \rightarrow y$ as $t \rightarrow \infty$ ($S(t) \rightarrow *y$ as $t \rightarrow \infty$).

REMARK. Theorem 2 gives a partial answer to a question raised on p. 550 of [7].

REFERENCES

1. J. B. Baillon, R. E. Bruck and S. Reich, *On the asymptotic behavior of nonexpansive mappings and semigroups in Banach spaces*, Houston J. Math. **4** (1978), 1–9.
2. N. Hirano, *Nonlinear ergodic theorems and weak convergence theorems*, J. Math. Soc. Japan **34** (1982), 35–46.
3. L. A. Karlovitz, *On nonexpansive mappings*, Proc. Amer. Math. Soc. **55** (1976), 321–325.
4. Z. Opial, *Weak convergence of the sequence of successive approximations for nonexpansive mappings*, Bull. Amer. Math. Soc. **73** (1967), 591–597.
5. A. Pazy, *On the asymptotic behavior of semigroups of nonlinear contractions in Hilbert space*, J. Funct. Anal. **27** (1978), 292–307.
6. S. Reich, *Weak convergence theorems for nonexpansive mappings in Banach space*, J. Math. Anal. Appl. **67** (1979), 274–276.
7. ———, *A note on the mean ergodic theorem for nonlinear semigroups*, J. Math. Anal. Appl. **91** (1983), 547–551.
8. S. Rolewicz, *Metric linear spaces*, PWN, Warsaw, 1972.
9. R. Schoeneberg, *Weak convergence of trajectories of nonexpansive semigroups in Hilbert space*, Israel J. Math. **30** (1978), 130–132.

INSTYTUT MATEMATYKI UMCS, PLAC MARII CURIE - SKŁODOWSKIEJ 1, 20 - 031 LUBIN, POLAND