A NOTE ON THE ABSOLUTE CONVERGENCE OF LACUNARY FOURIER SERIES

N. V. PATEL AND V. M. SHAH

ABSTRACT. P. B. Kennedy [3] studied lacunary Fourier series whose generating functions are of bounded variation on a subinterval I of $[-\pi, \pi]$ and satisfy a Lipschitz condition of order α on I . We show that the conclusion of one of his theorems on the absolute convergence of Fourier series remains valid when the function is merely of bounded r th variation in I and belongs to a class $\text{Lip}(\alpha, p)$ in I . Our results also generalize three theorems of S. M. Mazhar [4].

1. Introduction. Let

$$(1.1) \quad \sum_{k=1}^{\infty} (a_{n_k} \cos n_k x + b_{n_k} \sin n_k x)$$

be the Fourier series of a 2π -periodic function $f \in L[-\pi, \pi]$, where $\{n_k\}$ ($k \in N$) is a strictly increasing sequence of natural numbers such that

$$(1.2) \quad n_{k+1} - n_k \rightarrow \infty \quad \text{as } k \rightarrow \infty.$$

P. B. Kennedy [3] proved the following theorem which recalls a well-known result of Zygmund [7, p. 136].

THEOREM A [3, THEOREM V(iv)]. *If*

- (i) $\{n_k\}$ satisfies (1.2),
- (ii) f is of bounded variation in a subinterval $I = [x_0 - \delta, x_0 + \delta]$ of $[-\pi, \pi]$, and
- (iii) $f \in \text{Lip } \alpha$ in I , where $0 < \alpha < 1$, then

$$(1.3) \quad \sum_{k=1}^{\infty} (|a_{n_k}| + |b_{n_k}|) < \infty.$$

In [4, Theorem 1] S. M. Mazhar generalized a theorem of Noble [5], obtaining a variant of Theorem A in which (i) is replaced by a stronger condition, while (ii) is replaced by the weaker condition that f is of bounded r th variation. In this paper we prove a generalization of both theorems. The classes of functions we consider are defined below.

Received by the editors September 20, 1983 and, in revised form, March 20, 1984.

1980 *Mathematics Subject Classification*. Primary 42A28.

Key words and phrases. Lacunary Fourier series, absolute convergence, Lipschitz condition, bounded r th variation.

©1985 American Mathematical Society
0002-9939/85 \$1.00 + \$.25 per page

2. Definitions. (i) For $0 < \alpha \leq 1$ and $p \geq 1$, $\text{Lip}(\alpha, p, I)$ is defined to be the class of functions f in $L[-\pi, \pi]$ such that

$$\left(\int_I |f(x+h) - f(x)|^p dx \right)^{1/p} = O(|h|^\alpha) \quad \text{as } h \rightarrow 0.$$

Evidently $\text{Lip } \alpha \subseteq \text{Lip}(\alpha, p)$ for every $p \geq 1$.

(ii) Suppose a function f is defined on an interval $[a, b]$ where $a \leq x_0 < x_1 < \dots < x_{n-1} < x_n \leq b$. We write, as usual,

$$\Delta^1 f(x_i) = f(x_{i+1}) - f(x_i), \quad \Delta^q f(x_i) = \Delta^{q-1} f(x_{i+1}) - \Delta^{q-1} f(x_i) \quad (q \geq 2),$$

so that

$$\Delta^q f(x_i) = \sum_{\nu=0}^q (-1)^\nu \binom{q}{\nu} f(x_{i+q-\nu}).$$

Then f is said to be of bounded r th variation on $[a, b]$ (where r is a positive integer) if there is a constant M such that

$$\sum_{i=0}^{n-r} |\Delta^r f(x_i)| \leq M$$

whenever $x_0, x_1, \dots, x_n$ are in arithmetical progression.

It is clear that every function of bounded variation is of bounded r th variation, but the converse is false, as is pointed out by Mazhar [4].

We prove the following theorems.

3. Results.

THEOREM 1. *If*

- (i) $\{n_k\}$ satisfies (1.2),
- (ii) f is of bounded r th variation in I , and
- (iii) $f \in \text{Lip}(\alpha, p, I)$, where $0 < \alpha \leq 1, p > 2, \alpha p > 1$, then (1.3) holds.

This theorem clearly generalizes the results of Kennedy and Mazhar referred to in §1. It may also be compared to a theorem of Hardy and Littlewood [2] in which $r = 1, I = [-\pi, \pi]$, but lacunarity is not postulated.

The following extensions of Theorem 1 can also be proved; they contain Theorems 2 and 3 of [4].

THEOREM 2. *If*

- (i) $\{n_k\}$ satisfies (1.2),
- (ii) f satisfies Theorem 1(ii), and
- (iii) $f \in \text{Lip}(\alpha, p, I)$, where $0 < \alpha \leq 1, p > 2$, then

$$\sum_{k=1}^{\infty} \left(|a_{n_k}|^\beta + |b_{n_k}|^\beta \right) < \infty$$

for every β satisfying $2 > \beta > 2(p-1)/(2p + \alpha p - 3)$.

THEOREM 3. *Under the hypothesis of Theorem 2,*

$$\sum_{k=1}^{\infty} n_k^{\beta/2} (|a_{n_k}| + |b_{n_k}|) < \infty$$

for $\beta < (\alpha p - 1)/2(p - 1)$.

Observe that, for $\beta = 1$, Theorem 2 reduces to Theorem 1, and for $\beta = 0$, Theorem 3 also reduces to Theorem 1.

4. Proof of Theorem 1. We need the following lemmas. Lemma 1 is a special case of a very general theorem due to Paley and Wiener [6, Theorem XLII']. Lemma 2 is due to Kennedy [3, Lemma 1]. The result of Lemma 3 is given in the introduction of the paper [1] by Hardy and Littlewood.

LEMMA 1. *Suppose the function $f \in L[-\pi, \pi]$ has Fourier series (1.1). If $n_{k+1} - n_k \rightarrow \infty$ as $k \rightarrow \infty$ and $f \in L^2(I)$ for some subinterval I of $[-\pi, \pi]$, then $f \in L^2[-\pi, \pi]$.*

LEMMA 2. (i) *Let $\{\lambda_k\}$ ($-\infty < k < \infty$) be a sequence of real numbers satisfying $\lambda_{k+1} - \lambda_k \rightarrow \infty$ ($|k| \rightarrow \infty$), $\lambda_{k+1} - \lambda_k > 8\pi\delta^{-1}$ ($\delta > 0$).*

(ii) *Let $\{A_k\}$ ($-\infty < k < \infty$) be a sequence of complex numbers such that $\sum_{-\infty}^{\infty} |A_k| s^{|\lambda_k|} < \infty$ ($0 < s < 1$).*

(iii) *Let*

$$\phi(s, x) = \sum_{-\infty}^{\infty} A_k s^{|\lambda_k|} e^{i\lambda_k x} \quad (0 < s < 1, -\infty < x < \infty),$$

(iv) *Let*

$$\phi(x) = L^2\text{-}\lim_{s \rightarrow 1} \phi(s, x) \quad \text{in } I = [x_0 - \delta, x_0 + \delta].$$

Then

$$\sum_{-\infty}^{\infty} |A_k|^2 \leq 8\delta^{-1} \int_I |\phi(x)|^2 dx.$$

LEMMA 3. *If $f \in \text{Lip}(\alpha, p, I)$, $p \geq 1$, then $f \in L^p(I)$.*

Putting $n_0 = 0$, $n_k = -n_{-k}$ ($k < 0$), $C_{n_0} = 0$, $C_{n_k} = \frac{1}{2}(a_{n_k} - ib_{n_k})$ ($k > 0$), and $C_{n_k} = \bar{C}_{n_{-k}}$ ($k < 0$), we write (1.1) in the form

$$\sum_{k=-\infty}^{\infty} C_{n_k} e^{in_k x}.$$

Since $C_{n_k} \rightarrow 0$ as $k \rightarrow \infty$,

$$(4.1) \quad \sum_{k=-\infty}^{\infty} |C_{n_k}| s^{|\lambda_k|} < \infty \quad (0 < s < 1).$$

Put

$$f(s, x) = \sum_{k=-\infty}^{\infty} C_{n_k} e^{in_k x} s^{|\lambda_k|} \quad (0 < s < 1, -\infty < x < \infty).$$

Now, by Theorem 1(iii), $f \in \text{Lip}(\alpha, 2, I)$ and, therefore, by Lemma 3, $f \in L^2(I)$. Hence, by Lemma 1, $f \in L^2[-\pi, \pi]$, and so, by a known theorem [7, p. 87], it follows that

$$(4.2) \quad f(x) = L^2\text{-}\lim_{s \rightarrow 1} f(s, x) \quad (|x| \leq \pi).$$

Take integers K, j such that

$$(4.3) \quad n_K > 2r\pi/\delta \quad \text{and} \quad 0 \leq j \leq \delta n_K/8\pi.$$

In the following argument we let r be an even integer; the treatment for odd r is similar.

Put

$$(4.4) \quad g_j(x) = \sum_{\nu=0}^r (-1)^\nu \binom{r}{\nu} f\left(x + \frac{2j\pi}{n_K} + \frac{(r-2\nu)\pi}{2n_K}\right)$$

and

$$A_{kj} = C_{n_k} \left(2i \sin \frac{n_k \pi}{2n_K} \right)^r \exp\left(\frac{in_k 2j\pi}{n_K} \right).$$

Since $|A_{kj}| \leq |C_{n_k}| 2^r$, it follows from (4.1) that

$$(4.5) \quad \sum_{k=-\infty}^{\infty} |A_{kj}| s^{|n_k|} < \infty \quad (0 < s < 1).$$

If

$$(4.6) \quad g_j(s, x) = \sum_{k=-\infty}^{\infty} A_{kj} s^{|n_k|} e^{in_k x} \quad (0 < s < 1, -\infty < x < \infty),$$

we have

$$g_j(s, x) = \sum_{\nu=0}^r (-1)^\nu \binom{r}{\nu} f\left(s, x + \frac{2j\pi}{n_K} + \frac{(r-2\nu)\pi}{2n_K}\right);$$

from which, together with (4.2) and (4.4), we obtain

$$(4.7) \quad g_j(x) = L^2\text{-}\lim_{s \rightarrow 1} g_j(s, x) \quad \text{in } [x_0 - \tfrac{1}{2}\delta, x_0 + \tfrac{1}{2}\delta].$$

We may assume that

$$(4.8) \quad n_{k+1} - n_k > 16\pi\delta^{-1} \quad \text{for all } k.$$

In view of (1.2), this can be achieved, if necessary, by adding to $f(x)$ a polynomial in $\exp(in_k x)$, a process which affects neither the hypothesis nor the conclusion of the theorem. Hence, putting $I_1 = [x_0 - \tfrac{1}{2}\delta, x_0 + \tfrac{1}{2}\delta]$, we obtain from (4.5)–(4.8) and Lemma 2 that

$$(4.9) \quad \sum_{k=-\infty}^{\infty} |A_{kj}|^2 \leq 16\delta^{-1} \int_{I_1} |g_j(x)|^2 dx.$$

But

$$|A_{kj}| = |C_{n_k}| 2^r \left(\sin \frac{n_k \pi}{2n_K} \right)^r.$$

Therefore (4.9) gives

$$(4.10) \quad \sum_{n_K/2}^{n_K} 2^{2r} \left(\sin \frac{n_K \pi}{2n_K} \right)^{2r} |C_{n_K}|^2 \leq 16\delta^{-1} \int_{I_1} |g_j(x)|^2 dx.$$

But if $\frac{1}{2}n_K \leq |n_K| \leq n_K$, then

$$\left(\sin \frac{\pi n_K}{2n_K} \right)^{2r} \geq \left(\frac{1}{2} \right)^r.$$

Therefore (4.10) reduces to

$$(4.11) \quad \sum_{n_K/2}^{n_K} |C_{n_K}|^2 \leq \frac{16\delta^{-1}}{2^r} \int_{I_1} |g_j(x)|^2 dx.$$

Let $p' = p - 1$. Since $p > 2$, $p' > 1$. Define q by the equation $1/p' + 1/q = 1$.

We now denote, by $\sum_j$, summation over the range $0 \leq j \leq [\delta n_K / 8\pi]$ (where $[a]$ represents the integral part of a). It then follows from (4.11) that

$$(4.12) \quad \left\{ \sum_{n_K/2}^{n_K} |C_{n_K}|^2 \right\}^q \leq \frac{(16\delta^{-1}2^{-r})^q}{[\delta n_K / 8\pi]} \sum_j \left\{ \int_{I_1} |g_j(x)|^2 dx \right\}^q \\ \leq \frac{C}{n_K} \sum_j \left\{ \int_{I_1} |g_j(x)|^2 dx \right\}^q,$$

where the constant C depends on δ and r . C will henceforth denote positive constants not necessarily the same at each occurrence.

Since $2 = (p' + 1)/p' + 1/q$, an application of Hölder's inequality gives

$$\left\{ \int_{I_1} |g_j(x)|^2 dx \right\}^q = \left\{ \int_{I_1} |g_j(x)|^{(p'+1)/p'+1/q} dx \right\}^q \\ \leq \left[\left(\int_{I_1} |g_j(x)|^{(p'+1)p'/p'} dx \right)^{1/p'} \left(\int_{I_1} |g_j(x)|^{1/q \cdot q} dx \right)^{1/q} \right]^q \\ = \left\{ \int_{I_1} |g_j(x)|^p dx \right\}^{q/p'} \left\{ \int_{I_1} |g_j(x)| dx \right\}.$$

This, combined with (4.12), gives

$$(4.13) \quad \left\{ \sum_{n_K/2}^{n_K} |C_{n_K}|^2 \right\}^q \leq \frac{C}{n_K} \sum_j \left\{ \left(\int_{I_1} |g_j(x)|^p dx \right)^{q/p'} \int_{I_1} |g_j(x)| dx \right\}.$$

Putting $y = x + 2j\pi/n_K + r\pi/2n_K$ in (4.4), we evidently have

$$g_j(x) = \sum_{\nu=0}^{r-1} a_\nu \left\{ f\left(y - \frac{\nu\pi}{n_K}\right) - f\left(y - \frac{(\nu+1)\pi}{n_K}\right) \right\},$$

where $a_0, a_1, \dots, a_{r-1}$ are constants. Hence, by Minkowski's inequality and condition Theorem 1(iii),

$$\left[\int_{I_1} |g_j(x)|^p dx \right]^{1/p} \leq \sum_{\nu=0}^{r-1} |a_\nu| \left\{ \int_{I_1} \left| f\left(y - \frac{\nu\pi}{n_K}\right) - f\left(y - \frac{(\nu+1)\pi}{n_K}\right) \right|^p dx \right\}^{1/p} \\ = O(n_K^{-\alpha}).$$

Also, for $|x - x_0| \leq \delta/2$ and integers j and K satisfying (4.3), the points $y - \nu\pi/n_K$ ($\nu = 0, 1, \dots, r$) all lie in $[x_0 - \delta, x_0 + \delta]$. Therefore, if V is the total r th variation of f in I , it follows from Theorem 1(ii) that

$$\sum_j |g_j(x)| \leq V \quad \text{for } |x - x_0| \leq \frac{1}{2}\delta.$$

Hence, by (4.13),

$$\left\{ \sum_{n_K/2}^{n_K} |C_{n_k}|^2 \right\}^q \leq C n_K^{-1} n_K^{-\alpha p q / p'} V \delta = C n_K^{-1 - \alpha p q / p'},$$

which in turn gives

$$(4.14) \quad \sum_{n_K/2}^{n_K} |C_{n_k}|^2 \leq C n_K^{-1/q - \alpha p / p'}.$$

Let m be a positive integer. Either the set of integers k for which $2^m < |n_k| \leq 2^{m+1}$ is empty, or there is a member of this set, say $K = K(m)$, which has largest modulus; and in the latter case, the set is included in the set of k for which $\frac{1}{2}n_K \leq |n_k| \leq n_K$. Moreover, if m is large enough, K satisfies (4.3). Hence, in either case it follows from (4.14) that

$$\begin{aligned} \sum_{2^m < |n_k| \leq 2^{m+1}} \left\{ |C_{n_k}|^2 \right\} &\leq \sum_{n_K/2 \leq |n_k| \leq n_K} \left\{ |C_{n_k}|^2 \right\} = O(n_K^{-1/q - \alpha p / p'}) \\ &= O(2^{m(-1/q - \alpha p / p')}). \end{aligned}$$

Also, by (4.8), the number of terms in the above summation is $O(2^m)$. Hence, by Cauchy's inequality,

$$\begin{aligned} \sum_{2^m < |n_k| \leq 2^{m+1}} |C_{n_k}| &\leq \left\{ \sum_{2^m < |n_k| \leq 2^{m+1}} |C_{n_k}|^2 \right\}^{1/2} \left\{ \sum_{2^m < |n_k| \leq 2^{m+1}} 1 \right\}^{1/2} \\ &= O\{2^{m/2((1-\alpha p)/p')}\}, \end{aligned}$$

so

$$(4.15) \quad \sum_{k=-\infty}^{\infty} |C_{n_k}| \leq C \sum_{m=1}^{\infty} 2^{m(1-\alpha p)/2p'}.$$

Since $\alpha p > 1$, the series on the right side is convergent. This completes the proof of Theorem 1.

Theorems 2 and 3 can be proved in a similar manner.

Thanks are due to the referee for his useful comments.

REFERENCES

1. G. H. Hardy and J. E. Littlewood, *Some properties of fractional integrals*. I, Math. Z. **27** (1928), 565–606.
2. ———, *Notes on the theory of series*. IX: *On the absolute convergence of Fourier series*, J. London Math. Soc. **3** (1928), 250–253.
3. P. B. Kennedy, *Fourier series with gaps*, Quart. J. Math. Oxford Ser. (2) **7** (1956), 224–230.

4. S. M. Mazhar, *On the absolute convergence of a Fourier series with gaps*, Proc. Nat. Acad. Sci. India Sect. A **26** (1960), 104–109.
5. M. E. Noble, *Coefficient properties of Fourier series with a gap condition*, Math. Ann. **128** (1954), 55–62.
6. R. E. A. C. Paley and N. Wiener, *Fourier transforms in the complex domain*, Amer. Math. Soc. Colloq. Publ., Vol. 19, Amer. Math. Soc., Providence, R. I., 1934.
7. A. Zygmund, *Trigonometrical series*, PWN, Warsaw, 1935.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, THE MAHARAJA SAYAJIRAO UNIVERSITY OF BARODA, BARODA - 390 002, INDIA