MAXIMAL SETS OF ORTHOGONAL MEASURES ARE NOT ANALYTIC

DAVID PREISS AND JAN RATAJ

ABSTRACT. By proving the theorem given in the title we answer a question posed by D. Mauldin at the conference on Measure Theory held in Oberwolfach in 1981.

In this note we study maximal sets of pairwise orthogonal probability measures on $\langle 0, 1 \rangle$. The existence of such sets is guaranteed by the Kuratowski-Zorn principle. Our theorem says that such sets cannot be Borel in the weak topology, or even analytic. This answers a problem posed by D. Mauldin at the Conference on Measure Theory held in Oberwolfach in 1981 (see [2]). To study maximal sets of pairwise orthogonal measures is useful in the light of general properties of orthogonal kernels; see [2] for a survey of these results and further references.

First let us introduce some notation. Assume that K is a compact metric space. $\mathcal{M}(K)$ denotes the locally convex space of all finite Borel signed measures on K with weak topology. $\mathcal{M}^+(K)$ denotes its subspace of all positive measures and $\mathcal{P}(K)$ its subspace of all probability measures. It is well known that $\mathcal{P}(K)$ is a compact metrizable space.

If $x \in K$ we denote by ε_x the Dirac probability measure concentrated at x . Let $\mathcal{D}(K)$ denote the set of all finite convex combinations of Dirac probabilities on K . $\mathcal{D}(K)$ is a dense subset of $\mathcal{P}(K)$.

We will also use the norm

$$\|\mu\| = \sup\{|\mu(A)| : A \text{ is a Borel subset of } K\} \quad \text{for } \mu \in \mathcal{M}(K).$$

The function $\mu \mapsto \|\mu\|$ is lower semicontinuous on $\mathcal{M}(K)$. Consequently, the function $(\mu, \nu) \mapsto \|\mu - \nu\|$ is lower semicontinuous on $\mathcal{P}(K) \times \mathcal{P}(K)$.

If L is a compact subspace of K and $B \subseteq \mathcal{P}(K)$, we define the set of restrictions $B \upharpoonright L = \{\mu \upharpoonright L : \mu \in B\}$.

Now consider the special case $K = \langle 0, 1 \rangle$. Assume that $r \in \mathbb{N}$, $I_1, \dots, I_r$ are disjoint compact intervals in $\langle 0, 1 \rangle$; by an interval we always mean a nondegenerate interval, further, that $\lambda_1, \dots, \lambda_r$ are positive numbers such that $\sum_{i=1}^r \lambda_i = 1$ and

Received by the editors June 17, 1983 and, in revised form, August 22, 1983.
1980 *Mathematics Subject Classification*. Primary 28A33; Secondary 60B05.

$\Delta > 0$. Then we put

$$(1) \quad U(r; I_1, \dots, I_r; \lambda_1, \dots, \lambda_r; \Delta) \\ = \{ \mu \in \mathcal{P}(\langle 0, 1 \rangle) : \mu(\text{int } I_i) > \lambda_i - \Delta, \mu(I_i) \leq \lambda_i + \Delta, i = 1, \dots, r \}.$$

If $\mu \in \mathcal{D}(\langle 0, 1 \rangle)$, $\mu = \sum_{i=1}^r \lambda_i \varepsilon_{x_i}$, then the system of all sets

$$U(r; I_1, \dots, I_r; \lambda_1, \dots, \lambda_r; \Delta)$$

such that $x_i \in \text{int } I_i$, $i = 1, \dots, r$, and $\Delta > 0$ forms a basis of open neighbourhoods of μ . Consequently, every open set in $\mathcal{P}(\langle 0, 1 \rangle)$ contains an open set of the type (1).

For the proof we need the Banach-Mazur game as an instrument for proving residuality (see [3]). Assume that X is a complete metric space and $M \subseteq X$. The Banach-Mazur game is played by two players, (A) and (B). In the first step, (A) puts $U_1 = X$ and (B) chooses an open set $U'_1 \subseteq U_1$. In the n th step (A) chooses an open set $U_n \subseteq U'_{n-1}$ and (B) chooses an open $U'_n \subseteq U_n$. This defines a nonincreasing sequence of open sets. If their intersection contains an element of M , player (A) wins, in the opposite case (B) wins. It is well known that (A) has a winning strategy in this game if and only if M is residual in X ; the proof in [3] easily extends for the general case.

Our main result is given in the following

THEOREM. *No maximal set of pairwise orthogonal Borel probability measures on the interval $\langle 0, 1 \rangle$ is analytic.*

For the proof we will use two auxiliary lemmas:

LEMMA 1. *Assume that $K \subseteq \langle 0, 1 \rangle$ is a finite union of compact disjoint intervals, $V \subseteq \mathcal{P}(\langle 0, 1 \rangle)$ an open set, and $\varepsilon \in (0, 1)$. Then the set $Y = \{ \zeta \in \mathcal{P}(K) : \varepsilon \cdot \zeta \in V \upharpoonright K \}$ is open in $\mathcal{P}(K)$.*

PROOF. Let ζ_0 be an arbitrary measure in Y and let $\mu_0 \in V$ be an extension of $\varepsilon \cdot \zeta_0$. Clearly, if $\zeta \in \mathcal{P}(K)$ is sufficiently close to ζ_0 then the measure μ , defined by $\mu \upharpoonright K = \varepsilon \cdot \zeta$, $\mu \upharpoonright \langle 0, 1 \rangle \setminus K = \mu_0 \upharpoonright \langle 0, 1 \rangle \setminus K$, extends $\varepsilon \cdot \zeta$ and is arbitrarily close to μ_0 .

LEMMA 2. *Assume that K is the compact set from Lemma 1, $V \subseteq \mathcal{P}(\langle 0, 1 \rangle)$ an open set, $T \subseteq V$ is dense and open in V , $U \subseteq \mathcal{P}(K)$ is an open set, $\varepsilon \in (0, 1)$, and $\Delta > 0$. Assume further that for every $\zeta \in U$ the measure $\varepsilon \cdot \zeta$ can be extended into V (or, equivalently $\varepsilon \cdot \zeta \in V \upharpoonright K$). Then the set $Y = \{ \zeta \in U : \text{there exists } t \in (\varepsilon - \Delta, \varepsilon + \Delta), \text{ such that } t \cdot \zeta \in T \upharpoonright K \}$, is open and dense in U .*

PROOF. From Lemma 1 we see that Y is open. We prove that it is also dense in U . Let U_0 be any nonempty open subset of U . There is $\zeta_0 \in U_0$ such that (i) $\zeta_0(\text{bdry } K) = 0$. By the assumption, $\varepsilon \cdot \zeta_0$ can be extended at some $\mu_0 \in V$. Let μ_n be any sequence of elements of T converging to μ_0 . By (i), $\mu_n \upharpoonright K \rightarrow \mu_0 \upharpoonright K = \varepsilon \cdot \zeta_0$. If $\zeta_n = (\mu_n(K))^{-1} \cdot \mu_n \upharpoonright K$, then $\zeta_n \rightarrow \zeta_0$. Since $\mu_n(K) \rightarrow \varepsilon$, $\zeta_n \in U_0 \cap Y$ for all sufficiently large n .

The next lemma is the main step of the proof of our theorem.

LEMMA 3. Assume that $K \subseteq \langle 0, 1 \rangle$ is the compact set from Lemma 1, $0 \subseteq \mathcal{P}(K)$ is a nonempty open set in $\mathcal{P}(K)$, $B \subseteq \mathcal{P}(\langle 0, 1 \rangle)$ is residual in an open nonempty set $G \subseteq \mathcal{P}(\langle 0, 1 \rangle)$ and $\varepsilon \in (0, 1)$. Assume further that

(1) there exists $t_0 > \varepsilon$ such that, for every $\zeta \in 0$, $t_0 \cdot \zeta \in G \upharpoonright K$.

Then the set $X = \{\zeta \in 0: \text{there is } t > \varepsilon \text{ such that } t \cdot \zeta \in B \upharpoonright K\}$ is residual in 0.

PROOF. There are dense, open subsets H_n in G such that $\bigcap_{n=1}^{\infty} H_n \subseteq B$.

Let us fix a number $0 < \delta < |t_0 - \varepsilon|$. We prove the residuality of the set X in 0 by describing a winning strategy in the Banach-Mazur game. We use the same notation as in the beginning of this text. Put $U_1 = 0$. Player (B) chooses open sets $U_i' \subseteq 0$ such that

(i) $U_i' \subseteq U_i$, $i = 1, 2, \dots$, and (A) chooses open sets $U_i \subseteq U_{i-1}'$, $i = 2, 3, \dots$. We will even require that

(ii) $\bar{U}_i \subseteq U_{i-1}$, $i = 2, 3, \dots$. We will describe the moves of (A), no matter how (B) played in the preceding steps. In addition, we will construct open sets $V_i \subseteq \mathcal{P}(\langle 0, 1 \rangle)$, $i = 1, 2, \dots$, and numbers $\varepsilon_i > 0$, $i = 1, 2, \dots$, with the following properties:

(iii) $\bar{V}_i \subseteq V_{i-1} \cap H_i$, $i = 2, 3, \dots$,

(iv) $|\varepsilon_i - \varepsilon_{i-1}| < \delta \cdot 2^{-i}$, $i = 2, 3, \dots$,

(v) if $\zeta \in U_i$, then $\varepsilon_i \cdot \zeta \in V_i \upharpoonright K$, $i = 1, 2, \dots$,

(vi) $\text{diam } V_i < i^{-1}$, $i = 2, 3, \dots$,

(vii) if $\mu \in V_i$ then $\mu(\text{bdry } K) < i^{-1}$, $i = 2, 3, \dots$.

CONSTRUCTION. We put $V_1 = G$ and $\varepsilon_1 = t_0$. Then, by the assumption (1) of this lemma, the objects U_1, V_1, ε_1 satisfy the condition (v). The remaining conditions have no relation to the case $i = 1$.

Assume that open sets $U_1, U_1', V_1, \dots, U_{n-1}, U_{n-1}', V_{n-1}$ and positive numbers $\varepsilon_1, \dots, \varepsilon_{n-1}$ have already been defined and that the requirements (i)–(vii) hold. By Lemma 2 the set $X_n = \{\zeta \in U_{n-1}': \text{there is } s \in (\varepsilon_{n-1} - \delta \cdot 2^{-n}, \varepsilon_{n-1} + \delta \cdot 2^{-n}) \text{ such that } s \cdot \zeta \in (V_{n-1} \cap H_n) \upharpoonright K\}$, is dense and open in U_{n-1}' . Take $\zeta_n \in X_n$ such that $\zeta_n(\text{bdry } K) = 0$ and take $\varepsilon_n \in (\varepsilon_{n-1} - \delta \cdot 2^{-n}, \varepsilon_{n-1} + \delta \cdot 2^{-n})$ such that $\varepsilon_n \cdot \zeta_n$ can be extended to some $\mu_n \in V_{n-1} \cap H_n$; requirement (iv) holds for ε_n . There exists an open set $V_n \subseteq \mathcal{P}(\langle 0, 1 \rangle)$ such that $\mu_n \in V_n$ and the requirements (iii), (vi), (vii) hold. By Lemma 1, if U_n is a sufficiently small neighbourhood of ζ_n , the requirements (ii), (v) also hold.

Finally we verify that our strategy is really winning. Let ζ be a measure from the intersection $\bigcap_{n=1}^{\infty} U_n$; we show that $\zeta \in X$. By (iv), the sequence ε_n converges to some $\bar{\varepsilon} > \varepsilon$. By (v), $\varepsilon_n \cdot \zeta$ can be extended to some $\mu_n \in V_n$, $n = 1, 2, \dots$, and, by (iii) and (vi), there exists a measure $\mu \in \bigcap V_n \subseteq \bigcap H_n \subseteq B$ such that $\mu_n \rightarrow \mu$. By (vii), $\mu(\text{bdry } K) = 0$, hence $\mu_n \upharpoonright K \rightarrow \mu \upharpoonright K$. On the other hand, the sequence $\mu_n \upharpoonright K = \varepsilon_n \cdot \zeta$ converges to $\bar{\varepsilon} \cdot \zeta$; consequently, $\mu \upharpoonright K = \bar{\varepsilon} \cdot \zeta$. In other words, $\mu \in B$ is an extension of $\bar{\varepsilon} \cdot \zeta$.

We now see that this strategy of the Banach-Mazur game with the set $X \subseteq 0$ is winning for (A); hence, by the Theorem mentioned in the beginning of this paper, X is residual in 0 .

LEMMA 4. Assume that $G \subseteq \mathcal{P}(\langle 0, 1 \rangle)$ is a nonempty open convex set. Let $G_1 \subseteq G$ and $G_2 \subseteq G$ be open nonempty sets such that $G \subseteq \bar{G}_1 \cup \bar{G}_2$, and let $\tau \in (0, 1)$. Then

- (a) there exists measures $\mu_0 \in \bar{G}_1 \cap G$ and $\nu_0 \in G_2$ such that $\tau \cdot \mu_0 \leq \nu_0$, and
- (b) there exists a compact $K \subseteq (\langle 0, 1 \rangle)$ which is a finite union of compact disjoint intervals and there exists a nonempty open subset $0 \subseteq \mathcal{P}(K)$ such that, for every $\zeta \in 0$, $\tau \cdot \zeta \in G_1 \upharpoonright K \cap G_2 \upharpoonright K$.

PROOF. (a) Let $\mu \in G_1$ and $\nu \in G_2$ be arbitrary measures. Put (i)

$$\xi_k = \tau^k \cdot \mu + (1 - \tau^k) \cdot \nu, \quad k = 0, 1, 2, \dots$$

Clearly $\xi_k \in G$ for every k , because ξ_k are convex combinations of μ and ν . Let $k_0 \geq 1$ be the least natural number such that $\xi_{k_0} \in G_2$; the existence of such a number follows from the fact that $\xi_k \rightarrow \nu \in G_2$. We denote $\mu_0 = \xi_{k_0-1}$ (clearly $\mu_0 \in \bar{G}_1 \cap G$) and $\nu_0 = \xi_{k_0}$. From (i) we get

$$\nu_0 = \tau^{k_0} \cdot \mu + (1 - \tau^{k_0}) \cdot \nu = \tau \cdot \mu_0 + (1 - \tau) \cdot \nu,$$

hence $\tau \cdot \mu_0 \leq \nu_0$.

(b) We use the measures μ_0, ν_0 from (a). There exists a sequence $\mu_n, n = 1, 2, \dots$, in $G_1 \cap \mathcal{D}(\langle 0, 1 \rangle)$ such that $\mu_n \rightarrow \mu_0$. Define $\nu_n = \tau \cdot \mu_n + (\nu_0 - \tau \cdot \mu_0)$, $n = 1, 2, \dots$. We see that $\nu_n \rightarrow \nu_0$, hence there exists n_0 such that $\nu_{n_0} \in G_2$. Let μ_{n_0} have the form $\mu_{n_0} = \sum_{i=1}^r \lambda_i \cdot \varepsilon_{x_i}$, where $r \in \mathbb{N}$, $x_i \in \langle 0, 1 \rangle$, $\lambda_i > 0$, $i = 1, \dots, r$, $x_i \neq x_j$ for $i \neq j$ and $\sum_{i=1}^r \lambda_i = 1$. Let $\tilde{\mu} = \mu_{n_0}$. The measure $\nu_0 - \tau \cdot \mu_0$ can be approximated by a combination of Dirac measures $\sum_{i=r+1}^s \xi_i \cdot \varepsilon_{x_i}$ (where $x_i \in \langle 0, 1 \rangle \setminus \{x_1, \dots, x_r\}$, $\xi_i > 0$, $i = r+1, \dots, s$, $x_i \neq x_j$ for $i \neq j$) such that $\tilde{\nu} = \tau \cdot \mu_0 + \sum_{i=r+1}^s \xi_i \cdot \varepsilon_{x_i} \in G_2$.

There exist compact disjoint intervals $I_i \subseteq \langle 0, 1 \rangle$, $x_i \in \text{int } I_i$, $i = 1, \dots, s$, and a number $\Delta > 0$ such that neighbourhoods

$$U(\tilde{\mu}) = U(r; I_1, \dots, I_r; \lambda_1, \dots, \lambda_r; \Delta)$$

and

$$U(\tilde{\nu}) = U(s; I_1, \dots, I_s; \tau \cdot \lambda_1, \dots, \tau \cdot \lambda_r, \xi_{r+1}, \dots, \xi_s; \tau \cdot \Delta)$$

fulfill the inclusions $U(\tilde{\mu}) \subseteq G_1$ and $U(\tilde{\nu}) \subseteq G_2$. Take compact intervals $K_i \subseteq \text{int } I_i$, $i = 1, \dots, r$. Put $K = K_1 \cup \dots \cup K_r$ and

$$0 = \{ \zeta \in \mathcal{P}(K) : \zeta(K_i) \in (\lambda_i - \Delta/2, \lambda_i + \Delta/2), i = 1, \dots, r \}.$$

We now see that for every $\zeta \in 0$ the measure $\tau \cdot \zeta$ can be extended into both neighbourhoods $U(\tilde{\mu})$, $U(\tilde{\nu})$, and hence into both sets G_1 and G_2 .

PROOF OF THE THEOREM. Let A be a maximal set of pairwise orthogonal Borel probability measures in $\mathcal{P}(\langle 0, 1 \rangle)$. Assume, to the contrary, that A is analytic.

For every $k \in \mathbb{N}$ let E_k denote the space of all k -element subsets of A . We equip E_k with its usual topology (generated, e.g., by the Hausdorff metric). Clearly, each E_k is a continuous image of an open subset of the Cartesian product A^k , hence it is analytic.

Whenever $k \in \mathbb{N}$ and $\varepsilon \in (0, 1)$, the set of all $(\mu, j) \in \mathcal{P}(\langle 0, 1 \rangle) \times E_k$ such that $\|\mu - \nu\| \leq \varepsilon$ for every $\nu \in j$ is closed in $\mathcal{P}(\langle 0, 1 \rangle) \times E_k$. It follows that for every analytic set $C \subseteq E_k$ the set $\{(\mu, j) \in \mathcal{P}(\langle 0, 1 \rangle) \times C; \|\mu - \nu\| < \varepsilon \text{ for every } \nu \in j\}$ is analytic and hence its first projection $H_{k,\varepsilon}(C)$ is also analytic. In addition, let $\tau \in (0, \varepsilon)$. Define $F_{k,\varepsilon}^\tau(C) = H_{k,\varepsilon-\tau}(C) - H_{k+1,\varepsilon}(E_{k+1})$ and $U_{k,\varepsilon}^\tau = F_{k,\varepsilon}^\tau(E_k)$. Then:

(1) Whenever $k \in \mathbb{N}$, $\varepsilon \in (0, 1)$, $\varepsilon > \tau > 0$ and $C \subseteq E_k$ is analytic, the set $F_{k,\varepsilon}^\tau(C)$ is a difference of two analytic sets and hence it has the property of Baire.

(2) If $C_1 \subseteq E_k$ and $C_2 \subseteq E_k$ are disjoint sets, $\mu_1 \in F_{k,\varepsilon}^\tau(C_1)$ and $\mu_2 \in F_{k,\varepsilon}^\tau(C_2)$, then $\|\mu_1 - \mu_2\| \geq \tau$. (Otherwise we could find $\{\nu_1, \dots, \nu_k\} \in C_1$ and $\{\bar{\nu}_1, \dots, \bar{\nu}_k\} \in C_2$ such that $\|\mu_1 - \nu_i\| < \varepsilon$ and $\|\mu_1 - \bar{\nu}_i\| \leq \|\mu_1 - \mu_2\| + \|\mu_2 - \bar{\nu}_i\| < \varepsilon$ for every $i = 1, \dots, k$. The disjointedness of C_1 and C_2 would imply that $\mu_1 \in H_{k+1,\varepsilon}(E_{k+1})$.)

(3) Since A is a maximal set of orthogonal measures, for every $\nu \in \mathcal{P}(\langle 0, 1 \rangle)$ the set $\{\mu \in A; \|\nu - \mu\| < 1\}$ is nonempty and for every $t \in (0, 1)$ the set $\{\mu \in A; \|\nu - \mu\| < t\}$ is finite. We find $\varepsilon \in (0, 1)$ such that $\|\nu - \mu\| < \varepsilon$ for some $\mu \in A$. Let $j = \{\mu \in A; \|\nu - \mu\| < \varepsilon\}$. Since j is finite, there is $\tau \in (0, \varepsilon)$ such that $\|\mu - \nu\| < \varepsilon - \tau$ for each $\mu \in j$. Thus $\nu \in U_{k,\varepsilon}^\tau$, where k is the number of elements of j . Consequently,

$$\mathcal{P}(\langle 0, 1 \rangle) = \bigcup_{\varepsilon \in (0, 1)} \bigcup_{k=1}^{\infty} \bigcup_{\tau \in (0, \varepsilon)} U_{k,\varepsilon}^\tau.$$

We now prove that $U_{k,\varepsilon}^\tau$ is of first category for every $k \in \mathbb{N}$, $\varepsilon \in (0, 1)$, $\varepsilon > \tau > 0$. Assume, to the contrary, that $U_{k,\varepsilon}^\tau$ is of second category. Then, by the property of Baire, it is residual in some nonempty open set (see [1]). We can assume that $U_{k,\varepsilon}^\tau$ is residual in a convex nonempty open set $G \subseteq \mathcal{P}(\langle 0, 1 \rangle)$. Denote by Z the maximal open set in E_k such that $F_{k,\varepsilon}^\tau(Z)$ is of first category in G . We see that $F = E_k \setminus Z$ has at least two different elements, because the set $F_{k,\varepsilon}^\tau(\{j\})$ is nowhere dense for every $j \in E_k$. Take a partition $F = F_1 \cup F_2$, where F_1 and F_2 are Borel subsets of F with nonempty interiors. Put $A_1 = F_{k,\varepsilon}^\tau(F_1)$, $A_2 = F_{k,\varepsilon}^\tau(F_2)$. Clearly, A_1 and A_2 are both of second category in G ; by (1) they have the property of Baire, and hence they are residual in some open sets. Denote by G_1, G_2 the maximal open subsets in G , such that A_1, A_2 are residual in G_1, G_2 , respectively. Clearly $G \subseteq \overline{G_1} \cup \overline{G_2}$. By Lemma 4 there exists a compact subset $K \subseteq \langle 0, 1 \rangle$ which is a finite union of compact disjoint intervals, and a nonempty open set $0 \subseteq \mathcal{P}(K)$ such that for every $\zeta \in 0$ the measure $(1 - \tau) \cdot \zeta$ can be extended into G_1 and G_2 . By Lemma 3, the sets

$$X_1 = \{\zeta \in 0: \text{there is } t_1 > 1 - \tau \text{ such that } t_1 \cdot \zeta \in A_1 \upharpoonright K\}$$

and

$$X_2 = \{\zeta \in 0: \text{there is } t_2 > 1 - \tau \text{ such that } t_2 \cdot \zeta \in A_2 \upharpoonright K\}$$

are residual in 0 . Hence their intersection is nonempty and there exist $\zeta \in 0$, $\mu_1 \in A_1$, $\mu_2 \in A_2$ and $t_1, t_2 > 1 - \tau$ such that $\mu_1 \upharpoonright K = t_1 \cdot \zeta$ and $\mu_2 \upharpoonright K = t_2 \cdot \zeta$. Hence $\|\mu_1 - \mu_2\| < \tau$. But by (2) we see that $\|\mu_1 - \mu_2\| \geq \tau$. This contradiction proves that $U_{k,\varepsilon}^\tau$ is of first category.

In (3) it suffices to take only countable unions, hence the whole space $\mathcal{P}(\langle 0, 1 \rangle)$ is of first category. This contradiction proves our Theorem.

REFERENCES

1. K. Kuratowski, *Topology* I, Academic Press, New York and London, 1968.
2. D. Mauldin, D. Preiss and H. v. Weizsäcker, *A survey of problems and results concerning orthogonal transition kernels*, Measure Theory (Proc. Conf., Oberwolfach, 1981), Lecture Notes in Math., Vol. 945, Springer-Verlag, New York, 1982.
3. J. C. Oxtoby, *Measure and category*, Springer-Verlag, New York, 1980.

FACULTY OF MATHEMATICS AND PHYSICS, CHARLES UNIVERSITY, SOKOLOVSKA 83, 186 000 PRAGUE 8,
CZECHOSLOVAKIA