HIGH ORDER COEFFICIENT ESTIMATES IN THE CLASS Σ

Y. J. LEUNG AND G. SCHÖBER

ABSTRACT. Estimates are given for the coefficients b_n of functions in the class Σ in terms of $\operatorname{Re}\{1 - b_1\}$. As a consequence, there is an explicit finite number λ such that $\operatorname{Re}\{\lambda b_1 - b_n\} \leq \lambda$.

Introduction. Let Σ be the class of functions $g(z) = z + \sum_{n=2}^{\infty} b_n z^{-n}$ that are analytic and univalent for $|z| > 1$. The nonvanishing subclass Σ' consists of those functions $g \in \Sigma$ for which $g(z) \neq 0$, $|z| > 1$. Denote by S the familiar class of univalent analytic functions $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ in the open unit disk. It is easy to see that f belongs to S if and only if $g(z) = 1/f(1/z)$ belongs to Σ' .

The function $k(z) = z + 1/z$ is extremal for many problems in Σ and is closely related to the Koebe function $K(z) = z(1 - z)^{-2}$ in S . Its transform

$$k(z^{(n+1)/2})^{2\wedge(n+1)} = z(1 + z^{-n-1})^{2\wedge(n+1)}$$

is extremal for the coefficient problem $\max_{\Sigma} \operatorname{Re} b_n$ for $n = 1$ and $n = 2$, but not for $n \geq 3$.

It was conjectured by W. E. Kirwan [3] that

$$(1) \quad \operatorname{Re}\{nb_1 - b_n\} \leq n \quad (n \geq 2)$$

is true for all functions in the class Σ . Equality always occurs for the function k . This conjecture is known (Garabedian and Schiffer [2]) to be true for $n = 2$ and $n = 3$.

In general, we shall consider the inequality

$$(2) \quad \operatorname{Re}\{\lambda b_1 - b_n\} \leq \lambda.$$

Our principal result is that this inequality holds in Σ for some finite positive λ depending on n , but not on g . If $\lambda' > \lambda$, then

$$\operatorname{Re}\{\lambda' b_1 - b_n\} = \operatorname{Re}\{\lambda b_1 - b_n\} + (\lambda' - \lambda) \operatorname{Re} b_1 \leq \lambda + (\lambda' - \lambda) = \lambda';$$

that is, if (2) holds for some λ , then it holds for all $\lambda' > \lambda$ and equality with λ' occurs only for translations of the function k . The problem then becomes one of finding the least value λ for which (2) is true. It follows from the work in [3] that $\lambda = n$ is the least value for which inequality (2) holds when $n = 2$ or $n = 3$. We refer the reader to the articles [6,7] for further facts related to conjecture (1) and problem (2).

Received by the editors August 7, 1984 and, in revised form, September 28, 1984.
 1980 *Mathematics Subject Classification*. Primary 30C50.

The inequality (2) is contained in the following theorem.

THEOREM. *If $g(z) = z + \sum_{n=0}^{\infty} b_n z^{-n}$ belongs to Σ and $n \geq 2$, then there exists a finite positive number λ such that*

$$(3) \quad |\operatorname{Re} b_n| \leq \lambda \operatorname{Re}\{1 - b_1\}.$$

In particular, we have $\operatorname{Re}\{\lambda b_1 - b_n\} \leq \lambda$.

The number λ is given explicitly in formula (7). Afterwards, there is a section containing the numbers λ for $n = 2, 3, 4, 5$ and also simple estimates for λ that depend on n .

Note that (3) may be interpreted as coefficient estimates in a neighborhood of the function k . A generalization is stated as a concluding remark.

Proof of the theorem. Our proof is similar, but not identical, to that of R. N. Pederson [5] for a corresponding result in the class S . In addition, we shall keep track of the constants involved.

To estimate the coefficients of $g \in \Sigma$, we can assume without loss of generality that g is a slit mapping since such mappings are dense in Σ . If w_0 is any value omitted by g , then $g - w_0$ belongs to Σ' and we may associate with its reciprocal a function $f \in S$, as in the introduction. The function f may be embedded (cf. [1, Chapter 3]) into a Loewner chain $f(z, t)$ which satisfies a differential equation of the form

$$\partial f / \partial t = z[(1 + \kappa z)/(1 - \kappa z)] \partial f / \partial z, \quad t > 0,$$

where κ is a piecewise continuous function of t , $|\kappa| = 1$, and $f(z, 0) = f(z)$. In terms of $g(z, t) = 1/f(z, t)$ the differential equation has the similar form

$$(4) \quad \partial g / \partial t = z[(1 + \bar{\kappa} z)/(1 - \bar{\kappa} z)] \partial g / \partial z, \quad t > 0,$$

where $g(z, 0) = g(z) - w_0$.

From (4) the coefficients of $g(z, t) = e^{-t}[z + \sum_{n=0}^{\infty} b_n(t)z^{-n}]$ satisfy the differential equation

$b'_n(t) = (n+1)b_n(t) + 2(n-1)\kappa(t)b_{n-1}(t) + \cdots + 2\kappa^{n-1}(t)b_1(t) - 2\kappa^{n+1}(t)$, at least for $n \geq 1$. Repeated integration gives an iterated integral representation for the coefficients $b_n = b_n(0)$, $n \geq 1$, of $g - w_0$, hence of g , in the following form:

$$(5) \quad b_n = \Sigma(-1)^{m+1} 2^m \Gamma_{j_1 j_2 \cdots j_m} \int_0^\infty \int_{t_1}^\infty \cdots \int_{t_{m-1}}^\infty \kappa_1^{j_1} \kappa_2^{j_2} \cdots \kappa_m^{j_m} dt_m \cdots dt_2 dt_1,$$

where $\kappa_j = \kappa(t_j)e^{-t_j}$, $\Gamma_{j_1} = 1$, $\Gamma_{j_1 j_2 \cdots j_m} = (j_2 + \cdots + j_m - 1)(j_3 + \cdots + j_m - 1) \cdots (j_m - 1)$, and the summation is over all positive integers $j_1, j_2, \dots, j_m$ with $j_m \geq 2$ and $j_1 + j_2 + \cdots + j_m = n + 1$. Of course, b_1 is simply $2 \int_0^\infty \kappa(t)^2 e^{-2t} dt$. The coefficient $b_1 = 1$ if and only if $\kappa^2 \equiv 1$.

LEMMA 1. *For $j = 1, 2, \dots, N$, let θ_j be real and let δ_j be an integer multiple of π . Then we have the sharp inequality*

$$(6) \quad \begin{aligned} & |\cos(\theta_1 + \theta_2 + \cdots + \theta_N) - \cos(\delta_1 + \delta_2 + \cdots + \delta_N)| \\ & \leq N \left[\sum_{j=1}^N |\cos \theta_j - \cos \delta_j| \right]. \end{aligned}$$

PROOF. If $N = 1$, the inequality is trivial; assume therefore that $N \geq 2$. Let

$$\varphi(\theta_1, \dots, \theta_N) = N^2 - N(\cos \theta_1 + \dots + \cos \theta_N) + \cos(\theta_1 + \dots + \theta_N) - 1.$$

We shall show that $\varphi \geq 0$. If some θ_j satisfies $\cos \theta_j \leq 0$, then

$$\varphi \geq N^2 - N(N-1) - 1 - 1 = N - 2 \geq 0.$$

Thus we may restrict the domain of φ to the cube $|\theta_j| < \pi/2$, $1 \leq j \leq N$. The critical points of φ are found by setting

$$\partial \varphi / \partial \theta_j = N \sin \theta_j - \sin(\theta_1 + \dots + \theta_N) = 0 \quad \text{for } j = 1, 2, \dots, N.$$

That is, $\sin \theta_j$ has the value $(1/N)\sin(\theta_1 + \dots + \theta_N)$ for all j . Since the interval for θ_j is restricted to $(-\pi/2, \pi/2)$, we have $\theta_1 = \dots = \theta_N$. Thus at a critical point

$$\varphi = N^2 - N^2 \cos \theta_1 + \cos(N\theta_1) - 1.$$

This is nonnegative because of the inequality $1 - \cos(N\theta) \leq N(1 - \cos \theta)$. We conclude that $\varphi(\theta_1, \dots, \theta_N) \geq 0$ at all points. This proves (6) in case all $\delta_j = 0$.

The general case of (6) follows from this special case by replacing each θ_j by $\theta_j + \delta_j$ since $|\cos(\theta + \delta) - 1| = |\cos \theta - \cos \delta|$ whenever δ is an integer multiple of π .

Both sides of (6) vanish when all θ_j and δ_j are zero. To see that the constant N on the right side cannot be improved, choose all δ_j to be zero, set all θ_j equal to ε , and send ε to zero. $\square$

LEMMA 2. For each real-valued function θ , there exists a function δ whose values are either 0 or π such that

$$2|\cos \theta - \cos \delta| \leq [1 - \cos(2\theta)].$$

PROOF. Choose δ so that $\cos \delta = (\cos \theta)/|\cos \theta|$ if $\cos \theta \neq 0$ and $\cos \delta = 1$ if $\cos \theta = 0$. $\square$

We represent $\kappa(t) = e^{i\theta(t)}$ in the differential equation (4) and choose $\delta(t)$ from now on as in Lemma 2. Then

$$4 \int_0^\infty |\cos \theta(t) - \cos \delta(t)| e^{-2t} dt \leq 2 \int_0^\infty [1 - \cos 2\theta(t)] e^{-2t} dt = \operatorname{Re}\{1 - b_1\}.$$

We will also use the identity

$$2 \int_0^\infty |\cos 2\theta(t) - \cos 2\delta(t)| e^{-2t} dt = 2 \int_0^\infty [1 - \cos 2\theta(t)] e^{-2t} dt = \operatorname{Re}\{1 - b_1\}.$$

The choice $\theta = \delta$ in (5) gives coefficients $b_n = 0$ for $n \geq 2$. In general, therefore, we have

$$\begin{aligned} \operatorname{Re} b_n &= \sum (-1)^{m+1} 2^m \Gamma_{j_1 j_2 \dots j_m} \\ &\quad \times \int_0^\infty \int_{t_1}^\infty \dots \int_{t_{m-1}}^\infty \left[\cos \left(\sum_{k=1}^m j_k \theta(t_k) \right) - \cos \left(\sum_{k=1}^m j_k \delta(t_k) \right) \right] \\ &\quad \times e^{-\sum_{k=1}^m j_k t_k} dt_m \dots dt_2 dt_1 \end{aligned}$$

for $n \geq 2$. Let I denote just the iterated integral. Then Lemma 1 implies

$$\begin{aligned} |I| &\leq m \sum_{k=1}^m \int_0^\infty \int_{t_1}^\infty \cdots \int_{t_{m-1}}^\infty |\cos(j_k \theta(t_k)) - \cos(j_k \delta(t_k))| e^{-\sum_{i=1}^m j_i t_i} dt_m \cdots dt_2 dt_1 \\ &= m \sum_{k=1}^m I_k, \end{aligned}$$

where

$$\begin{aligned} I_k &= \int_0^\infty \int_{t_1}^\infty \cdots \int_{t_{k-1}}^\infty |\cos(j_k \theta(t_k)) - \cos(j_k \delta(t_k))| \int_{t_k}^\infty \cdots \int_{t_{m-1}}^\infty \\ &\quad \times e^{-\sum_{i=k+1}^m j_i t_i} dt_m \cdots dt_1 \\ &= \int_0^\infty \int_{t_1}^\infty \cdots e^{-\sum_{i=1}^{k-1} j_i t_i} \int_{t_{k-1}}^\infty |\cos(j_k \theta(t_k)) - \cos(j_k \delta(t_k))| \\ &\quad \times e^{-(j_k + \cdots + j_m)t_k} dt_k \cdots dt_1 / B_{j_1 j_2 \cdots j_m}(k) \end{aligned}$$

and

$$B_{j_1 j_2 \cdots j_m}(m) = 1, \quad B_{j_1 j_2 \cdots j_m}(k) = (j_{k+1} + \cdots + j_m)(j_{k+2} + \cdots + j_m) \cdots j_m.$$

If j_k is even, then from $t_k \geq t_{k-1}$ and Lemma 1 we have

$$\begin{aligned} &\int_{t_{k-1}}^\infty |\cos(j_k \theta(t_k)) - \cos(j_k \delta(t_k))| e^{-(j_k + \cdots + j_m)t_k} dt_k \\ &\leq e^{-(j_k + \cdots + j_m - 2)t_{k-1}} \int_0^\infty |\cos(j_k \theta(t_k)) - \cos(j_k \delta(t_k))| e^{-2t_k} dt_k \\ &\leq e^{-(j_k + \cdots + j_m - 2)t_{k-1}} (j_k/2)^2 \int_0^\infty |\cos(2\theta(t_k)) - \cos(2\delta(t_k))| e^{-2t_k} dt_k \\ &= e^{-(j_k + \cdots + j_m - 2)t_{k-1}} (j_k^2/8) \operatorname{Re}\{1 - b_1\}. \end{aligned}$$

Similarly, if j_k is odd, then

$$\begin{aligned} &\int_{t_{k-1}}^\infty |\cos(j_k \theta(t_k)) - \cos(j_k \delta(t_k))| e^{-(j_k + \cdots + j_m)t_k} dt_k \\ &\leq e^{-(j_k + \cdots + j_m - 2)t_{k-1}} \int_0^\infty |\cos(j_k \theta(t_k)) - \cos(j_k \delta(t_k))| e^{-2t_k} dt_k \\ &\leq e^{-(j_k + \cdots + j_m - 2)t_{k-1}} ((j_k + 1)/2) \\ &\quad \times \left\{ ((j_k - 1)/2) \int_0^\infty |\cos(2\theta(t_k)) - \cos(2\delta(t_k))| e^{-2t_k} dt_k \right. \\ &\quad \left. + \int_0^\infty |\cos \theta(t_k) - \cos \delta(t_k)| e^{-2t_k} dt_k \right\} \\ &\leq e^{-(j_k + \cdots + j_m - 2)t_{k-1}} ((j_k + 1)/2) \\ &\quad \times \{ ((j_k - 1)/4) \operatorname{Re}\{1 - b_1\} + (1/4) \operatorname{Re}\{1 - b_1\} \} \\ &= e^{-(j_k + \cdots + j_m - 2)t_{k-1}} (j_k(j_k + 1)/8) \operatorname{Re}\{1 - b_1\}. \end{aligned}$$

If $[x]$ denotes the greatest integer less than or equal to x , then we can combine these two cases by writing

$$\int_{t_{k-1}}^{\infty} |\cos(j_k \theta(t_k)) - \cos(j_k \delta(t_k))| e^{-(j_k + \dots + j_m)t_k} dt_k \\ \leq e^{-(j_k + \dots + j_m - 2)t_{k-1}} (j_k/4) [(j_k + 1)/2] \operatorname{Re}\{1 - b_1\}.$$

It follows that

$$I_k \leq \operatorname{Re}\{1 - b_1\} (j_k/4) [(j_k + 1)/2] \int_0^{\infty} \dots \int_{t_{k-2}}^{\infty} e^{-\sum_{i=1}^k j_i t_i} \\ \times e^{-(j_k + \dots + j_m - 2)t_{k-1}} dt_{k-1} \dots dt_1 / B_{j_1 j_2 \dots j_m}(k) \\ = \operatorname{Re}\{1 - b_1\} (j_k/4) [(j_k + 1)/2] / C_{j_1 j_2 \dots j_m}(k),$$

where

$$C_{j_1 j_2 \dots j_m}(k) = A_{j_1 j_2 \dots j_m}(k) B_{j_1 j_2 \dots j_m}(k)$$

and

$$A_{j_1 j_2 \dots j_m}(1) = 1,$$

$$A_{j_1 j_2 \dots j_m}(k) = (j_1 + \dots + j_m - 2)(j_2 + \dots + j_m - 2) \dots (j_{k-1} + \dots + j_m - 2).$$

In summary, we have the estimate $|\operatorname{Re} b_n| \leq \lambda \operatorname{Re}\{1 - b_1\}$, where

$$(7) \quad \lambda = \sum m 2^{m-2} \Gamma_{j_1 j_2 \dots j_m} \sum_{k=1}^m j_k [(j_k + 1)/2] / C_{j_1 j_2 \dots j_m}(k)$$

and the first summation is over all positive integers $j_1, j_2, \dots, j_m$ with $j_m \geq 2$ and $j_1 + j_2 + \dots + j_m = n + 1$. Therefore (3) is proved and also

$$\operatorname{Re}\{\lambda b_1 - b_n\} \leq \lambda \operatorname{Re} b_1 + |\operatorname{Re} b_n| \leq \lambda \operatorname{Re} b_1 + \lambda \operatorname{Re}\{1 - b_1\} = \lambda. \quad \square$$

Estimates for λ . Direct computation of (7) leads, for example, to the estimates

$$|\operatorname{Re} b_2| \leq 8 \operatorname{Re}\{1 - b_1\}, \quad |\operatorname{Re} b_3| \leq (115/3) \operatorname{Re}\{1 - b_1\},$$

$$|\operatorname{Re} b_4| \leq (1963/12) \operatorname{Re}\{1 - b_1\}, \quad |\operatorname{Re} b_5| \leq (6421/10) \operatorname{Re}\{1 - b_1\}.$$

The first inequality is not as good as the sharp inequality $|\operatorname{Re} b_2| \leq 2 \operatorname{Re}\{1 - b_1\}$ proved by Garabedian and Schiffer [2]. They also proved that $\operatorname{Re}\{b_3 - 3b_1\} \geq -3$. Together with the inequality $\operatorname{Re}\{b_3 + 3b_1\} \leq 3$, which is a special case of what is proved in [4, Theorem 3.3], this implies that $|\operatorname{Re} b_3| \leq 3 \operatorname{Re}\{1 - b_1\}$. However, the inequalities for the higher coefficients are the only ones of this form known to the authors. For example, they imply

$$\operatorname{Re}\{(1963/12)b_1 - b_4\} \leq 1963/12 \quad \text{and} \quad \operatorname{Re}\{(6421/10)b_1 - b_5\} \leq 6421/10.$$

It is desirable to have an estimate for (7) which is of a simpler form. Each factor in $\Gamma_{j_1 j_2 \dots j_m}$ is dominated by the corresponding factor in $C_{j_1 j_2 \dots j_m}(k)$; hence $\Gamma_{j_1 j_2 \dots j_m} / C_{j_1 j_2 \dots j_m}(k) \leq 1$. In addition, each $j_k \leq n - m + 2$. Therefore

$$\lambda \leq \sum m^2 2^{m-3} (n - m + 2)(n - m + 3),$$

where the summation is over all positive integers $j_1, j_2, \dots, j_m$ with $j_m \geq 2$ and $j_1 + j_2 + \dots + j_m = n + 1$. For fixed m there are $\binom{n-1}{m-1}$ such terms; thus

$$\begin{aligned}\lambda &\leq \sum_{m=1}^n m^2 2^{m-3} (n-m+2)(n-m+3) \binom{n-1}{m-1} \\ &= (1/4)(4n^4 + 50n^3 + 149n^2 + 277n + 6)3^{n-5} \equiv \mu_n.\end{aligned}$$

As a consequence, with μ_n defined just above, our development gives the weaker, but very explicit inequalities

$$|\operatorname{Re} b_n| \leq \mu_n \operatorname{Re}\{1 - b_1\} \quad \text{and} \quad \operatorname{Re}\{\mu_n b_1 - b_n\} \leq \mu_n.$$

Concluding remark. Just as in Pederson's article [5], we can obtain the following more general theorem.

THEOREM. *Let P be a polynomial of n variables with real coefficients, and assume that $g(z) = z + \sum_{n=0}^{\infty} b_n z^{-n}$ belongs to Σ . Then there exists a constant C , independent of g , such that*

$$|\operatorname{Re}\{P(b_1, b_2, \dots, b_n) - P(1, 0, \dots, 0)\}| \leq C \operatorname{Re}\{1 - b_1\}.$$

REFERENCES

1. P. L. Duren, *Univalent functions*, Springer-Verlag, New York, 1983.
2. P. R. Garabedian and M. Schiffer, *A coefficient inequality for schlicht functions*, Ann. of Math. (2) **61** (1955), 116–136.
3. W. E. Kirwan and G. Schober, *New inequalities from old ones*, Math. Z. **180** (1982), 19–40.
4. Y. J. Leung and G. Schober, *Low order coefficient estimates in the class Σ* , Ann. Acad. Sci. Fenn. Ser. A I Math. (to appear).
5. R. N. Pederson, *A note on the local coefficient problem*, Proc. Amer. Math. Soc. **20** (1969), 345–347.
6. G. Schober and J. K. Williams, *On coefficient estimates and conjectures for the class Σ* , Math. Z. **186** (1984), 309–320.
7. G. Schober, *Some conjectures for the class Σ* , Contemporary Math. (to appear).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF DELAWARE, NEWARK, DELAWARE 19716

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON, INDIANA 47405