A CHARACTERIZATION OF POTENTIAL SPACES

JOSÉ R. DORRONSORO

ABSTRACT. A mean oscillation characterization, valid for all $\alpha > 0$, of the spaces L_α^p of Bessel potentials of L^p functions is given and is used to relate the known characterizations for $0 < \alpha < 2$ via Marcinkiewicz integrals, due to E. M. Stein, and via vector-valued means of differences, due to R. S. Strichartz.

1. Introduction. The Bessel potential of order α , $\alpha > 0$, of a smooth function g is defined as $J_\alpha g = K_\alpha * g$, where

$$(K_\alpha)^\wedge(\xi) = (1 + 4\pi^2|\xi|^2)^{-\alpha/2}.$$

The kernel K_α can be shown to be integrable and, for $1 \leq p \leq \infty$, the potential spaces L_α^p , introduced by Aronszajn and Smith and by Calderón, are then the spaces of Bessel potentials of order α of L^p functions: that is,

$$L_\alpha^p = \{J_\alpha g : g \in L^p\}.$$

With the norm $\|J_\alpha g\|_{p,\alpha} = \|g\|_p$, L_α^p becomes a Banach space.

This paper is concerned with the question of when a given function f is the Bessel potential of an L^p function. An obviously necessary condition is that f be in L^p , and Stein proved [7], for $0 < \alpha < 2$ and $2n/n + 2\alpha < p < \infty$, that an L^p function f belongs to L_α^p if and only if its Marcinkiewicz integral

$$D_\alpha f(x) = \left(\int_{\mathbf{R}^n} |\Delta_y^{[\alpha]+1} f(x)|^2 |y|^{-2\alpha-n} dy \right)^{1/2}$$

is in L^p and, furthermore, $\|f\|_{p,\alpha} \sim \|f\|_p + \|D_\alpha f\|_p$ (by $A \sim B$ we mean there is a constant C independent of A or B such that $C^{-1}A \leq B \leq CA$; also, $\Delta_y f(x) = f(x+y) - f(x)$).

With p now such that $1 < p < \infty$, the following characterization of Bessel potentials for the same range of α is due to Strichartz [9]. Let $[\alpha]$ denote the integral part of α .

THEOREM 1. *Let $0 < \alpha < 2$ and $1 < p < \infty$. An L^p function f belongs to L_α^p if and only if the function*

$$S_\alpha f(x) = \left(\int_0^\infty \left(t^{-\alpha} \int_{|y| \leq 1} |\Delta_{ty}^{[\alpha]+1} f(x)| dy \right)^2 t^{-1} dt \right)^{1/2}$$

belongs to L^p . Furthermore, $\|f\|_{p,\alpha} \sim \|f\|_p + \|S_\alpha f\|_p$.

This result was extended by Bagby [1] to the range $0 < \alpha < n$.

Received by the editors June 8, 1984 and, in revised form, September 21, 1984.
1980 *Mathematics Subject Classification.* Primary 46E35; Secondary 26A16.

We will show here how Bessel potentials of L^p functions, $1 < p < \infty$, can be characterized for a general $\alpha > 0$. This will be done by means of a mixed norm estimate of the approximation of these functions by polynomials. To be more precise, let f be a locally integrable function and Q a cube in $\mathbf{R}^n$. We denote by $P_Q^k f$ the unique polynomial of degree k such that

$$\int_Q (f(y) - P_Q^k f(y)) y^\gamma dy = 0$$

for each n -tuple $\gamma = (\gamma_1, \dots, \gamma_n) \in \mathbf{N}^n$ such that $|\gamma| = \gamma_1 + \dots + \gamma_n \leq k$ (for instance, if $k = 0$, $P_Q^0 f$ is just the mean $f_Q = |Q|^{-1} \int_Q f$). We will write $P_Q f$ or even P_Q if there is no chance of confusion. Then, for $x \in \mathbf{R}^n$ and $t > 0$, we define

$$\Omega_f^k(x, t) = \Omega_f(x, t) = \sup \left\{ |Q|^{-1} \int_Q |f - P_Q^k f| : x \in Q, |Q| = t^n \right\}.$$

The main result of this paper is

THEOREM 2. *Let $\alpha > 0$ and $f \in L^p$, $1 < p < \infty$. Then $f \in L_\alpha^p$ if and only if the function*

$$G_\alpha f(x) = \left(\int_0^\infty (t^{-\alpha} \Omega_f^{[\alpha]}(x, t))^2 t^{-1} dt \right)^{1/2}$$

is in L^p . Furthermore, $\|f\|_{p, \alpha} \sim \|f\|_p + \|G_\alpha f\|_p$.

The paper is organized as follows: §2 contains the proof of Theorem 2 when $0 < \alpha < 1$; in view of Strichartz's theorem, we will prove that $G_\alpha f \in L^p$ if and only if $S_\alpha f \in L^p$. In §3 we deal with the case $\alpha > 0$, α nonintegral, by reducing it to that of §2. Complex interpolation is then used in §4 to deal with the remaining case $\alpha > 0$, α integral. Finally, §5 contains the relationship between $G_\alpha f$ and the Marcinkiewicz integral $D_\alpha f$, and some further results.

2. The case $0 < \alpha < 1$. We begin with some remarks about the polynomials P_Q . Let f be a locally integrable function. An easy homogeneity argument gives [6]

$$(1) \quad \operatorname{ess\,sup}_Q |P_Q^k f| \leq C |Q|^{-1} \int_Q |f|,$$

for any cube Q and $k \geq 0$, and where C does not depend on Q or f (throughout the paper C will denote any absolute constant independent of the particular functions, points or sets considered). As a consequence, for any n -tuple $\gamma \in \mathbf{N}^n$, we have for the γ -derivatives of P_Q (see [3, 4])

$$(2) \quad \begin{aligned} \operatorname{ess\,sup}_Q |D^\gamma(P_Q^k f)| &\leq C |Q|^{-\gamma/n} \operatorname{ess\,sup}_Q |P_Q^k f| \\ &\leq C |Q|^{-\gamma/n-1} \int_Q |f|. \end{aligned}$$

It also follows from (1) that, since $P_Q^k(f + R) = P_Q^k f + R$ for any polynomial R of degree $\leq k$,

$$(3) \quad |Q|^{-1} \int_Q |f - P_Q^k f| \leq C |Q|^{-1} \int_Q |f - R|,$$

and that if $Q \subset Q'$,

$$(4) \quad |Q|^{-1} \int_Q |f - P_Q^k f| \leq C(|Q'|/|Q|)|Q'|^{-1} \int_{Q'} |f - P_{Q'}^k f|.$$

The cube with centre x and side length t will be denoted by $Q_{x,t}$. From (4) we obtain

$$(5) \quad \Omega_f^k(x, t) \leq Ct^{-n} \int_{Q_{x,2t}} |f - P_{Q_{x,2t}}^k|,$$

and therefore, if $t \leq s \leq 2t$, $\Omega_f^k(x, t) \leq C\Omega_f^k(x, s) \leq C\Omega_f^k(x, 2t)$. In particular, the series

$$\left(\sum_{-\infty}^{\infty} (2^{-i\alpha} \Omega_f(x, 2^i))^2 \right)^{1/2}$$

is comparable to G_α , and we can also use balls instead of cubes to define Ω_f and $G_\alpha f$. We have now

THEOREM 3. *Let $\alpha > 0$, α nonintegral, and $f \in L^p$, $1 < p < \infty$. Then $G_\alpha f \in L^p$ if and only if for a.e. $x \in \mathbf{R}^n$ there is a polynomial P_x of degree $\leq [\alpha]$, depending on f , such that if*

$$\tilde{G}_\alpha f(x) = \left(\int_0^\infty \left(t^{-\alpha-n} \int_{Q_{x,t}} |f(y) - P_x(y)| dy \right)^2 t^{-1} dt \right)^{1/2},$$

$\tilde{G}_\alpha f \in L^p$. Furthermore, $\|G_\alpha f\|_p \sim \|\tilde{G}_\alpha f\|_p$. The polynomial P_x will be a.e. the Taylor polynomial of f .

PROOF. The inequality $G_\alpha f \leq C\tilde{G}_\alpha f$ is an immediate consequence of (5) and (3). Conversely, if $G_\alpha f \in L^p$, fix $x \in \mathbf{R}^n$, $\gamma \in \mathbf{N}^n$ with $|\gamma| \leq [\alpha]$ and write $P_{Q_{x,t}}^{[\alpha]} f(y) = P_{Q_{x,t}}(y)$ as

$$P_{Q_{x,t}}(y) = \sum_{|\beta| \leq [\alpha]} a_\beta(x, t)(y - x)^\beta / \beta!.$$

If $t > s > 0$, let $Q_0 \subset Q_1 \subset \dots \subset Q_m$ be a sequence of cubes such that $Q_0 = Q_{x,s}$, $Q_m = Q_{x,t}$ and $|Q_i| = 2^n |Q_{i-1}|$. Then, by (2),

$$\begin{aligned} |a_\gamma(x, t) - a_\gamma(x, s)| &= |D^\gamma(P_{Q_0} - P_{Q_m})(x)| \\ &\leq \sum_{i=1}^m |D^\gamma(P_{Q_i} - P_{Q_{i-1}})(x)| \\ &\leq C \sum_{i=1}^m |Q_i|^{-1-|\gamma|/n} \int_{Q_i} |f - P_{Q_i}| \\ &\leq C \int_s^t \Omega_f(x, u) u^{-|\gamma|-1} du. \end{aligned}$$

Thus, if $\alpha > |\gamma|$, Schwartz's inequality gives

$$(6) \quad |a_\gamma(x, t) - a_\gamma(x, s)| \leq Ct^{\alpha-|\gamma|} G_\alpha f(x).$$

In addition, we have

$$(7) \quad \begin{aligned} |a_\gamma(x, t)| &\leq |a_\gamma(x, t) - a_\gamma(x, 1)| + |a_\gamma(x, 1)| \\ &\leq C(G_\alpha f(x) + Mf(x)), \end{aligned}$$

with M the Hardy-Littlewood maximal operator. Therefore, as t goes to 0, the functions $a_\gamma(x, t)$ form a Cauchy sequence of L^p functions. We denote its L^p limit by f_γ , which will also be a.e. the pointwise limit of some subsequence $a_\gamma(x, t_j)$. Hence, for a.e. x

$$(8) \quad \begin{aligned} |f_\gamma(x) - a_\gamma(x, t)| &= \lim |a_\gamma(x, t_j) - a_\gamma(x, t)| \\ &\leq C \int_0^t \Omega_f(x, u) u^{-|\gamma|-1} du \leq C t^{\alpha-|\gamma|} G_\alpha f(x), \end{aligned}$$

and, in particular, $\|f_\gamma\|_p \leq \|f_\gamma - a_\gamma(\cdot, 1)\|_p + \|a_\gamma(\cdot, 1)\|_p \leq C(\|G_\alpha f\|_p + \|f\|_p)$.

If we now define $P_x(y) = \sum_{|\gamma| \leq k} f_\gamma(x)(y-x)^\gamma/\gamma!$, we obtain

$$\begin{aligned} \int_{Q_{x,t}} |f - P_x| dy &\leq \int_{Q_{x,t}} |f - P_{Q_{x,t}}| dy + C \sum_{|\gamma| \leq k} |a_\gamma(x, t) - f_\gamma(x)| t^{|\gamma|} \\ &\leq C t^{-n} \left(\Omega_f(x, t) + \sum_{i=0}^k t^i \int_0^t \Omega_f(x, u) u^{-i-1} du \right). \end{aligned}$$

Assuming that $k = [\alpha] < \alpha$, Hardy's inequality implies

$$\tilde{G}_\alpha f(x) \leq C G_\alpha f(x) \quad \text{for a.e. } x \in \mathbf{R}^n.$$

In particular, since $a_0(x, t) = f_{Q_{x,t}}$ tends to $f(x)$ a.e., it follows that for $0 < \alpha < 1$, $\tilde{G}_\alpha f(x) = S_\alpha f(x)$ a.e., and Theorem 2 is proved in this case.

3. The case $\alpha > 0$, α nonintegral. The following is a well-known result in the theory of potential spaces (see, for instance [8, p. 136]).

THEOREM 4. *Let $f \in L^p$, $1 < p < \infty$. Then $f \in L^p_\alpha$, $\alpha \geq 1$, if and only if f and its weak partials $\partial f/\partial x_i$ belong to $L^p_{\alpha-1}$. Furthermore, $\|f\|_{p,\alpha} \sim \|f\|_{p,\alpha-1} + \sum_1^n \|\partial f/\partial x_i\|_{p,\alpha-1}$.*

A similar result also holds for L^p functions f such that $G_\alpha f \in L^p$; concretely,

THEOREM 5. *Let $f \in L^p$, $1 < p < \infty$. Then, for $\alpha > 1$, $G_\alpha f \in L^p$ if and only if $G_{\alpha-1} f \in L^p$, the weak partials $\partial f/\partial x_i \in L^p$ and $G_{\alpha-1}(\partial f/\partial x_i) \in L^p$. Furthermore,*

$$\|f\|_p + \|G_\alpha f\|_p \sim \|f\|_p + \|G_{\alpha-1} f\|_p + \sum_1^n (\|\partial f/\partial x_i\|_p + \|G_{\alpha-1}(\partial f/\partial x_i)\|_p).$$

PROOF. We begin with the only if part. Suppose first that f is a C^1 function such that f and $\partial f/\partial x_i$ are in L^p , and fix x and $t > 0$. For each $i = 1, \dots, n$ and $y \in Q_{0,t}$, consider the polynomials

$$P_i(x+y) = \int_0^1 P_{Q_{x, st}}^{k-1}(\partial f/\partial x_i)(x+sy) ds,$$

with $k = [\alpha]$. Then, by Taylor's formula,

$$\begin{aligned} & t^{-n} \int_{Q_{x,t}} \left| f(z) - f(x) - \sum_1^n P_i(z)(z_i - x_i) \right| dz \\ & \leq t^{-n} \sum_1^n \int_0^1 t \int_{Q_{0,t}} |\partial f / \partial x_i(x + sy) - P_{Q_{x,st}}^{k-1}(\partial f / \partial x_i)(x + sy)| dy ds \\ & \leq C \sum_1^n t \int_0^1 (st)^{-n} \int_{Q_{0,st}} |\partial f / \partial x_i(x + z) - P_{Q_{x,st}}^{k-1}(x + z)| dz ds, \end{aligned}$$

which implies

$$\Omega_f^k(x, t) \leq C \sum_1^n \int_0^{2t} \Omega_{\partial f / \partial x_i}^{k-1}(x, u) du.$$

This estimate can be extended to any $f \in L^p$ with weak partials in L^p by a standard regularization argument and, in the conditions of the theorem, Hardy's inequality yields

$$G_\alpha f(x) \leq C \sum_1^n G_{\alpha-1}(\partial f / \partial x_i)(x).$$

Next, we will divide the proof of the if part into three lemmas.

LEMMA 1. *Let $f \in L^p$ such that $G_\alpha f \in L^p$, and $\gamma \in \mathbf{N}^n$, $|\gamma| < \alpha$. Then, with f_γ as in Theorem 3, $\|G_{\alpha-|\gamma|} f_\gamma\|_p \leq C \|G_\alpha f\|_p$.*

PROOF. Fix $x \in \mathbf{R}^n$ and a cube Q with $x \in Q$ and $|Q| = t^n$. If $y \in Q$, $Q_{y,t} \subset Q_{x,2t}$, and by (8) and (2),

$$\begin{aligned} |f_\gamma(y) - D^\gamma P_{Q_{x,2t}}(y)| & \leq |f_\gamma(y) - a_\gamma(y, t)| + |D^\gamma(P_{Q_{y,t}} - P_{Q_{x,2t}})(y)| \\ & \leq C \int_0^t \Omega_f(y, s) s^{-|\gamma|-1} ds + Ct^{-|\gamma|} \Omega_f(x, 2t). \end{aligned}$$

Therefore,

$$\begin{aligned} \Omega_{f_\gamma}(x, t) & \leq Ct^{-n} \int_{Q_{x,2t}} dy \left(\int_0^t \Omega_f(y, s) s^{-|\gamma|-1} ds \right) + Ct^{-|\gamma|} \Omega_f(x, 2t) \\ & \leq C \int_0^t M(\Omega_f(\cdot, s))(x) s^{-|\gamma|-1} ds + Ct^{-|\gamma|} \Omega_f(x, 2t), \end{aligned}$$

where M denotes the Hardy-Littlewood maximal operator. Now, Hardy's inequality gives

$$G_{\alpha-|\gamma|}(f_\gamma)(x) \leq C \left(G_\alpha f(x) + \left(\int_0^\infty (t^{-\alpha} M \Omega_f(\cdot, t))(x)^2 t^{-1} dt \right)^{1/2} \right),$$

and the lemma follows by the Fefferman-Stein theorem on vector-valued maximal operators [5].

LEMMA 2. Let $f \in L^p$ be such that $G_\alpha f \in L^p$, $\alpha > 1$. Then the weak partials $\partial f / \partial x_i$ exist and coincide with f_{e_i} , where e_i denotes the n -tuple with 1 in the i th place and 0 in the others.

PROOF. It is enough to show that $(f(x + te_i) - f(x))/t$ tends to $f_{e_i}(x)$ in L^p as t goes to 0. First, we have by (8),

$$\begin{aligned} & |t^{-1}(f(x + te_i) - f(x)) - f_{e_i}(x)| \\ & \leq t^{-1}|f(x + te_i) - a_0(x + te_i, 2t)| + t^{-1}|f(x) - a_0(x, 2t)| \\ & \quad + |t^{-1}(a_0(x + te_i, 2t) - a_0(x, 2t)) - f_{e_i}(x)| \\ & \leq Ct^{\alpha-1}(G_\alpha f(x + te_i) + G_\alpha f(x)) \\ & \quad + |t^{-1}(a_0(x + te_i, 2t) - a_0(x, 2t)) - f_{e_i}(x)|. \end{aligned}$$

Next, since $a_0(y, t) = P_{Q_{y,t}}(y)$,

$$\begin{aligned} & a_0(x + te_i, 2t) - a_0(x, 2t) \\ & = P_{Q_{x+te_i, 2t}}(x + te_i) - P_{Q_{x, 2t}}(x + te_i) + P_{Q_{x, 2t}}(x + te_i) - P_{Q_{x, 2t}}(x) \\ & = P_{Q_{x+2te_i, 2t}}(x + te_i) - P_{Q_{x, 2t}}(x + te_i) + \sum_{j=1}^{[\alpha]} a_{je_i}(x, 2t)t^j/j!; \end{aligned}$$

thus,

$$\begin{aligned} & |t^{-1}(a_0(x + te_i, 2t) - a_0(x, 2t)) - f_{e_i}(x)| \\ & \leq |t^{-1}(a_0(x + te_i, 2t) - a_0(x, 2t)) - a_{e_i}(x, 2t)| + |a_{e_i}(x, 2t) - f_{e_i}(x)| \\ & \leq t^{-1}|P_{Q_{x+te_i, 2t}}(x + te_i) - P_{Q_{x, 2t}}(x + te_i)| \\ & \quad + C \sum_{j=2}^{[\alpha]} |a_{je_i}(x, 2t)|t^{j-1} + Ct^{\alpha-1}G_\alpha f(x) \\ & \leq Ct^{-1}t^{-n} \int_{Q_{x+te_i, t}} |P_{Q_{x+te_i, 2t}}(y) - P_{Q_{x, 2t}}(y)| dy \\ & \quad + C \sum_{j=2}^{[\alpha]} t^{j-1}|a_{je_i}(x, 2t)| + Ct^{\alpha-1}G_\alpha f(x), \end{aligned}$$

and taking L^p norms, we obtain

$$\begin{aligned} & \|t^{-1}(f(\cdot + te_i) - f(\cdot)) - f_{e_i}(\cdot)\|_p \\ & \leq Ct^{-1}\|\Omega_f(\cdot, 2t)\|_p + C \sum_{j=2}^{[\alpha]} t^{j-1}\|a_{je_i}(\cdot, 2t)\|_p + Ct^{\alpha-1}\|G_\alpha f\|_p. \end{aligned}$$

But $\|\Omega_f(\cdot, 2t)\|_p \leq Ct^\alpha\|G_\alpha f\|_p$ and, if $j < [\alpha]$, $\|a_{je_i}(\cdot, 2t)\|_p \leq C(\|G_\alpha f\|_p + \|f\|_p)$, by (7), whereas if $j = [\alpha] = \alpha$, an easy modification of (7) gives

$$\|a_{je_i}(\cdot, 2t)\|_p \leq C(\log t^{-1})(\|G_\alpha f\|_p + \|f\|_p).$$

In any case $\|t^{-1}(f(\cdot + te_i) - f(\cdot)) - f_{e_i}(\cdot)\|_p$ tends to 0 as t goes to 0.

To finish Theorem 5 we will prove

LEMMA 3. *Let f be an L^p function such that $G_\alpha f \in L^p$, $\alpha > 1$. Then $G_{\alpha-1}f \in L^p$ and $\|G_{\alpha-1}f\|_p \leq C(\|G_\alpha f\|_p + \|f\|_p)$.*

PROOF. Let $k = [\alpha]$. We will follow the lines of the proofs of similar facts in [3] or [6] (see also [4]). Fix $x \in \mathbf{R}^n$ and $t > 0$. For each cube Q containing x and such that $|Q| = t^n$, write $P_Q^k f$ as

$$P_Q^k f(y) = \sum_{|\gamma| < k} a_\gamma(Q)(y-x)^\gamma + \sum_{|\gamma| = k} a_\gamma(Q)(y-x)^\gamma = R_Q(y) + S_Q(y).$$

Then, by (3),

$$(9) \quad \begin{aligned} |Q|^{-1} \int_Q |f - P_Q^{k-1} f| &\leq C|Q|^{-1} \int_Q |f - R_Q| \\ &\leq C \left(\Omega_f^k(x, t) + \text{ess sup}_Q |S_Q| \right). \end{aligned}$$

Denoting by Q_i the cube with same centre as Q and side length $2^i t$, (2) implies, for each $|\gamma| = k$,

$$|a_\gamma(Q_i)| \leq C(2^i t)^{-k} |Q_i|^{-1} \int_{Q_i} |f| \leq C(2^i t)^{-k-n/p} \|f\|_p;$$

that is, $a_\gamma(Q_i)$ tends to 0 as i goes to ∞ and, again by (2),

$$|a_\gamma(Q)| \leq \sum_{i=0}^{\infty} |a_\gamma(Q_{i+1}) - a_\gamma(Q_i)| \leq C \sum_{i=0}^{\infty} (2^i t)^{-k} \Omega_f^k(x, 2^i t).$$

Therefore, inserting this estimate in (9), we obtain

$$\Omega_f^{k-1}(x, t) \leq C \left(\Omega_f^k(x, t) + t^k \int_t^\infty \Omega_f^k(x, s) s^{-k-1} ds \right).$$

Now Hardy's inequality gives

$$\begin{aligned} G_{\alpha-1}f(x) &\leq C \left(\int_0^\infty (t^{-\alpha+1} \Omega_f^k(x, t))^2 t^{-1} dt \right)^{1/2} \\ &\leq C \left(\left(\int_0^1 \right)^{1/2} \right) + C \left(\left(\int_1^\infty \right)^{1/2} \right) \\ &\leq C \left(\int_0^1 (t^{-\alpha} \Omega_f^k(x, t))^2 t^{-1} dt \right)^{1/2} + CMf(x), \end{aligned}$$

and the lemma follows.

Theorems 4 and 5 imply that, if the Theorem 2 holds for a given α , it also holds for $\alpha + 1$. Therefore, the already proved case $0 < \alpha < 1$ implies that Theorem 2 is true for all $\alpha > 0$, α nonintegral, and that if we can prove it for $\alpha = 1$, it will hold for α integral as well. We do this in the following section.

4. The case α integral. Let f be a locally integrable function and fix an integer k and a cube Q with side length $t > 0$. For any $y \in Q$ and any cube Q' such that $y \in Q' \subset Q$, let $Q' = Q_0 \subset Q_1 \subset \dots \subset Q_m = Q$ be a sequence of cubes such that $|Q_i| = 2^n |Q_{i-1}|$. By (1) and (4) we have

$$|Q'|^{-1} \int_{Q'} |f - P_Q^k f| \leq C \sum_0^m |Q_i|^{-1} \int_{Q_i} |f - P_{Q_i}| \leq C \int_0^t \Omega_f(y, s) s^{-1} ds$$

which by Lebesgue's differentiation theorem gives, for a.e. $y \in Q$,

$$(10) \quad \begin{aligned} |f(y) - P_Q(y)| &= \lim_{Q' \rightarrow y} |Q'|^{-1} \int_{Q'} |f - P_Q| \\ &\leq C \int_0^t \Omega_f(y, s) s^{-1} ds. \end{aligned}$$

Now, the second difference operator Δ_h^2 annihilates polynomials of degree 1, and for a.e. x we obtain by (10)

$$\begin{aligned} t^{-n} \int_{Q_{0,t}} |\Delta_h^2 f(x)| dh &= t^{-n} \int_{Q_{0,t}} |\Delta_h^2 (f - P_{Q_{x,2t}}^1)(x)| dh \\ &\leq C \int_0^{2t} M\Omega_f^1(\cdot, s)(x) s^{-1} ds, \end{aligned}$$

which by Hardy's inequality implies

$$\begin{aligned} S_1 f(x) &= \left(\int_0^\infty \left(t^{-1} t^{-n} \int_{Q_{0,t}} |\Delta_h^2 f(x)| dh \right)^2 t^{-1} dt \right)^{1/2} \\ &\leq C \left(\int_0^\infty (t^{-1} M\Omega_f^1(\cdot, t)(x))^2 t^{-1} dt \right)^{1/2}. \end{aligned}$$

Thus, if $G_1 f \in L^p$, $1 < p < \infty$, the L^p boundedness of the vector-valued Hardy-Littlewood maximal operator gives that $\|S_1 f\|_p \leq C \|G_1 f\|_p$.

Conversely, to prove that $f \in L_1^p$ implies $G_1 f \in L^p$, we will use complex interpolation. If $(A, B)_{[\theta]}$ denotes the intermediate space between the Banach spaces A, B obtained by the complex method for the index θ , it is well known (see, for instance [2]) that $(L_{\alpha_1}^p, L_{\alpha_2}^p)_{[\theta]} = L_{\alpha}^p$, with $\alpha = \theta\alpha_1 + (1 - \theta)\alpha_2$. Suppose now that $0 < \alpha_1 < 1 < \alpha_2 < 2$ and let H_{α_i} , $i = 1, 2$, denote the Banach space of functions $F(y, t)$ defined in the product $(0, \infty) \times Q_{0,1}$ such that

$$\|F\|_{H_{\alpha_i}} = \left(\int_0^\infty \left(t^{-\alpha_i} \int_{Q_{0,1}} |F(y, t)| dy \right)^2 t^{-1} dt \right)^{1/2} < \infty.$$

By the results in §§2 and 3 and the inequality $\Omega_f^1 \leq C\Omega_f^0$ derived from (4), the map

$$f \rightarrow F(y, t; x) = f(x + ty) - P_{Q_{x,t}}^1 f(x + ty)$$

is a bounded linear map from $L_{\alpha_i}^p$ to the L^p space of H_{α_i} valued functions $L^p(H_{\alpha_i})$ and, therefore, from $L_1^p = (L_{\alpha_1}^p, L_{\alpha_2}^p)_{[\theta]}$ into $(L^p(H_{\alpha_1}), L^p(H_{\alpha_2}))_{[\theta]}$ with $\theta =$

$\alpha_2 - 1/\alpha_2 - \alpha_1$. But $(L^p(H_{\alpha_1}), L^p(H_{\alpha_2}))_{[\theta]} = L^p((H_{\alpha_1}, H_{\alpha_2})_{[\theta]})$ and $(H_{\alpha_1}, H_{\alpha_2})_{[\theta]} = H_1$ [2, pp. 107 and 122]. Therefore, if $f \in L_1^p$,

$$\|G_1 f\|_p \leq C \|(\|F(\cdot, \cdot; x)\|_{H_1})\|_p \leq C \|f\|_{p,1}.$$

This concludes the proof of Theorem 2 when $\alpha = 1$ and, by the preceding remarks, for all $\alpha > 0$.

5. Further results. First, we will show how Theorem 2 relates with the Marcinkiewicz integral characterization of L_α^p when $p > 2n/n + 2\alpha$. For simplicity, we will give the details only for $0 < \alpha < 1$. In this range of α , the Marcinkiewicz integral of a function f can be written as

$$\begin{aligned} (D_\alpha f(x))^2 &= \int_{\mathbf{R}^n} |f(x+y) - f(x)|^2 |y|^{-2\alpha-n} dy \\ &= \int_{\mathbf{R}^n} |f(x+y) - f(x)|^2 \int_{|y|}^\infty t^{-2\alpha-n-1} dt dy \\ &= \int_0^\infty t^{-2\alpha-n} \left(\int_{|y| \leq t} |f(x+y) - f(x)|^2 dy \right) t^{-1} dt, \end{aligned}$$

which can be seen as a variant of $G_\alpha f$. This suggests the possibility of replacing in the definition of G_α the L^1 means $\Omega_f(x, t)$ by the L^r means

$$\Omega_{f,r}(x, t) = \sup \left\{ \left(|Q|^{-1} \int_Q |f - P_Q|^r \right)^{1/r} : x \in Q, |Q| = t^n \right\}.$$

The following theorem tells us when this can be done.

THEOREM 6. *Let $\alpha > 0$ and f be an L^p function, $1 < p < \infty$, such that $G_\alpha f \in L^p$. Then*

(i) *if $1 < p \leq 2$ and $\alpha \leq n/p$, the function $G_{\alpha,r} f$ obtained by replacing Ω_f with $\Omega_{f,r}$ in the definition of $G_\alpha f$ is also in L^p for $r < pn/n - p\alpha$; if $\alpha > n/p$, $G_{\alpha,\infty} \in L^p$;*

(ii) *if $2 \leq p < \infty$ and $\alpha \leq n/2$, $G_{\alpha,r} f \in L^p$ for $r < 2n/n - 2\alpha$; if $\alpha > n/2$, $G_{\alpha,\infty} f \in L^p$.*

Finally, in all cases $\|G_{\alpha,r} f\|_p \sim \|G_\alpha f\|_p$.

PROOF. Fix $x \in \mathbf{R}^n$, $t > 0$ and a cube Q such that $x \in Q$, $|Q| = t^n$. By (10) we have a.e. in Q

$$\begin{aligned} (11) \quad |f(y) - P_Q(y)| &\leq C \int_0^t \Omega_f(y, s) s^{-1} ds \\ &\leq C \int_0^t \left(s^{-n} \int_{Q_{y,s}} \Omega_f(z, s) dz \right) s^{-1} ds. \end{aligned}$$

Suppose now $1 < p \leq 2$ and $\alpha \leq n/p$, and for $r < pn/n - p\alpha$, fix $q < p$ and $\beta < \alpha$ such that $1/r = 1/q - \beta/n$. By (11),

$$|f(y) - P_Q(y)| \leq C \int_0^t s^{-\beta} M_\beta(\Omega_f(\cdot, s) \chi_Q)(y) s^{-1} ds,$$

where M_β denotes the maximal operator $M_\beta g(z) = \sup\{|Q|^{-1+\beta/n} \int_Q |g| : z \in Q\}$. As it is well known, M_β maps L^q into L^r and, therefore,

$$\begin{aligned} \left(\int_Q |f - P_Q|^r \right)^{1/r} &\leq C \int_0^t s^{-\beta} \left(\int_Q (M_\beta(\Omega_f(\cdot, s)\chi_Q))^r dy \right)^{1/r} s^{-1} ds \\ &\leq C \int_0^t s^{-\beta} \left(\int_Q \Omega_f(z, s)^q dz \right)^{1/q} s^{-1} ds \end{aligned}$$

which implies

$$\Omega_{f,r}(x, t) \leq Ct^\beta \int_0^t s^{-\beta} M_q \Omega_f(\cdot, s)(x) s^{-1} ds,$$

with $M_q(g) = (M(|g|^q))^{1/q}$. Hardy's inequality gives now

$$G_{\alpha,r}f(x) \leq C \left(\int_0^\infty (t^{-\alpha} M_q(\Omega_f(\cdot, t))(x))^2 t^{-1} dt \right)^{1/2}$$

and the estimate $\|G_{\alpha,r}f\|_p \leq C\|G_\alpha f\|_p$ follows from the Fefferman-Stein theorem. The same proof works for the case $\alpha \leq n/2$ of (ii): taking $r < 2n/n - 2\alpha$ allows us to choose $q < 2$, and to apply the Fefferman-Stein theorem again.

If $p \leq 2$ and $\alpha > n/p$, fix β and q such that $\alpha > \beta > n/q > n/p$. From (11) we deduce for a.e. $y \in Q$

$$\begin{aligned} |f(y) - P_Q(y)| &\leq C \int_0^t s^{\beta-n/q} s^{-\beta} \left(\int_{Q_{y,s}} \Omega_f(z, s)^q dt \right)^{1/q} s^{-1} ds \\ (12) \quad &\leq Ct^{\beta-n/q} \int_0^t s^{-\beta} \left(\int_Q \Omega_f(z, s)^q dz \right)^{1/q} s^{-1} ds \\ &\leq Ct^\beta \int_0^t s^{-\beta} M_q \Omega_f(\cdot, s)(x) s^{-1} ds \end{aligned}$$

and the inequality $\|G_{\alpha,\infty}f\|_p \leq C\|G_\alpha f\|_p$ follows as before. The same argument works for the case $\alpha > n/2$ of (ii). Since the estimate $\|G_\alpha f\|_p \leq C\|G_{\alpha,r}f\|_p$ is obvious, Theorem 6 is proved.

As a consequence, $G_{\alpha,2}f$ characterizes L_α^p for $p > 2n/n + 2\alpha$ and, since $D_\alpha f \sim G_{\alpha,2}f$, so does $D_\alpha f$.

Finally, the characterization of potential spaces in terms of the function G_α provides easy proofs of the well-known imbeddings $L_\alpha^p \subset L_\beta^q$, $1/p - \alpha/n = 1/q - \beta/n$, and can be used to show that, for $\alpha > n/p$, L_α^p is an algebra under multiplication. Also, Besov spaces $B_\alpha^{p,q}$ of L^p functions (see [8] for the definitions) can be characterized by the finiteness of the expression

$$\left(\int_0^\infty (t^{-\alpha} \|\Omega_f^{[\alpha]}(\cdot, t)\|_p)^q t^{-1} dt \right)^{1/q}.$$

Combining this with Theorem 2, easy proofs of the imbeddings $B_\alpha^{p,p} \subset L_\alpha^p \subset B_\alpha^{p,2}$, $1 < p \leq 2$, $B_\alpha^{p,2} \subset L_\alpha^p \subset B_\alpha^{p,p}$, $2 \leq p < \infty$ and $B_\alpha^{p,\infty} \subset L_{\alpha-\varepsilon}^p$ for all $\varepsilon > 0$ follow.

REFERENCES

1. R. J. Bagby, *A characterization of Riesz potentials and an inversion formula*, Indiana Univ. Math. J. **29** (1980), 581–595.
2. J. Bergh and J. Löfström, *Interpolation spaces: An introduction*, Springer-Verlag, Berlin, 1976.
3. S. Campanato, *Proprieta di una famiglia di spazi funzionali*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **18** (1964), 137–160.
4. R. De Vore and R. Sharpley, *Maximal operators and smoothness*, Mem. Amer. Math. Soc. No. 293 (1984).
5. C. Fefferman and E. M. Stein, *Some maximal inequalities*, Amer. J. Math. **93** (1971), 107–115.
6. S. Janson, M. Taibleson and G. Weiss, *Elementary characterizations of Morrey-Campanato spaces*, Lecture Notes in Math., vol. 992, Springer-Verlag, 1983, pp. 101–114.
7. E. M. Stein, *The characterization of functions arising as potentials*, Bull. Amer. Math. Soc. **67** (1961), 102–104.
8. ———, *Singular integrals and differentiability properties of functions*, Princeton Univ. Press, Princeton, N.J., 1970.
9. R. S. Strichartz, *Multipliers on fractional Sobolev spaces*, J. Math. Mech. **16** (1967), 1031–1061.

DIVISIÓN DE MATEMATICAS, FACULTAD DE CIENCIAS, UNIVERSIDAD AUTONOMA,
28049 MADRID, SPAIN