A q -POLYNOMIAL IDENTITY

KUNG-WEI YANG

ABSTRACT. We show that the q -polynomial coefficients $\left[\begin{smallmatrix} n \cdot m \\ r \end{smallmatrix} \right]$ (see the definition preceding Theorem 1), which are the generating functions of the number of inversions between multisets, satisfy the beautiful identity

$$\prod_{i=1}^n \left(\sum_{j=0}^m q^{jm(i-1) + \binom{j}{2}} t^j \right) = \sum_{r=0}^{nm} \left[\begin{smallmatrix} n \cdot m \\ r \end{smallmatrix} \right] q^{\binom{r}{2}} t^r.$$

1. Inversions between multisets. Let $(m_1, \dots, m_n)$ denote the *multiset* on $\{1, \dots, n\}$, in which the multiplicity of i is m_i . The number of elements in $(m_1, \dots, m_n)$ is $m_1 + \dots + m_n$ and is denoted by $|(m_1, \dots, m_n)|$. We abbreviate the multiset $(m_1, \dots, m_n)$, where $m_1 = \dots = m_n = m$, to $(n \cdot m)$.

A *multisubset* $(a_1, \dots, a_n)$ of $(n \cdot m)$ satisfies $a_i \leq m$, for $i = 1, \dots, n$, and it uniquely defines a *complementary multisubset* $(\bar{a}_1, \dots, \bar{a}_n)$ satisfying $a_i + \bar{a}_i = m$ ($i = 1, \dots, n$).

An *inversion* between the multisets $(a_1, \dots, a_n)$ and $(b_1, \dots, b_n)$, in that order, is a pair (i, j) , where i is an element of $(a_1, \dots, a_n)$, and j is an element of $(b_1, \dots, b_n)$, and $i > j$.

Let $(a_1, \dots, a_n)$ be a multisubset of $(n \cdot m)$. We denote by $I(a_1, \dots, a_n)$ the number of inversions between $(a_1, \dots, a_n)$ and $(\bar{a}_1, \dots, \bar{a}_n)$. We let $\left[\begin{smallmatrix} n \cdot m \\ r \end{smallmatrix} \right]$ denote the generating function

$$\left[\begin{smallmatrix} n \cdot m \\ r \end{smallmatrix} \right] = \sum_{|(a_1, \dots, a_n)|=r} q^{I(a_1, \dots, a_n)}.$$

Our first theorem is a generalization of both the Gaussian recurrence for the q -binomial (Gaussian) coefficients [2, p. 100] and the polynomial coefficient recurrence [1, p. 77].

THEOREM 1 [3].

$$\left[\begin{smallmatrix} n \cdot m \\ r \end{smallmatrix} \right] = \sum_{i=0}^m q^{i((n-1)m - (r-i))} \left[\begin{smallmatrix} (n-1) \cdot m \\ r-i \end{smallmatrix} \right].$$

Received by the editors September 27, 1984 and, in revised form, December 21, 1984.

1980 *Mathematics Subject Classification*. Primary 05A19; Secondary 05A15.

Key words and phrases. q -polynomial identity, q -polynomial coefficient, inversion between multisets.

PROOF.

$$\begin{aligned}
 \begin{bmatrix} n \cdot m \\ r \end{bmatrix} &= \sum_{|(a_1, \dots, a_n)|=r} q^{I(a_1, \dots, a_n)} \\
 &= \sum_{i=0}^m \sum_{|(a_1, \dots, a_{n-1}, i)|=r} q^{I(a_1, \dots, a_{n-1}, i)} \\
 &= \sum_{i=0}^m q^{i(\bar{a}_1 + \dots + \bar{a}_{n-1})} \sum_{|(a_1, \dots, a_{n-1})|=r-i} q^{I(a_1, \dots, a_{n-1})} \\
 &= \sum_{i=0}^m q^{i((n-1)m - (r-i))} \begin{bmatrix} (n-1) \cdot m \\ r-i \end{bmatrix},
 \end{aligned}$$

where the third equality is explained by

$$\begin{aligned}
 I(a_1, a_2, \dots, a_{n-1}, i) \\
 &= a_2 \bar{a}_1 + a_3(\bar{a}_1 + \bar{a}_2) + \dots + a_{n-1}(\bar{a}_1 + \dots + \bar{a}_{n-2}) + i(\bar{a}_1 + \dots + \bar{a}_{n-1}) \\
 &= I(a_1, \dots, a_{n-1}) + i(\bar{a}_1 + \dots + \bar{a}_{n-1}). \quad \text{Q.E.D.}
 \end{aligned}$$

Because of Theorem 1, we will call $\begin{bmatrix} n \cdot m \\ r \end{bmatrix}$, a q -polynomial coefficient.

2. Identity.

THEOREM 2.

$$\prod_{i=1}^n \left(\sum_{j=0}^m q^{jm(i-1) + \binom{j}{2}} t^j \right) = \sum_{r=0}^{nm} \begin{bmatrix} n \cdot m \\ r \end{bmatrix} q^{\binom{r}{2}} t^r.$$

PROOF. We prove the identity by induction on n , using Theorem 1. The identity obviously holds when $n = 1$. Assume that it holds for $n - 1$. Then,

$$\begin{aligned}
 &\prod_{i=0}^n \left(\sum_{j=0}^m q^{jm(i-1) + \binom{j}{2}} t^j \right) \\
 &= \left(\prod_{i=0}^{n-1} \left(\sum_{j=0}^m q^{jm(i-1) + \binom{j}{2}} t^j \right) \right) \left(\sum_{j=0}^m q^{jm(n-1) + \binom{j}{2}} t^j \right) \\
 &= \left(\sum_{k=0}^{(n-1)m} \begin{bmatrix} (n-1) \cdot m \\ k \end{bmatrix} q^{\binom{k}{2}} t^k \right) \left(\sum_{j=0}^m q^{jm(n-1) + \binom{j}{2}} t^j \right) \\
 &= \sum_{r=0}^{nm} \left(\sum_{i=0}^m \begin{bmatrix} (n-1) \cdot m \\ r-i \end{bmatrix} q^{i((n-1)m - (r-i))} \right) q^{\binom{r}{2}} t^r \\
 &= \sum_{r=0}^{nm} \begin{bmatrix} n \cdot m \\ r \end{bmatrix} q^{\binom{r}{2}} t^r. \quad \text{Q.E.D.}
 \end{aligned}$$

If we set $m = 1$ in Theorem 2, we obtain the well-known identity [2, p. 101]

$$(1+t)(1+tq) \cdots (1+tq^{n-1}) = \sum_{r=0}^n \begin{bmatrix} n \\ r \end{bmatrix} q^{\binom{r}{2}} t^r.$$

REFERENCES

1. L. Comtet, *Advanced combinatorics*, Reidel, Boston, Mass., 1974.
2. I. P. Goulden and D. M. Jackson, *Combinatorial enumeration*, Wiley, New York, 1983.
3. K.-W. Yang, *Generalized Euler and Chu-Vandermonde identities*, J. Combin. Theory Ser. A (to appear).

DEPARTMENT OF MATHEMATICS, WESTERN MICHIGAN UNIVERSITY, KALAMAZOO,
MICHIGAN 49008