

ON A CONJECTURE OF KÀTAI CONCERNING WEAKLY COMPOSITE NUMBERS

JANOS GALAMBOS

ABSTRACT. A number is called weakly composite if the sum of the reciprocals of its prime divisors is bounded by two. In this note it is proved that, for $n \geq n_0$, there is a weakly composite number between n and $n + \log \log \log n$.

We call a number weakly composite if the sum of the reciprocals of its distinct prime divisors is bounded by two. We prove that, for all integers $n \geq n_0$, there is a weakly composite number between n and $n + \log \log \log n$.

Let $q_1 < q_2 < \dots < q_N$ be the prime divisors of n . We say that n is weakly composite if

$$(1) \quad g(n) = \sum_{j=1}^N \frac{1}{q_j} \leq 2.$$

One of the consequences of the result of Hausman [2] is that, for fixed k , there are infinitely many integers n such that none of $n, n+1, n+2, \dots, n+k$ is weakly composite. In other words, if $n_t, t \geq 1$, is the sequence of weakly composite numbers then, as $t \rightarrow +\infty$,

$$(2) \quad \limsup(n_{t+1} - n_t) = +\infty.$$

On the other hand, it easily follows from the most elementary results of probabilistic number theory (see, e.g., Elliott [1, Chapter 5, in particular the concluding remarks]) that the average of the gaps $n_{t+1} - n_t$ is bounded, and so (2) is very slowly diverging. When lecturing at Temple University on related topics, I. Kàtai of Budapest formulated the conjecture that the gaps $n_{t+1} - n_t$ must be bounded by a function of the magnitude of $\log \log \log t$. We prove this conjecture by establishing the following result.

THEOREM. *For all sufficiently large real numbers n , there is a weakly composite number between n and $n + \log \log \log n$.*

PROOF. We use an idea of Kàtai [3], in which he generalizes the result of Hausman. This proof also incorporates simplifications by the referee with the referee's kind permission.

Set $k = \log \log \log n$. It suffices to show that

$$(3) \quad S = \sum_{n < m \leq n+k} g(m) < 2k.$$

Received by the editors March 28, 1985.

1980 *Mathematics Subject Classification.* Primary 10A05, 10H25.

©1986 American Mathematical Society
 0002-9939/86 \$1.00 + \$.25 per page

We write the left side as $S_1 + S_2$, where

$$S_1 = \sum_{n < m \leq n+k} \sum_{\substack{p|m \\ p < \log n}} \frac{1}{p}$$

and

$$S_2 = \sum_{n < m \leq n+k} \sum_{\substack{p|m \\ p \geq \log n}} \frac{1}{p}.$$

Now

$$\begin{aligned} S_1 &= \sum_{p < \log n} \frac{1}{p} \sum_{\substack{n < m \leq n+k \\ p|m}} 1 \leq \sum_{p < \log n} \frac{1}{p} \left(\frac{k}{p} + 1 \right) \\ &< k \sum_p p^{-2} + \sum_{p < \log n} p^{-1}. \end{aligned}$$

From the asymptotic formula

$$\sum_{p < x} \frac{1}{p} = \log \log x + O(1)$$

and the elementary inequality

$$\sum_p p^{-2} < \sum_{p < 51} p^{-2} + \frac{1}{51} < \frac{1}{2}$$

we see that

$$(4) \quad S_1 < 3k/2 + O(1).$$

Next we treat S_2 . Observe that if $h(m)$ is the number of prime divisors of m exceeding $\log n$, then

$$S_2 \leq \frac{1}{\log n} \sum_{n < m \leq n+k} h(m).$$

For any $m \leq n+k$, we have

$$(\log n)^{h(m)} \leq \prod_{p|m} p \leq n+k < 2n,$$

from which, upon taking the logarithm, we see that

$$h(m) < (\log 2 + \log n)/(\log \log n).$$

Consequently, for n large, $S_2 < 2k/(\log \log n)$. Combining this with (4), we get

$$S \leq k(3/2 + O(1/\log \log n)).$$

The inequality at (3) then follows for n sufficiently large. The proof is completed.

REFERENCES

1. P. D. T. A. Elliott, *Probabilistic number theory*. I, Springer-Verlag, New York, 1979.
2. M. Hausman, *Generalization of a theorem of Landau*, Pacific J. Math. **84** (1979), 91–95.
3. I. Kátai, *A minimax theorem for additive functions*, Publ. Math. Debrecen **30** (1983), 249–252.

DEPARTMENT OF MATHEMATICS, TU 038-16, TEMPLE UNIVERSITY, PHILADELPHIA, PENNSYLVANIA 19122