THE n TH ROOTS OF SOLUTIONS OF LINEAR ORDINARY DIFFERENTIAL EQUATIONS¹

WILLIAM A. HARRIS, JR. AND YASUTAKA SIBUYA²

ABSTRACT. In this paper we shall prove the following theorem: Let K be a differential field of characteristic zero. Let φ and ψ be elements of a differential field extension of K such that (i) $\varphi \neq 0$ and $\psi \neq 0$; (ii) φ and ψ satisfy nontrivial linear differential equations with coefficients in K , say, $P(\varphi) = 0$ and $Q(\psi) = 0$; (iii) $\varphi = \psi^n$ for some positive integer n such that $n \geq \text{ord } P$. Then the logarithmic derivatives of φ and ψ are algebraic over K . (Note that $\varphi'/\varphi = n(\psi'/\psi)$.)

1. Introduction. Let K be a differential field of characteristic zero. We denote by $\mathcal{D}_K$ the ring of linear ordinary differential operators with coefficients in K , that is

$$\mathcal{D}_K = \left\{ \sum_{k=0}^m a_k D^k; a_k \in K, m \in N \right\},$$

where D denotes differentiation in any differential field extension of K and N is the set of all nonnegative integers. In a previous paper [2] we characterized those functions which together with their reciprocal satisfy linear differential equations:

THEOREM A. *Let φ be an element of a differential field extension of K such that*

- (i) $\varphi \neq 0$;
- (ii) $P(\varphi) = 0$ for some $P \in \mathcal{D}_K - \{0\}$;
- (iii) $Q(1/\varphi) = 0$ for some $Q \in \mathcal{D}_K - \{0\}$.

Then, the logarithmic derivative of φ ($= \varphi'/\varphi$) is algebraic over K .

In the present paper, utilizing a similar method, we shall prove the following result:

THEOREM B. *Let φ and ψ be elements of a differential field extension of K such that*

- (i) $\varphi \neq 0$ and $\psi \neq 0$;
- (ii) $P(\varphi) = 0$ for some $P \in \mathcal{D}_K - \{0\}$;
- (iii) $Q(\psi) = 0$ for some $Q \in \mathcal{D}_K - \{0\}$;
- (iv) $\varphi = \psi^n$ for some positive integer n such that

$$(1.1) \quad n \geq \text{ord } P.$$

Then $\varphi'/\varphi = n(\psi'/\psi)$ is algebraic over K .

Received by the editors June 24, 1985.

1980 *Mathematics Subject Classification.* Primary 12H05, 13B20, 13J05, 34A30.

¹This paper was presented in a special session on ordinary differential equations at the 91st annual meeting of the AMS on January 10, 1985 at Anaheim, California.

²Both authors were supported in part by grants from the National Science Foundation.

Theorem B gives the following two corollaries:

COROLLARY C. *In case K is the field of rational functions of x with coefficients in $\mathbf{C}$, if φ and ψ satisfy the hypotheses of Theorem B, then $\varphi'/\varphi = n(\psi'/\psi)$ is an algebraic function of x .*

COROLLARY D. *In case K is the quotient field of $\mathbf{C}\{x\}$ (the ring of convergent power series in x with coefficients in $\mathbf{C}$), if φ and ψ are formal power series in x with coefficients in $\mathbf{C}$, and if φ and ψ satisfy the hypotheses of Theorem B, then φ and $\psi \in \mathbf{C}\{x\}$ (i.e., φ and ψ are convergent).*

The following examples illustrate these results.

EXAMPLE 1. The power series $\{\cos x\}^{1/n}$ and $J_\nu(x)^{1/n}$ ($n \geq 2$) do not satisfy any linear ordinary differential equation with polynomial coefficient (cf. also [1]).

EXAMPLE 2. The divergent power series $\{\sum_{m=0}^\infty m!x^m\}^{1/n}$ ($n \geq 2$) do not satisfy any linear ordinary differential equation with coefficients in $\mathbf{C}\{x\}$.

2. A fundamental lemma. The following two important steps were utilized in our previous paper [2].

(1) Set

$$(2.1) \quad P = \sum_{h=0}^{m+1} p_h D^h, \quad Q = \sum_{h=0}^{n+1} q_h D^h,$$

where m and $n \in N$; p_h and $q_h \in K$; in particular,

$$(2.2) \quad p_{m+1} \neq 0, \quad q_{n+1} \neq 0.$$

For an element of f of K , let us set

$$\hat{f} = \sum_{h=0}^{\infty} \frac{f^{(h)}}{h!} x^h \in K[[x]],$$

where $K[[x]]$ is the ring of formal power series in x with coefficients in K . Then, $T(f) = \hat{f}$ defines an injective homomorphism of rings $T: K \rightarrow K[[x]]$ such that $T(f') = dT(f)/dx$. Corresponding to two operators P and Q of (2.1), let us consider two operators

$$\hat{P} = \sum_{h=0}^{m+1} \hat{p}_h (d/dx)^h, \quad \hat{Q} = \sum_{h=0}^{n+1} \hat{q}_h (d/dx)^h.$$

We assume that φ is an element of a differential field extension of K . Denote this extension by $\tilde{K}$. Then, $P(\varphi) = 0$ and $Q(1/\varphi) = 0$ imply, respectively, that the formal power series

$$\hat{\varphi} = \sum_{h=0}^{\infty} \frac{\varphi^{(h)}}{h!} x^h \in \tilde{K}[[x]]$$

satisfies $\hat{P}(\hat{\varphi}) = 0$ and $\hat{Q}(1/\hat{\varphi}) = 0$.

Observe that (2.2) implies $1/\hat{p}_{m+1} \in K[[x]]$ and $1/\hat{q}_{n+1} \in K[[x]]$. Therefore

$$(2.3) \quad y = \hat{\varphi}/\varphi = 1 + \sum_{h=1}^{\infty} \frac{\varphi^{(h)}}{\varphi h!} x^h$$

satisfies the differential equation

$$(2.4) \quad y^{(m+1)} + \sum_{h=0}^m \frac{\hat{p}_h}{\hat{p}_{m+1}} y^{(h)} = 0,$$

and $u = \varphi/\hat{\varphi}$ satisfies the differential equation

$$(2.5) \quad u^{(n+1)} + \sum_{h=0}^n \frac{\hat{q}_h}{\hat{q}_{n+1}} u^{(h)} = 0.$$

In this manner, the general case was reduced to the case of formal power series.

(2) Let k be a field. We denote by $k[y_1, \dots, y_p]$ the ring of polynomials in p indeterminates $y_1, \dots, y_p$ with coefficients in k . For $F \in k[y_1, \dots, y_p]$ we set

$$(2.6) \quad w(F) = \deg_t F(\alpha_1 t, \alpha_2 t^2, \dots, \alpha_p t^p),$$

where we regard $F(\alpha_1 t, \alpha_2 t^2, \dots, \alpha_p t^p)$ as a polynomial in t whose coefficients are polynomials in $\alpha_1, \dots, \alpha_p$. The following lemma was the fundamental algebraic tool in the proof of Theorem A.

LEMMA E. *Let n and p be positive integers, and let $F_1, \dots, F_p \in k[y_1, \dots, y_p]$. Assume that*

- (i) $w(F_p - y_p^n) < np$;
- (ii) $w(F_j) = nj$ ($j = 1, \dots, p-1$);
- (iii) $w(F_j(y_1, \dots, y_j, 0, \dots, 0) - y_j^n) < nj$ ($j = 1, \dots, p-1$).

Then, the system of algebraic equations

$$(2.7) \quad F_j(y_1, \dots, y_p) = 0 \quad (j = 1, \dots, p)$$

admits only a finite number of solutions in any field extension of k and these solutions are algebraic over k .

3. A lemma on differential equations. Let k be a field of characteristic zero, and let $k[[x]]$ be the ring of formal power series in x with coefficients in k . Then a linear ordinary differential equation

$$(3.1) \quad y^{(m+1)} + \sum_{h=0}^m a_h(x) y^{(h)} = 0 \quad (a_h \in k[[x]])$$

admits a canonical basis of $m+1$ solutions of the form

$$(3.2) \quad f_j = \frac{1}{j!} x^j + \sum_{h=m+1}^{\infty} f_{j,h} x^h \quad (j = 0, 1, \dots, m),$$

where $f_{j,h} \in k$.

Let $\tilde{k}$ be a field extension of k . If a formal power series

$$(3.3) \quad \varphi = \sum_{h=0}^m c_h x^h$$

with coefficients in $\tilde{k}$ satisfies (3.1), then we have

$$(3.4) \quad \varphi = \sum_{j=0}^m (j!) c_j f_j.$$

Let us consider another linear ordinary differential equation:

$$(3.5) \quad u^{(p+1)} + \sum_{h=0}^p b_h(x) u^{(h)} = 0 \quad (b_h \in k[[x]]).$$

This equation also has a canonical basis of $p+1$ solutions of the form

$$(3.6) \quad g_j = \frac{1}{j!} x^j + \sum_{h=p+1}^{\infty} g_{j,h} x^h \quad (j = 0, 1, \dots, p),$$

where $g_{j,h} \in k$.

As in our previous paper [2], in order to prove Theorem B it suffices to prove the following result:

LEMMA F. *There are at most a finite number of solutions of the form*

$$(3.7) \quad \begin{cases} y = f_0 + \sum_{j=1}^m \alpha_j f_j & (\alpha_j \in \tilde{k}), \\ u = g_0 + \sum_{j=1}^p \beta_j g_j & (\beta_j \in \tilde{k}) \end{cases}$$

of (3.1) and (3.5) respectively such that

$$(3.8) \quad y = u^n$$

if

$$(3.9) \quad n \geq m+1.$$

For these solutions, the coefficients $\alpha_1, \dots, \alpha_m, \beta_1, \dots, \beta_p$ are algebraic over k .

PROOF. Set $y = \sum_{h=0}^{\infty} \lambda_h x^h$ and $u = \sum_{h=0}^{\infty} \mu_h x^h$, where λ_h and $\mu_h \in \tilde{k}$. Then

$$(3.10) \quad \begin{cases} \lambda_0 = 1, & \mu_0 = 1, \\ \lambda_j = (1/j!) \alpha_j & (j = 1, \dots, m), \\ \mu_j = (1/j!) \beta_j & (j = 1, \dots, p), \\ \lambda_h = f_{0,h} + \sum_{j=1}^m (j!) f_{j,h} \lambda_j & (h \geq m+1), \\ \mu_h = g_{0,h} + \sum_{j=1}^p (j!) g_{j,h} \mu_j & (h \geq p+1). \end{cases}$$

In order that

$$(3.11) \quad \sum_{h=0}^{\infty} \lambda_h x^h = \left(\sum_{h=0}^{\infty} \mu_h x^h \right)^n,$$

it is necessary and sufficient that

$$(3.12) \quad \lambda_h = \sum_{h_1 + \dots + h_n = h, h_j \geq 0} \mu_{h_1} \cdots \mu_{h_n} \quad \text{for } h \geq 0.$$

Let us define $w(F)$ for $F \in k[\mu_1, \dots, \mu_p]$ in the same way as (2.6), and let us make the following observations:

(1) Utilizing the last formulas of (3.10), we regard μ_h ($h \geq p+1$) as elements of $k[\mu_1, \dots, \mu_p]$. Then

$$(3.13) \quad w(\mu_h) \leq p < h \quad \text{for } h \geq p+1.$$

(2) Utilizing $\lambda_h = \sum \mu_{h_1} \cdots \mu_{h_n}$ for $h = 1, \dots, m$ (cf. (3.12)), we can regard $\lambda_1, \dots, \lambda_m$ as elements of $k[\mu_1, \dots, \mu_p]$. Then

$$(3.14) \quad w(\lambda_j) = j \quad (j = 1, \dots, m).$$

(3) Utilizing the formulas for λ_h ($h \geq m+1$) of (3.10), we can regard λ_h ($h \geq m+1$) as elements of $k[\mu_1, \dots, \mu_p]$. Then

$$(3.15) \quad w(\lambda_h) \leq m < h \quad \text{for } h \geq m+1.$$

Now, regarding (3.12) for $h \geq m+1$ as a system of algebraic equations on $\mu_1, \dots, \mu_p$, let us consider the system of p equations

$$(3.16) \quad \lambda_h - \sum_{h_1 + \dots + h_n = h, h_j \geq 0} \mu_{h_1} \cdots \mu_{h_n} = 0$$

where $h = n, 2n, \dots, pn$. Set

$$(3.17) \quad F_j(\mu_1, \dots, \mu_p) = \lambda_{jn} - \sum_{h_1 + \dots + h_n = nj} \mu_{h_1} \cdots \mu_{h_n} \quad (j = 1, \dots, p).$$

Note that, if (3.9) is satisfied, we have

$$(3.18) \quad w(\lambda_{jn}) < nj \quad (j = 1, \dots, p)$$

(cf. (3.15)). Therefore,

$$(3.19) \quad w(F_j) \leq nj \quad (j = 1, \dots, p)$$

if (3.9) is satisfied. Furthermore, since

$$F_j(\mu_1, \dots, \mu_j, 0, \dots, 0) = \lambda_{jn} - \sum \mu_{h_1} \cdots \mu_{h_n}$$

if $\mu_{j+1} = 0, \dots, \mu_p = 0$, where the sum is over all n -tuples $(h_1, \dots, h_n)$ such that $h_1 + \dots + h_n = nj$, $h_j \geq 0$, it follows that

$$(3.20) \quad w(F_j(\mu_1, \dots, \mu_j, 0, \dots, 0) - \mu_j^n) < nj \quad (j = 1, \dots, p).$$

Thus applying Lemma E to (3.16) we can complete the proof of Lemma F, and thus Theorem B.

Using different methods, our results have been substantially generalized in [3].

REFERENCES

1. R. P. Stanley, *Differentiably finite power series*, European J. Combin. **1** (1980), 175–188.
2. W. A. Harris, Jr. and Y. Sibuya, *The reciprocals of solutions of linear ordinary differential equations*, IMA Preprint Series No. 72, Univ. of Minnesota, May 1984; Adv. in Math. (to appear).
3. S. Sperber, *On solutions of differential equations which satisfy certain algebraic relations*, Pacific J. of Math. (to appear).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN CALIFORNIA, LOS ANGELES, CALIFORNIA 90089-1113

SCHOOL OF MATHEMATICS, UNIVERSITY OF MINNESOTA, MINNEAPOLIS, MINNESOTA 55455