A NEW INEQUALITY FOR COMPLEX-VALUED POLYNOMIAL FUNCTIONS

THEMISTOCLES M. RASSIAS

ABSTRACT. Let $f_1, f_2, \dots, f_n: \mathbb{C} \rightarrow \mathbb{C}$ be complex-valued polynomial functions of degrees $d_1, d_2, \dots, d_n$, respectively, of a complex variable z . Then

$$M_{f_1} M_{f_2} \cdots M_{f_n} \geq M_{f_1 f_2 \cdots f_n} \geq k M_{f_1} M_{f_2} \cdots M_{f_n}$$

where

$$k = \left(\sin \frac{2}{n} \frac{\pi}{8d_1} \right)^{d_1} \left(\sin \frac{2}{n} \frac{\pi}{8d_2} \right)^{d_2} \cdots \left(\sin \frac{2}{n} \frac{\pi}{8d_n} \right)^{d_n}.$$

Let $\mathbb{C}$ be the field of complex numbers and $f: \mathbb{C} \rightarrow \mathbb{C}$ be a complex-valued polynomial function of a complex variable z . Define $M_f = \max_{|z|=1} |f(z)|$.

THEOREM. Let $f_1, f_2, \dots, f_n: \mathbb{C} \rightarrow \mathbb{C}$ be polynomial functions of degrees $d_1, d_2, \dots, d_n$, respectively, of a complex variable z . Then

$$M_{f_1} \cdot M_{f_2} \cdots M_{f_n} \geq M_{f_1 \cdot f_2 \cdots f_n} \geq k \cdot M_{f_1} \cdot M_{f_2} \cdots M_{f_n}$$

where

$$(1) \quad k = \left(\sin \frac{2}{n} \frac{\pi}{8d_1} \right)^{d_1} \left(\sin \frac{2}{n} \frac{\pi}{8d_2} \right)^{d_2} \cdots \left(\sin \frac{2}{n} \frac{\pi}{8d_n} \right)^{d_n}.$$

PROOF. Of course the left inequality is obvious.

Let $f: \mathbb{C} \rightarrow \mathbb{C}$ be an arbitrary polynomial function of degree d with leading coefficient one, among the given functions $\{f_i\}$. Then we can factor $f(z)$ in the form

$$f(z) = (z - z_1)(z - z_2) \cdots (z - z_d),$$

so that

$$(2) \quad \begin{aligned} M_f &= \max_{|z|=1} |f(z)| = \max_{|z|=1} |z - z_1| |z - z_2| \cdots |z - z_d| \\ &\leq (1 + |z_1|)(1 + |z_2|) \cdots (1 + |z_d|). \end{aligned}$$

We will prove that for each $\varepsilon > 0$ and $n > 0$, we have for most θ that

$$(3) \quad |f(e^{i\theta})| = |e^{i\theta} - z_1| |e^{i\theta} - z_2| \cdots |e^{i\theta} - z_d| \geq M_f \cdot \left(\sin \frac{2}{n} \frac{\pi}{8d + \varepsilon} \right)^d.$$

However by (2), we see that it is enough to show that

$$(4) \quad \frac{|e^{i\theta} - z_1| |e^{i\theta} - z_2| \cdots |e^{i\theta} - z_d|}{(1 + |z_1|)(1 + |z_2|) \cdots (1 + |z_d|)} \geq \left(\sin \frac{2}{n} \frac{\pi}{8d + \varepsilon} \right)^d$$

Received by the editors June 11, 1984 and, in revised form, June 12, 1985.

1980 *Mathematics Subject Classification.* Primary 30A10; Secondary 30C10.

Key words and phrases. Complex polynomial function, connected subset, length, inequalities.

for all values of θ in the complement of the union U_d of intervals on $[-\pi, \pi]$ with total length $l(U_d) \leq (2/n)8d\pi/(8d + \varepsilon)$. Let z be an arbitrary element in $\{z_i\}$ and set $z = re^{ia}$, $r > 0$, and $\phi = (2/n)\pi/(8d + \varepsilon)$. Then for θ satisfying the inequality $2\phi \leq |\theta| \leq \pi - 2\phi$, we have $(\rho = 2r/(r^2 + 1) < 1)$

$$\begin{aligned} \frac{|1 + re^{i\theta}|^2}{(1 + r)^2} &= \frac{1 + r^2 + 2r \cos \theta}{1 + r^2 + 2r} \geq \frac{1 + r^2 - 2r \cos 2\phi}{1 + r^2 + 2r} \\ &= \frac{1 - (2r/(1 + r^2)) \cos 2\phi}{1 + (2r/(1 + r^2))} = \frac{1 - \rho \cos 2\phi}{1 + \rho} \geq \frac{1 - \cos 2\phi}{2} = \sin^2 \phi. \end{aligned}$$

Thus $|1 + re^{i\theta}|/(1 + r) \geq \sin \phi$ for all values of θ in the complement of the union V of intervals with total length $l(V) \leq 8\phi$. Therefore, on a similar set we have that

$$(5) \quad |e^{i\theta} - re^{ia}|/(1 + |re^{ia}|) \geq \sin \phi$$

and so (4) follows. Applying the estimate (2) for each polynomial function f_i of degree d_i , we obtain n inequalities of the form (3) (with $f = f_1, f_2, \dots, f_n$), with each holding for all values of θ in the complement of the union U_{d_i} of intervals with total length

$$l(U_{d_i}) \leq (2/n)8d_i\pi/(8d_i + \varepsilon),$$

respectively, for $i = 1, 2, \dots, n$.

We add this for all n , and deduce that all n inequalities (3) hold for some

$$\beta \in [-\pi, \pi] - \bigcup_{i=1}^n U_{d_i}.$$

Therefore

$$\begin{aligned} M_{f_1 \cdot f_2 \cdots f_n} &\geq |f_1(e^{i\beta}) \cdot f_2(e^{i\beta}) \cdots f_n(e^{i\beta})| \\ (6) \quad &= |f_1(e^{i\beta})| |f_2(e^{i\beta})| \cdots |f_n(e^{i\beta})| \geq M_{f_1} \cdot M_{f_2} \cdots M_{f_n} \\ &\quad \cdot \left(\sin \frac{2}{n} \frac{\pi}{8d_1 + \varepsilon} \right)^{d_1} \left(\sin \frac{2}{n} \frac{\pi}{8d_2 + \varepsilon} \right)^{d_2} \cdots \left(\sin \frac{2}{n} \frac{\pi}{8d_n + \varepsilon} \right)^{d_n}. \end{aligned}$$

Taking the limit as $\varepsilon \rightarrow 0$ (6) implies

$$(7) \quad M_{f_1 f_2 \cdots f_n} \geq k \cdot M_{f_1} M_{f_2} \cdots M_{f_n}$$

where k satisfies (1).

REMARK. See [1–5] for other results and some research problems concerning inequalities on the derivative of a complex-valued function.

ACKNOWLEDGEMENT. I would like to express my thanks to the referee for helpful suggestions.

REFERENCES

1. Th. M. Rassias, *On the derivative of a complex valued function*, Bull. Inst. Math. Acad. Sinica **12** (1984), 423–425.
2. G. M. Rassias, J. M. Rassias, and Th. M. Rassias, *A counterexample to a conjecture by P. Erdős*, Proc. Japan Acad. Sci. Ser. A **53** (1977), 119–121.
3. S. Smale, *On the efficiency of algorithms of analysis*, Bull. Amer. Math. Soc. (N.S.) **13** (1985), 87–121.

4. ———, *The fundamental theorem of algebra and complexity theory*, Bull. Amer. Math. Soc. (N.S.) **4** (1981), 1–36.
5. M. J. Todd, *Polynomial expected behavior of a pivoting algorithm for linear complementary and linear programming problems*, Technical Rep. No. 595, School of Operations Research and Industrial Engineering, Cornell Univ., Ithaca, N. Y., 1983.

Current address: 4 Zagoras Street, Paradissos, Amaroussion, 15125 Athens, Greece