SHORTER NOTES

The purpose of this department is to publish very short papers of unusually polished character, for which there is no other outlet.

RANGES OF JOINT LAPLACE-FOURIER TRANSFORMS

DANIEL J. GRUBB

ABSTRACT. If $f \in L^1([0, \infty) \times \mathbf{R})$ and $\hat{f}(z, s) = \int_{-\infty}^{\infty} \int_0^{\infty} f(u, v) e^{iuz} e^{isv} du dv$, where $z \in H = \{z \in \mathbf{C} : \text{Im } z \geq 0\}$, $s \in \mathbf{R}$, then $\hat{f}(H \times \mathbf{R}) \cup \{0\} = \hat{f}(\mathbf{R} \times \mathbf{R}) \cup \{0\}$.

The proof is based on the following facts: (a) $\hat{f}$ is continuous on $H \times \mathbf{R}$, (b) for fixed $s \in \mathbf{R}$, $\hat{f}(z, s)$ is holomorphic on the interior of H , and (c) $\hat{f}$ vanishes at infinity on $H \times \mathbf{R}$. Since these facts are well known [1], I only check (c).

Define $e_x(u, v) = e^{-xu}$ for $x \in \mathbf{R}$ and $(u, v) \in [0, \infty) \times \mathbf{R}$. Fix $\varepsilon > 0$. By dominated convergence, there is an $R > 0$ such that $\|e_R f\|_1 < \varepsilon$. Then, for $\text{Im } z \geq R$, we have

$$|\hat{f}(z, s)| \leq \int_{-\infty}^{\infty} \int_0^{\infty} |f(u, v)| e_{\text{Im } z}(u, v) du dv \leq \|e_R f\|_1 < \varepsilon.$$

Again, by dominated convergence, the map $x \mapsto e_x f$ from $[0, R]$ into $L^1([0, \infty) \times \mathbf{R})$ is continuous. By compactness, there are $0 \leq x_1 < x_2 < \dots < x_n \leq R$ such that, for any $0 \leq x \leq R$, there is an x_j with $\|e_x f - e_{x_j} f\|_1 < \varepsilon/2$. Since $\widehat{e_{x_j} f}$ is the Fourier transform, $\widehat{e_{x_j} f}$ vanishes at infinity on $\mathbf{R} \times \mathbf{R}$. Thus there is a compact $K \times L \subseteq \mathbf{R} \times \mathbf{R}$ such that $(r, s) \notin K \times L$ implies $|\widehat{e_{x_j} f}(r, s)| < \varepsilon/2$ for all x_j . Then, if $0 \leq y = \text{Im } z \leq R$ and $(\text{Re } z, s) \notin K \times L$, we have

$$\begin{aligned} |\hat{f}(z, s)| &= |\widehat{e_y f}(\text{Re } z, s)| \leq |\widehat{e_y f}(\text{Re } z, s) - \widehat{e_{x_j} f}(\text{Re } z, s)| + |\widehat{e_{x_j} f}(\text{Re } z, s)| \\ &\leq \|e_y f - e_{x_j} f\|_1 + |\widehat{e_{x_j} f}(\text{Re } z, s)| < \varepsilon/2 + \varepsilon/2 = \varepsilon \end{aligned}$$

for appropriate x_j . Thus $(z, s) \notin (K \times [0, R]) \times L$ implies that $|\hat{f}(z, s)| < \varepsilon$.

Evidently, we have $\hat{f}(\mathbf{R} \times \mathbf{R}) \cup \{0\} \subseteq \hat{f}(H \times \mathbf{R}) \cup \{0\}$. To prove the reverse inclusion, fix $r \notin \hat{f}(\mathbf{R} \times \mathbf{R}) \cup \{0\}$. Since $\hat{f}$ vanishes at infinity on $H \times \mathbf{R}$, there is an $s_0 \in \mathbf{R}$ such that $r \notin \hat{f}(H \times \{s_0\})$. Now set $A = \{s \in \mathbf{R} : r \notin \hat{f}(H \times \{s\})\}$. Then A is open. In fact, if $s \in A$, $\hat{f}(H \times \{s\})$ is bounded away from r . By uniform continuity, there is an $\varepsilon > 0$ such that $|s - t| < \varepsilon$ implies $r \notin \hat{f}(H \times \{t\})$, i.e., $t \in A$.

The complement of A is also open since, if $s \notin A$, there is a $z_0 \in H$ with $\hat{f}(z_0, s) = r$. By assumption on r , $\text{Im } z_0 > 0$. Since $z \mapsto \hat{f}(z, s)$ is nonconstant and

Received by the editors November 13, 1985.

1980 *Mathematics Subject Classification*. Primary 44A30.

holomorphic, there is a circle C contained in the interior of H such that $\hat{f}(z, s)$ is never r for z in C . Then $a = \inf\{|\hat{f}(z, s) - r| : z \in C\} > 0$. By uniform continuity of $\hat{f}$, there is an $\varepsilon > 0$ such that $|s - t| < \varepsilon$ implies $|\hat{f}(z, s) - \hat{f}(z, t)| < a$ for all z in C . By Rouché's Theorem, $\hat{f}(z, t) - r$ has as many roots inside C as does $\hat{f}(z, s) - r$, i.e., at least one. Then $|s - t| < \varepsilon$ implies $t \notin A$.

Since $\mathbf{R}$ is connected and $s_0 \in A$, A is equal to $\mathbf{R}$, and so r does not belong to $\hat{f}(H \times \mathbf{R}) \cup \{0\}$.

REMARK. The foregoing proof is valid with $[0, \infty[\times \mathbf{R}$ replaced by $[0, \infty[\times \mathbf{R}^n$, and $H \times \mathbf{R}$ by $H \times \mathbf{R}^n$, etc.

Rewording the theorem gives

COROLLARY. For $f \in L^1([0, \infty[\times \mathbf{R})$, the spectrum of f as an element of the Banach algebra $L^1([0, \infty[\times \mathbf{R})$ is the same as the spectrum of f as an element of the Banach algebra $L^1(\mathbf{R} \times \mathbf{R})$.

REMARK. Note that the maximal ideal space of $L^1([0, \infty[\times \mathbf{R})$ is (homeomorphic to) $H \times \mathbf{R}$ and that of $L^1(\mathbf{R} \times \mathbf{R})$ is $\mathbf{R} \times \mathbf{R}$ (see [1]). Thus, the above corollary provides a Banach algebra B and a closed subalgebra A such that

(i) the spectrum of any element of A as an element of A coincides with the spectrum as an element of B , while

(ii) not every multiplicative linear functional on A extends to a multiplicative linear functional on B .

For another example where the range of a Fourier type transform is attained on a small portion of the domain see [2].

BIBLIOGRAPHY

1. R. F. Arens and I. M. Singer, *Generalized analytic functions*, Trans. Amer. Math. Soc. **81** (1956), 379–393, **MR 17**, 1226.
2. Edwin Hewitt and J. H. Williamson, *Note on absolutely convergent Dirichlet series*, Proc. Amer. Math. Soc. **8** (1957), 863–868, **MR 19**, 851.

DEPARTMENT OF MATHEMATICS, KANSAS STATE UNIVERSITY, MANHATTAN, KANSAS
66506