CUBE SLICING IN $\mathbf{R}^n$

KEITH BALL

ABSTRACT. We prove that every $(n - 1)$ -dimensional section of the unit cube in $\mathbf{R}^n$ has volume at most $\sqrt{2}$. This upper bound is clearly best possible.

1. Introduction. Hensley [3] showed that if $Q_n = [-\frac{1}{2}, \frac{1}{2}]^n \subset \mathbf{R}^n$ is the central unit cube in $\mathbf{R}^n$, and H is an $(n - 1)$ -dimensional subspace of $\mathbf{R}^n$, then $|H \cap Q_n|$ lies between 1 and 5. (Here and henceforth $|\cdot|$ will denote the usual volume in a Euclidean space.) Hensley conjectured that the volume is at most $\sqrt{2}$. Besides proving this conjecture, we shall provide a somewhat simpler proof that the volume is at least 1.

The bound 5, obtained by Hensley, can be improved in a more general setting. In [1], we show that there is an absolute constant, c , so that for all n , every n -dimensional Banach space has a representation on $\mathbf{R}^n$, with unit ball, B , say, with the property that if H and K are $(n - 1)$ -dimensional subspaces of $\mathbf{R}^n$, $|H \cap B| \leq c|K \cap B|$. We can take c to be $\sqrt{6}$.

Let H be an $(n - 1)$ -dimensional subspace of $\mathbf{R}^n$ and $a = (a_1, \dots, a_n) \in \mathbf{R}^n$ be the unit vector normal to H . Assume $n \geq 2$ and $|a_i| < 1$, for all i . Define $f = f_H: \mathbf{R} \rightarrow [0, \infty)$ by $f(r) = |(H + ra) \cap Q_n|$, so $f(r)$ is the volume of the intersection of Q_n with a hyperplane parallel to H , at distance $|r|$ from H . The assumption that $n \geq 2$, $|a_i| < 1$ for all i ensures that H is not parallel to a face of Q_n and hence that f is continuous.

As in [3] we shall use probabilistic methods to obtain an expression for $|H \cap Q_n| = f(0)$. Let $X_1, \dots, X_n$ be independent random variables, each uniformly distributed on $[-\frac{1}{2}, \frac{1}{2}]$, with respect to a probability, P . Then the random vector $(X_1, \dots, X_n)$ induces the usual measure on Q_n , and, since f is continuous,

$$\begin{aligned} f(r) &= \lim_{s \rightarrow 0} \frac{1}{2s} \int_{r-s}^{r+s} f(t) dt \\ &= \lim_{s \rightarrow 0} \frac{1}{2s} \left| \left\{ x \in Q_n : \left| \sum_{i=1}^n a_i x_i - r \right| \leq s \right\} \right| \\ &= \lim_{s \rightarrow 0} \frac{1}{2s} P \left(\left| \sum_{i=1}^n a_i X_i - r \right| \leq s \right), \end{aligned}$$

Received by the editors July 10, 1985.

1980 *Mathematics Subject Classification*. Primary 52A20, 52A40, 60E05.

©1986 American Mathematical Society
0002-9939/86 \$1.00 + \$.25 per page

so f is the continuous probability density function of the random variable $X = \sum a_i X_i$.

It was observed by Hensley that f attains its maximum at 0. This is also an immediate consequence of the Brunn-Minkowski inequality, namely,

$$|\tfrac{1}{2}A + \tfrac{1}{2}B| \geq \left(\tfrac{1}{2}|A|^{1/k} + \tfrac{1}{2}|B|^{1/k} \right)^k$$

for convex sets A and B in $\mathbf{R}^k$. For, taking $k = n - 1$,

$$A = H \cap (Q_n + ra) \subset H = \mathbf{R}^k, \quad \text{and} \quad B = H \cap (Q_n - ra) = -A,$$

the convexity of Q_n gives

$$H \cap Q_n \supset \tfrac{1}{2}(A + B) = \tfrac{1}{2}(A - A);$$

so using the Brunn-Minkowski inequality gives

$$f(0) = |H \cap Q_n| \geq |\tfrac{1}{2}(A - A)| \geq |A| = f(r).$$

The upper bound, $\sqrt{2}$, will thus apply to all sections of Q_n by hyperplanes, not merely those by subspaces.

Our principal aim is to prove best possible bounds on $\|f\|_\infty = f(0) = |H \cap Q_n|$ with $(a_i)_1^n$, $(X_i)_1^n$ and f as above.

2. The lower bound. The lower bound will be an immediate consequence of the following simple probabilistic inequality, which compares a random variable with a uniformly distributed random variable of the same L_p -norm.

LEMMA 1. *Let Y be a random variable with a probability density function $g: \mathbf{R} \rightarrow [0, \infty)$. Then for $p > 0$,*

$$\|g\|_\infty \|Y\|_p \geq \tfrac{1}{2}(p+1)^{-1/p}.$$

Equality holds in the above if and only if Y is uniformly distributed on some subinterval $[-t, t]$ of $\mathbf{R}$.

PROOF. Without loss of generality, we may assume that X is symmetric. Then setting

$$G(x) = \int_0^x g(t) dt \quad \text{for } x \geq 0,$$

we have $G(0) = 0$, $G(\infty) = 1/2$, and $G(x) \leq x\|g\|_\infty$, so

$$\begin{aligned} 2^{-p} &= 2(G(\infty))^{p+1} = 2 \int_0^\infty (G(x)^{p+1})' dx \\ &= 2(p+1) \int_0^\infty g(x) G(x)^p dx \\ &\leq (p+1) \|g\|_\infty^p \cdot 2 \int_0^\infty g(x) \cdot x^p dx \\ &= (p+1) \|g\|_\infty^p \|Y\|_p^p. \end{aligned}$$

There is equality if and only if $g(t) = \|g\|_\infty$ or 0 almost everywhere.

THEOREM 2. *Every section of the unit cube, Q_n , in $\mathbf{R}^n$ by an $(n-1)$ -dimensional subspace, H , has volume at least 1. The volume is 1 if and only if H is parallel to a face of the cube.*

PROOF. Plainly it suffices to show that, with $(a_i)_1^n, (X_i)_1^n$ as above ($|a_i| < 1$ for all i) and f the continuous probability density function of $X = \sum a_i X_i$, we have $|H \cap Q_n| = f(0) > 1$. Since the X_i 's are independent and identically distributed,

$$EX^2 = \left(\sum a_i^2 \right) EX_1^2 = EX_1^2 = 1/12.$$

So $\|X\|_2 = 1/2\sqrt{3}$, and the result follows by applying Lemma 1 with $p = 2$ and $g = f$.

We should remark that Theorem 2 was conjectured by A. Good, as was the following extension to subspaces of arbitrary dimension. Let U be a subspace of $\mathbf{R}^n$: then $|U \cap Q_n| \geq 1$. This extension was proved by Vaaler [4].

3. The upper bound. The proof of the principal result depends on the following lemma, whose proof we postpone until after the theorem itself.

LEMMA 3.

$$\frac{1}{\pi} \int_{-\infty}^{\infty} \left| \frac{\sin t}{t} \right|^p dt \leq \frac{\sqrt{2}}{\sqrt{p}} \quad \text{if } p \geq 2,$$

and there is equality if and only if $p = 2$.

THEOREM 4. *Every section of the unit cube, Q_n , by an $(n-1)$ -dimensional subspace, H , has volume at most $\sqrt{2}$. This upper bound is attained if and only if H contains an $(n-2)$ -dimensional face of Q_n .*

PROOF. Let $(a_i)_1^n$ be as above and assume $a_i \geq 0$ for all i . Suppose first that for some i , $a_i > 1/\sqrt{2}$; say $a_1 > 1/\sqrt{2}$. Consider $H \cap C_n$, where C_n is the "cylinder" $\{x \in \mathbf{R}^n: |x_i| \leq 1/2, 2 \leq i \leq n\}$. Let $T: \mathbf{R}^n \rightarrow \mathbf{R}^n$ be the orthogonal projection onto the subspace $\{x \in \mathbf{R}^n: x_1 = 0\}$, and S the orthogonal projection onto H . Then $T(H \cap C_n)$ is the unit cube in $\mathbf{R}^{n-1}$ and

$$|H \cap C_n| = |T(H \cap C_n)| \frac{\|Se\|}{\|TSe\|} = \frac{\|Se\|}{\|TSe\|},$$

where e is the unit vector $(1, 0, \dots, 0)$ and $\|\cdot\|$ is the usual Euclidean norm in $\mathbf{R}^n$.

Since $Se = (1 - a_1^2, -a_1a_2, \dots, -a_1a_n)$ and $TSe = (0, -a_1a_2, \dots, -a_1a_n)$, we obtain

$$|H \cap C_n| = \frac{\left((1 - a_1^2)^2 + a_1^2 \sum_{i=2}^n a_i^2 \right)^{1/2}}{\left(a_1^2 \sum_{i=2}^n a_i^2 \right)^{1/2}} = \frac{1}{a_1} < \sqrt{2}.$$

Hence $|H \cap Q_n| \leq |H \cap C_n| < \sqrt{2}$.

Assume otherwise that $a_i \leq 1/\sqrt{2}$ for all i . Since, if some a_i is zero, the problem reduces to that in $\mathbf{R}^{n-1}$, we may assume inductively that $a_i > 0$ for all i , and with this assumption the case of equality will follow if we show that $|H \cap Q_n| < \sqrt{2}$ unless $n = 2$ and $a_1 = a_2 = 1/\sqrt{2}$. With $(X_i)_1^n$ independent random variables uniformly distributed on $[-\frac{1}{2}, \frac{1}{2}]$ and f the continuous p.d.f. of $X = \sum a_i X_i$ as

above, the random variable $a_i X_i$ has characteristic function

$$\phi_i(t) = (2 \sin \tfrac{1}{2} a_i t) / a_i t,$$

and so the characteristic function of X is

$$\phi(t) = \prod_1^n \phi_i(t) = \prod_1^n \frac{2 \sin \frac{1}{2} a_i t}{a_i t} = \int_{-\infty}^{\infty} \cos xt \cdot f(x) dx.$$

By the standard Fourier inversion formula,

$$\begin{aligned} |H \cap Q_n| &= f(0) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \prod_1^n \frac{2 \sin \frac{1}{2} a_i t}{a_i t} dt \\ &= \frac{1}{\pi} \int_{-\infty}^{\infty} \prod_1^n \frac{\sin a_i t}{a_i t} dt \leq \frac{1}{\pi} \int_{-\infty}^{\infty} \prod_1^n \left| \frac{\sin a_i t}{a_i t} \right| dt. \end{aligned}$$

Set $p_i = a_i^{-2}$ so that $p_i \geq 2$ for all i and $\sum p_i^{-1} = 1$. Then by Hölder's inequality with n exponents,

$$\begin{aligned} |H \cap Q_n| &\leq \prod_1^n \left(\frac{1}{\pi} \int_{-\infty}^{\infty} \left| \frac{\sin a_i t}{a_i t} \right|^{p_i} dt \right)^{1/p_i} \\ &= \prod_1^n \left(\frac{1}{a_i \pi} \int_{-\infty}^{\infty} \left| \frac{\sin t}{t} \right|^{p_i} dt \right)^{1/p_i}. \end{aligned}$$

By Lemma 3 this gives

$$|H \cap Q_n| \leq \prod_1^n \left(\frac{\sqrt{2}}{a_i \sqrt{p_i}} \right)^{1/p_i} = \prod_1^n (\sqrt{2})^{1/p_i} = \sqrt{2},$$

and there is equality only if $p_i = 2$ for all i ; i.e., $n = 2$ and $a_1 = a_2 = 1/\sqrt{2}$.

PROOF OF LEMMA 3. We divide the proof into two cases, $p \geq 4$ and $2 \leq p < 4$.

(a) $p \geq 4$. Observe that

$$\begin{aligned} \frac{\sin t}{t} &= 1 - \frac{t^2}{6} + \frac{t^4}{120} - \frac{t^6}{7!} + \frac{t^8}{9!} - \dots \\ &< 1 - \frac{t^2}{6} + \frac{t^4}{120} \quad \text{if } t^2 < 72, \\ e^{-t^2/6} &= 1 - \frac{t^2}{6} + \frac{t^4}{72} - \frac{t^6}{1296} + \frac{t^8}{6^4 \cdot 4!} - \frac{t^{10}}{6^5 \cdot 5!} + \dots \\ &> 1 - \frac{t^2}{6} + \frac{t^4}{72} - \frac{t^6}{1296} \quad \text{if } t^2 < 30. \end{aligned}$$

Since

$$\frac{t^4}{72} - \frac{t^6}{1296} - \frac{t^4}{120} = \frac{t^4}{180} - \frac{t^6}{1296} \geq 0 \quad \text{if } t^2 \leq \frac{36}{5}$$

and $36/5 \leq \pi^2$, we have

$$0 \leq (\sin t)/t \leq e^{-t^2/6} \quad \text{if } t^2 \leq 36/5.$$

Let $m = 6/\sqrt{5}$. Then

$$\begin{aligned} \frac{1}{\pi} \int_{-\infty}^{\infty} \left| \frac{\sin t}{t} \right|^p dt &< \frac{1}{\pi} \int_{-\infty}^{\infty} e^{-pt^2/6} dt + \frac{2}{\pi} \int_m^{\infty} t^{-p} dt \\ &= \frac{\sqrt{6}}{\sqrt{p} \cdot \sqrt{\pi}} + \frac{2}{\pi(p-1)m^{p-1}} \\ &\leq \frac{1}{\sqrt{p}} \left(\frac{\sqrt{6}}{\sqrt{\pi}} + \frac{4}{3\pi m^3} \right) \quad \text{for } p \geq 4 \\ &< \frac{\sqrt{2}}{\sqrt{p}}. \end{aligned}$$

(b) $2 \leq p < 4$. Setting $s = p/2 - 1$, we wish to show that

$$(1) \quad \frac{1}{\pi} \int_{-\infty}^{\infty} \left(\frac{\sin^2 t}{t^2} \right)^{1+s} dt \leq \frac{1}{\sqrt{1+s}} \quad \text{for } 0 \leq s < 1,$$

with equality if and only if $s = 0$.

For $n = 0, 1, 2, \dots$ define

$$\alpha_n = \frac{1}{\pi} \int_{-\infty}^{\infty} e^{-t^2/\pi} (1 - e^{-t^2/\pi})^n, \quad \beta_n = \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{\sin^2 t}{t^2} \left(1 - \frac{\sin^2 t}{t^2} \right)^n dt.$$

It is easily checked that

$$\beta_0 = \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{\sin^2 t}{t^2} dt = 1.$$

Now

$$\begin{aligned} \left(\frac{\sin^2 t}{t^2} \right)^{1+s} &= \frac{\sin^2 t}{t^2} \left(1 - \left(1 - \frac{\sin^2 t}{t^2} \right) \right)^s \\ &= \frac{\sin^2 t}{t^2} \sum_0^{\infty} \frac{(-1)^n s(s-1) \cdots (s-n+1)}{n!} \left(1 - \frac{\sin^2 t}{t^2} \right)^n \\ &= \frac{\sin^2 t}{t^2} \left(1 - \sum_1^{\infty} \frac{|s(s-1) \cdots (s-n+1)|}{n!} \left(1 - \frac{\sin^2 t}{t^2} \right)^n \right), \end{aligned}$$

since $(-1)^n s(s-1) \cdots (s-n+1) \leq 0$ if $n \geq 1$.

Hence by the monotone convergence theorem,

$$\begin{aligned} \frac{1}{\pi} \int_{-\infty}^{\infty} \left(\frac{\sin^2 t}{t^2} \right)^{1+s} dt &= \beta_0 - \sum_1^{\infty} \frac{|s(s-1) \cdots (s-n+1)|}{n!} \beta_n \\ &= 1 - \sum_1^{\infty} \frac{|s(s-1) \cdots (s-n+1)|}{n!} \beta_n. \end{aligned}$$

Now we may write the right side of (1) as

$$\frac{1}{\sqrt{1+s}} = \frac{1}{\pi} \int_{-\infty}^{\infty} (e^{-t^2/\pi})^{1+s} dt,$$

and analogously to the above argument, we have

$$\frac{1}{\sqrt{1+s}} = 1 - \sum_1^{\infty} \frac{|s(s-1) \cdots (s-n+1)|}{n!} \alpha_n.$$

Hence it will suffice to show that $\alpha_n < \beta_n$ for $n \in \mathbb{N}$.

The first six values of α_n and β_n (calculated using the residue theorem in the case of β_n), correct to two decimal places, are shown in Table 1.

TABLE 1

n	α_n	β_n
1	0.29	0.33
2	0.16	0.22
3	0.11	0.17
4	0.08	0.15
5	0.07	0.13
6	0.05	0.12

It is easily checked that for $n \geq 7$,

$$\frac{1}{\sqrt{2\pi} \cdot n} < \frac{1}{2\sqrt{\pi} \sqrt{n+1}} - \frac{1}{2en},$$

so to complete the proof it suffices to show that

$$(2) \quad \alpha_n \leq \frac{1}{\sqrt{2\pi} \cdot n},$$

$$(3) \quad \beta_n \geq \frac{1}{2\sqrt{\pi} \sqrt{n+1}} - \frac{1}{2en} \quad \text{for } n \geq 7.$$

We obtain these estimates as follows.

For $0 < x \leq 1$,

$$\begin{aligned} |\log x| &\geq 1 - x + \frac{1}{2}(1-x)^2 + \frac{1}{3}(1-x)^3 + \frac{1}{4}(1-x)^4 \\ &\geq 2(1-x)^2 + (1-x) - \frac{3}{2}(1-x)^2 + \frac{1}{2}(1-x)^4 \\ &= 2(1-x)^2 + \frac{1}{2}x^2(1-x)(3-x) \\ &\geq 2(1-x)^2. \end{aligned}$$

Hence

$$\begin{aligned} \alpha_n &= \frac{1}{\pi} \int_{-\infty}^{\infty} e^{-x^2/\pi} (1 - e^{-x^2/\pi})^n dx \\ &= \frac{2}{\sqrt{\pi}} \int_0^{\infty} e^{-u^2} (1 - e^{-u^2})^n du \\ &= \frac{1}{\sqrt{\pi}} \int_0^1 \frac{(1-x)^n}{\sqrt{|\log x|}} dx \leq \frac{1}{\sqrt{2\pi}} \int_0^1 (1-x)^{n-1} dx \\ &= \frac{1}{\sqrt{2\pi} \cdot n}, \quad \text{proving (2).} \end{aligned}$$

Now plainly,

$$\begin{aligned}\beta_n &= \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{\sin^2 t}{t^2} \left(1 - \frac{\sin^2 t}{t^2}\right)^n dt \\ &\geq \frac{2}{\pi} \int_1^{\infty} \frac{\sin^2 t}{t^2} \left(1 - \frac{\sin^2 t}{t^2}\right)^n dt \\ &\geq \frac{2}{\pi} \int_1^{\infty} \frac{\sin^2 t}{t^2} \left(1 - \frac{1}{t^2}\right)^n dt.\end{aligned}$$

Observe that $(1/t^2)(1 - 1/t^2)^n$ is increasing for $1 \leq t \leq \sqrt{n+1}$ and decreasing for $t \geq \sqrt{n+1}$.

Hence, if $n \geq 6$ so that $\sqrt{n+1} > 1 + \pi/2$,

$$\int_{1+\pi/2}^{\sqrt{n+1}} \frac{\sin^2 t}{t^2} \left(1 - \frac{1}{t^2}\right)^n dt \geq \int_1^{\sqrt{n+1}-\pi/2} \frac{\cos^2 t}{t^2} \left(1 - \frac{1}{t^2}\right)^n dt$$

and

$$\int_{\sqrt{n+1}}^{\infty} \frac{\sin^2 t}{t^2} \left(1 - \frac{1}{t^2}\right)^n dt \geq \int_{\sqrt{n+1}+\pi/2}^{\infty} \frac{\cos^2 t}{t^2} \left(1 - \frac{1}{t^2}\right)^n dt.$$

So

$$\begin{aligned}\int_1^{\infty} \frac{\sin^2 t}{t^2} \left(1 - \frac{1}{t^2}\right)^n dt &\geq \frac{1}{2} \int_1^{\infty} \frac{1}{t^2} \left(1 - \frac{1}{t^2}\right)^n dt - \frac{1}{2} \int_{\sqrt{n+1}-\pi/2}^{\sqrt{n+1}+\pi/2} \frac{\cos^2 t}{t^2} \left(1 - \frac{1}{t^2}\right)^n dt \\ &\geq \frac{1}{2} \int_1^{\infty} \frac{1}{t^2} \left(1 - \frac{1}{t^2}\right)^n dt - \frac{\pi}{4} \max_t \left(\frac{1}{t^2} \left(1 - \frac{1}{t^2}\right)^n \right) \\ &= \frac{1}{2} \int_0^1 (1-u^2)^n du - \frac{\pi}{4(n+1)} \left(1 - \frac{1}{n+1}\right)^n \\ &= \frac{4^n}{2(2n+1)} \binom{2n}{n}^{-1} - \frac{\pi}{4n} \left(1 - \frac{1}{n+1}\right)^{n+1} \\ &\geq \frac{\sqrt{\pi}}{4\sqrt{n+1}} - \frac{\pi}{4en}.\end{aligned}$$

Hence

$$\beta_n \geq \frac{1}{2\sqrt{\pi} \cdot \sqrt{n+1}} - \frac{1}{2en},$$

which is (3). $\square$

The series comparison technique, used in part (b), above, was employed by Haagerup in his determination of the best constants in Khintchine's inequality [2]. We are indebted to J. Lindenstrauss for pointing out the similarity between the L_p -integral to be evaluated here and those of Haagerup.

Given the identity

$$\frac{1}{\pi} \int_{-\infty}^{\infty} \left(\frac{\sin^2 t}{t^2} \right)^{1+s} dt = 1 + \sum_1^{\infty} (-1)^n \frac{s(s-1) \cdots (s-n+1)}{n!} \beta_n, \quad 0 \leq s < 1,$$

one could proceed with the proof of Lemma 3, part (b), in a number of ways. One possibility would be to consider the derivative with respect to s , as follows.

Let

$$I(s) = \frac{1}{\pi} \int_{-\infty}^{\infty} \left(\frac{\sin^2 t}{t^2} \right)^{1+s} dt, \quad J(s) = \frac{1}{\sqrt{1+s}}.$$

Then

$$\begin{aligned} I'(s) &= \frac{d}{ds} \left(1 + \sum_1^{\infty} (-1)^n \frac{s(s-1) \cdots (s-n+1)}{n!} \beta_n \right) \\ &= - \sum_1^{\infty} \frac{\beta_n}{n} + \sum_2^{\infty} s \cdot \frac{(1-s) \cdots (n-1-s)}{n!} \beta_n \cdot \sum_{k=1}^{n-1} \frac{1}{k-s} \\ &= - \sum_1^{\infty} \frac{\beta_n}{n} + \sum_2^{\infty} \frac{s}{n} \cdot \prod_{j=1}^{n-1} \left(1 - \frac{s}{j} \right) \cdot \beta_n \sum_{k=1}^{n-1} \frac{1}{k-s} \\ &\leq - \sum_1^{\infty} \frac{\beta_n}{n} + \sum_2^{\infty} \frac{s}{n} \beta_n \cdot (3 + \log(n-1)) \quad \text{for } 0 \leq s \leq \frac{1}{2}. \end{aligned}$$

A simple computation shows that $\beta_n \leq 1/\sqrt{n}$ for all n , and that the above then gives

$$I'(s) \leq - \sum_0^{\infty} \frac{\beta_n}{n} + 10s \quad \text{for } 0 \leq s \leq \frac{1}{2}.$$

Substituting the values of β_n for $1 \leq n \leq 6$ gives

$$- \sum_0^{\infty} \frac{\beta_n}{n} < -0.58,$$

so

$$I'(s) < -1/2 \leq J'(s) \quad \text{for } 0 \leq s \leq 0.008.$$

This gives $I(s) < J(s)$ for $0 < s \leq 0.008$. The fact that I and J are decreasing functions enables us to prove that $I(s) < J(s)$ for $0.008 < s \leq 1$, using finitely many iterations of the following scheme. Let $s_0 = 0.008$. Using sufficiently accurate values for β_n for sufficiently many n , find t_0 so that $I(s_0) \leq t_0 < J(s_0)$. Choose s_1 so that $J(s_1) \geq t_0$ and repeat to obtain $t_1, s_2, t_2, s_3, \dots$. For some k we obtain $s_k > 1$, and the result is proven.

4. Additional remarks. Despite the similarity of Theorem 4 to the classical Khintchine inequality, the two results do not seem to be readily deducible from one another. However, Theorem 4 and Lemma 1 together provide a simple analogue of Khintchine's inequality for $0 < p < 2$ for independent, ununiformly distributed random variables.

PROPOSITION 5. Let $X_1, \dots, X_n$ be independent random variables, uniformly distributed on $[-\frac{1}{2}, \frac{1}{2}]$ and $p > 0$. Then

$$\|\sum a_i X_i\|_p \geq \frac{\sqrt{3}}{\sqrt{2}(p+1)^{1/p}} \|\sum a_i X_i\|_2 = \frac{1}{2\sqrt{2}(p+1)^{1/p}} (\sum a_i^2)^{1/2}$$

for any sequence, $(a_i)_1^n$, of scalars.

It is natural to ask what is the maximum volume (in terms of k and n) of the intersection of Q_n with a k -dimensional subspace of $\mathbf{R}^n$. While for $k = 1$ the answer is trivially $\sqrt{n}$ and for $k = n - 1$ the answer is provided by Theorem 3, it is not clear what should be the result for general k .

This work will form part of a doctoral thesis being written under the supervision of Dr. B. Bollobás, whose constant advice and encouragement have been invaluable.

REFERENCES

1. K. M. Ball, *Sections of convex sets in $\mathbf{R}^n$* (in preparation).
2. U. Haagerup, *The best constant in the Khintchine inequality*, Studia Math. **70** (1982), 231–283.
3. D. Hensley, *Slicing the cube in $\mathbf{R}^n$ and probability*, Proc. Amer. Math. Soc. **73** (1979), 95–100.
4. J. D. Vaaler, *A geometric inequality with applications to linear forms*, Pacific J. Math. **83** (1979), 543–553.

DEPARTMENT OF PURE MATHEMATICS AND MATHEMATICAL STATISTICS, MILL LANE, CAMBRIDGE, CB2 1SB, ENGLAND