KINEMATIC FORMULAS FOR WEYL'S CURVATURE INVARIANTS OF SUBMANIFOLDS IN COMPLEX PROJECTIVE SPACE

TORU ISHIHARA¹

ABSTRACT. It is shown in [5] that Weyl's curvature invariants $k_{2p}(M)$ can be expressed by $\gamma_q \wedge F^{m-q}[M]$, where M is a $2m$ -dimensional Kähler submanifold with compact closure in a space of constant holomorphic curvature, γ_q is the q th Chern form of M and F is the Kähler form of M . In this paper, we shall show that each $\gamma_q \wedge F^{m-q}[M]$ is expressible in terms of F and $k_{2p}(M)$. Using this result, we get kinematic formulas for $k_{2p}(M)$ from Shifrin's [8] kinematic formulas for Chern classes.

1. Introduction. In this paper we rely on the notation, terminology and results in [3 and 5]. Let M be a Kähler submanifold of complex dimension m with compact closure in a space $N(\lambda)$ of constant holomorphic curvature 4λ ($\lambda > 0$). Let F be the Kähler form of M and put $K = R^M - R^{N(\lambda)}$ on M , where R^M and $R^{N(\lambda)}$ are the curvature tensors of M and $N(\lambda)$, respectively. Then K is a Kähler curvature-like tensor field on M . For $p = 1, \dots, m$, let $k_{2p}(M)$ be the curvature invariant defined for the tensor field K by Weyl in [9] (see also (7.6) in [4]). For $p = 1, \dots, m$, let γ_p be the p th Chern form of M . Put $\Gamma_p(M) = (\gamma_p \wedge F^{m-p})[M]$ for $1 \leq p \leq m$ and $\Gamma_0(M) = F^m[M]$. Then we have (see (3.26) and (3.27) in [5] and also Lemma 4.1 in [6]).

LEMMA 1 (A. GRAY [5]). For $1 \leq p \leq m$, we have

$$(1.1) \quad k_{2p}(M) = \sum_{q=0}^p \frac{(2\pi)^p}{(m-p)!} \binom{m-q+1}{p-q} \left(-\frac{\lambda}{\pi}\right)^{p-q} \Gamma_q(M).$$

Conversely we will get

LEMMA 2. For $1 \leq p \leq m$, we have

$$(1.2) \quad \Gamma_p(M) = \sum_{q=1}^p \frac{(m-q)!}{(2\pi)^q} \binom{m-q+1}{p-q} \left(\frac{\lambda}{\pi}\right)^{p-q} k_{2q}(M) + \binom{m+1}{p} \left(\frac{\lambda}{\pi}\right)^p F^m[M].$$

Next we state Shifrin's kinematic formulas (see (3.5) and (4.4) in [8]).

THEOREM 1 (T. SHIFRIN [8]). Let M^m and N^l be Kähler submanifolds of complex dimensions m and l , respectively, in a Fubini-Study space $N(\lambda)$. Assume

Received by the editors February 20, 1985 and, in revised form, June 10, 1985.

1980 *Mathematics Subject Classification.* Primary 53C65; Secondary 53C40, 53C55.

Key words and phrases. Kinematic formulas, Chern forms, Weyl's curvature invariants.

¹ Research partially supported by the Grant in Aid for Scientific Research (No. 59540045).

that $k = m + l - n \geq 1$. Then, for $1 \leq h \leq k$,

(1.3)

$$\int_{U(n+1) \rightarrow g} \Gamma_h(M^m \cap gN^l) dg \\ = \text{Vol.}(U(n+1)) \sum_{0 \leq i, 0 \leq j}^{i+j \leq n} (-1)^h \binom{n+h-i-j}{n} \left(\frac{\lambda}{\pi}\right)^{n+h-i-j} \Gamma_i(M^m) \Gamma_j(N^l)$$

whenever both sides make sense. Let $G(l, n)$ be the space of projective l -planes in $CP^n (= N(\lambda))$. Then, for $0 \leq h \leq k$,

$$(1.4) \quad \int_{G(l, n) \ni L} \Gamma_h(M^m \cap L) dL \\ = \text{Vol.}(G(l, n)) \sum_{j=0}^h (-1)^{h-j} \binom{n+h-l-j-1}{h-j} \left(\frac{\lambda}{\pi}\right)^{n+h-l-j} \Gamma_j(M^m),$$

whenever both sides makes sense.

The normalized Kähler form $\omega = \lambda F/\pi$ is used in [8] so that the coefficients in Shifrin's original kinematic formulas are slightly different from those in the above formulas. Using Lemmas 1 and 2, we obtain from Theorem 1 the following kinematic formulas for Weyl's curvature invariants k_{2p} .

THEOREM 2. Let M^m and N^l be Kähler submanifolds with compact closure in a Fubini-Study space $N(\lambda)$ of constant holomorphic curvature 4λ ($\lambda > 0$). Assume that $k = m + l - n \geq 1$. Then, for $1 \leq p \leq k$,

$$(1.5) \quad \int_{U(n+1)} k_{2p}(M^m \cap gN^l) dg = \text{Vol.}(U(n+1)) \sum_{0 \leq a, b}^{a+b \leq p} F_{pab} k_{2a}(M^m) k_{2b}(N^l),$$

$$(1.6) \quad \int_{G(l, n)} k_{2p}(M^m \cap L) dL = \text{Vol.}(G(l, n)) \left(\frac{\lambda}{\pi}\right)^{n-l} \frac{(m-p)!}{(k-p)!} k_{2p}(M^m),$$

where

$$(1.7) \quad \begin{cases} f_{pab} = (-1)^p \left(\frac{\lambda}{\pi}\right)^n \frac{(2\sqrt{\lambda})^{2(p-a-b)}}{(p-a-b)!} \frac{(m-a)!(l-b)!}{(k-p)!} \frac{(n+k-a-b+2)!}{(n+k-p+2)!}, \\ F_{pab} = f_{pab} \quad \text{for } a \geq 1 \text{ and } b \geq 1, \\ F_{p0b} = f_{p0b}/m! \quad \text{for } p \geq b \geq 1, \\ F_{pa0} = f_{pa0}/l! \quad \text{for } p \geq a \geq 1, \quad F_{p00} = f_{p00}/m!l!. \end{cases}$$

Weyl's curvature invariants have relations with other invariants. In fact, let $\mu_{2p}(K)$ be the Chern's curvature invariants for the curvature-like tensor K (see [2]). Put $(2p-1)!! = (2p-1) \cdot (2p-3) \cdots 3 \cdot 1$. Then it holds that

$$\mu_{2p}(K) = \left(\binom{2m}{2p} (2p-1)!! \right)^{-1} k_{2p}(M).$$

Also, $k_{2p}(M)$ are closely related to mean curvatures of M in $N(\lambda)$ (see, for example, [1]).

2. Proof of Lemma 2. We may put $k_0(M) = (1/m!)\Gamma_0(M)$. Let, for $0 \leq p \leq m$ and $0 \leq q \leq m$,

$$a_p = \frac{(m-p)!}{(2\pi)^p} \left(-\frac{\pi}{\lambda}\right)^p k_{2p}(M), \quad b_q = \left(-\frac{\pi}{\lambda}\right)^q \Gamma_q(M).$$

Then, from (1.1) we have, for $0 \leq p \leq m$, $a_p = \sum_{q=0}^p \binom{m+1-q}{p-q} b_q$. Now, we use formula (7a) (on p. 53 in [7]), and get

$$b_p = \sum_{q=0}^p (-1)^{p+q} \binom{m+1-q}{p-q} a_q.$$

This implies (1.2).

3. Proof of Theorem 2. First we will show (1.5). If we write (1.3) as

$$\int \Gamma_h(M^m \cap gN^l) dg = \sum_{\substack{i+j \leq h \\ 0 \leq i, 0 \leq j}} D_{hij} \Gamma_i(M^m) \Gamma_j(N^l),$$

by means of (2.1) and (2.2), we have, for $1 \leq p \leq k$,

$$\int k_{2p}(M^m \cap gN^l) dg = \sum_{h=0}^p \sum_{\substack{i+j \leq h \\ 0 \leq i, 0 \leq j}} \sum_{a=0}^i \sum_{b=0}^j A_{ph} D_{hij} B_{ia} B_{jb} k_{2a}(M^m) k_{2b}(N^l).$$

Thus we obtain

$$(3.1) \quad \text{Vol.}(U(n+1)) F_{pab} = \sum_{h=a+b}^p \sum_{\substack{i+j \leq h \\ a \leq i, b \leq j}} A_{ph} D_{hij} B_{ia} B_{jb}.$$

Put

$$(3.2) \quad f_{pab} = (-1)^p \left(\frac{\lambda}{\pi}\right)^n (2\lambda)^{p-a-b} \frac{(m-a)!(l-b)!}{(k-p)!} I_p,$$

(3.3)

$$I_p = \sum_{h=a+b}^p \sum_{\substack{i+j \leq j \\ a \leq i, b \leq j}} \binom{k-h+1}{p-h} \binom{n+h-i-j}{n} \binom{m-a+1}{i-a} \binom{l-b+1}{j-b}.$$

Using (1.1)–(1.3), from (3.1) we get, for $1 \leq a$, $1 \leq b$, $a+b \leq p$,

$$(3.4) \quad \begin{aligned} F_{pab} &= f_{pab}, & F_{p0b} &= f_{j0b}/m!, \\ F_{pa0} &= f_{pa0}/l!, & F_{p00} &= f_{p00}/m!l!. \end{aligned}$$

To calculate I_p , we write them as

$$(3.5) \quad I_p = \sum_{h'=0}^{p'} \sum_{t=0}^{h'} \binom{k'-h+1}{p'-h} \binom{n+h-t}{n} \binom{m'+l'+2}{t},$$

where $m' = m - a$, $l' = l - b$, $k' = k - a - b$ and $p' = p - a - b$. We use the identity

$$\sum_{i+j=t} \binom{m'+1}{i} \binom{l'+1}{j} = \binom{m'+l'+2}{t}$$

to get (3.5). We know the identities (see, for example, [7])

$$(3.6) \quad \sum_{h=t}^{p'} \binom{k' - h + 1}{p' - h} \binom{n + h - t}{n} = \binom{n + k' - t + 2}{p' - t},$$

$$(3.7) \quad \sum_{t=0}^{p'} \binom{n + k' + 2}{t} \binom{n + k' + 2 - t}{p' - t} = 2^{p'} \binom{n + k' + 2}{p'}.$$

Hence, from (3.5), it follows that

$$I_p = 2^{p'} \binom{n + k' + 2}{p'} = 2^{p-a-b} \binom{n + k - a - b + 2}{p - a - b}.$$

Combining this with (3.2) and (3.4), we get (1.7).

Next, we prove (1.6). We may put, for $1 \leq p \leq k$,

$$\begin{aligned} \int \Gamma_p(M^m \cap L) dL &= \sum_{q=0}^p E_{pq} \Gamma_q(M^m), \\ \int k_{2p}(M^m \cap L) dL &= \sum_{q=1}^p G_{pq} k_{2q}(M^m). \end{aligned}$$

Using (2.1) and (2.2), we get

$$(3.8) \quad G_{pq} = \sum_{h=q}^p \sum_{i=q}^h A_{ph} E_{hi} B_{iq}.$$

Assume that $1 \leq q < p$. Then, using (1.1), (1.2) and (1.4), we get

$$(3.9) \quad G_{pq} = \text{Vol.}(G(l, n)) (2\lambda)^{p-q} \left(\frac{\lambda}{\pi}\right)^{n-l} \frac{(m-q)!}{(k-p)!} H_{pq},$$

$$(3.10) \quad H_{pq} = \sum_{i'=0}^{p'} \sum_{h'=i'}^{p'} (-1)^{p'-i'} \binom{k' - h' + 1}{p' - h'} \binom{n - l + h' - i' - 1}{h' - i'} \binom{m' + 1}{i'},$$

where $i' = i - q$, $p' = p - q$, $h' = h - q$ and $m' = m - q$. Similar to (3.6), we have

$$\sum_{h'=i'}^{p'} \binom{k' - h' + 1}{p' - h'} \binom{n - l + h' - i' - 1}{h' - i'} = \binom{m' - i' + 1}{m' - p' + 1}.$$

Hence, from (3.10), it follows that

$$H_{pq} = (-1)^{p'} \sum_{i'=0}^{p'} (-1)^{i'} \binom{m' + 1 - i'}{m' + 1 - p'} \binom{m' + 1}{i'} = 0.$$

This gives $G_{pq} = 0$ for $1 \leq q < p$. Similarly we get $G_{p0} = 0$. Thus we obtain $G_{pq} = 0$ for $0 \leq q < p$. On the other hand, we have

$$G_{pp} = A_{pp} E_{pp} B_{pp} = \text{Vol.}(G(l, n)) \left(\frac{\lambda}{\pi}\right)^{n-l} \frac{(m-p)!}{(k-p)!}.$$

Thus we proved (1.6).

ACKNOWLEDGMENT. The proof of Lemma 2 in §2 is due to the referee. I would like to thank him for valuable suggestions.

REFERENCES

1. B. Y. Chen, *On the total curvature of immersed manifolds. I*, Amer. J. Math. **93** (1971), 148–162.
2. S. S. Chern, *On the kinematic formula in integral geometry*, J. Math. Mech. **16** (1966), 101–118.
3. A. Gray, *Chern numbers and curvature*, Amer. J. Math. **100** (1978), 463–476.
4. ———, *Comparison theorems for the volumes of tubes as generalizations of the Weyl tube formula*, Topology **21** (1982), 201–228.
5. ———, *Volumes of tubes about Kähler submanifolds expressed in terms of Chern classes*, J. Math. Soc. Japan **36** (1984), 23–35.
6. ———, *A product formula for the volumes of tubes about Kähler submanifolds of complex euclidean space*, Riv. Mat. Univ. Parma (4) **10** (1984), 109–121.
7. J. Riordan, *Combinatorial identities*, Wiley, New York, 1968.
8. T. Shifrin, *The kinematic formula in complex integral geometry*, Trans. Amer. Math. Soc. **264** (1981), 255–293.
9. H. Weyl, *On the volume of tubes*, Amer. J. Math. **61** (1939), 461–472.

DEPARTMENT OF MATHEMATICS, TOKUSHIMA UNIVERSITY, JOSANJIMA TOKUSHIMA
770, JAPAN