ON THE DIOPHANTINE EQUATION $x^{2n} - \mathcal{D}y^2 = 1$

CAO ZHENFU

ABSTRACT. In this paper, it has been proved that if $n > 2$ and Pell's equation $u^2 - \mathcal{D}v^2 = -1$ has integer solution, then the equation $x^{2n} - \mathcal{D}y^2 = 1$ has only solution in positive integers $x = 3$, $y = 22$ (when $n = 5$, $\mathcal{D} = 122$). That is proved by studying the equations $x^p + 1 = 2y^2$ and $x^p - 1 = 2y^2$ (p is an odd prime). In addition, some applications of the above result are given.

1. Introduction. For the Diophantine equation

$$(1) \quad x^{2n} - \mathcal{D}y^2 = 1$$

$\mathcal{D} > 0$ and is not a perfect square, and $n > 1$ is an integer number. When $n = 2$, its solvability was discussed by Ljunggren, Cohn, Ko Chao and Sun Chi [1], and the author [2], etc. In this paper, we discuss the solvability of equation (1) in the case $n > 2$.

THEOREM 1. *If $n > 2$ and Pell's equation $u^2 - \mathcal{D}v^2 = -1$ has integer solution, then the Diophantine equation (1) has only solution in positive integers $x = 3$, $y = 22$ (when $n = 5$, $\mathcal{D} = 122$).*

Tartakowski [3] has proved that if $n > 2$ and the equation $u^2 - \mathcal{D}v^2 = -1$ has solution, then the equation $x^{2n} - \mathcal{D}y^{2n} = 1$ has no solution in positive integers. Clearly, the result of Tartakowski is included in Theorem 1.

THEOREM 2. *If $\eta = U_0 + V_0\sqrt{\mathcal{D}}$ is the fundamental solution of Pell's equation $U^2 - \mathcal{D}V^2 = 1$, then positive integer solutions of equation (1) do not satisfy*

$$x^n + y\sqrt{\mathcal{D}} = \eta^{4m}, \quad n > 2, m > 0.$$

The proof of the above result is obtained by studying equations

$$(2) \quad x^p + 1 = 2y^2, \quad p \text{ odd prime} > 3,$$

and

$$(3) \quad x^p - 1 = 2y^2, \quad p \text{ odd prime} > 3,$$

with Ko Chao's elementary method (see Ko Chao [4]). The method is helpful to the research of Erdős' conjecture on combinatorial

Received by the editors October 24, 1984 and, in revised form, May 29, 1985.
 1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 11D15; Secondary 11B10.

2. Two lemmas on equations (2) and (3).

LEMMA 1. *If equation (2) has positive integer solution, then $2p|y$ except $x = y = 1$.*

PROOF. Suppose (x, y) is any positive integer solution of (2). Then

$$(x+1)((x^p+1)/(x+1)) = 2y^2.$$

Since $(x+1, (x^p+1)/(x+1)) = 1$ or p , we have

$$(4) \quad x+1 = 2y_1^2, \quad (x^p+1)/(x+1) = y_2^2, \quad y = y_1y_2,$$

or

$$(5) \quad x+1 = 2py_1^2, \quad (x^p+1)/(x+1) = py_2^2, \quad y = py_1y_2.$$

By the result of Ljunggren [5], $(x^p+1)/(x+1) = y_2^2$, therefore $x = 1$. Thus (4) gives $x = y = 1$.

For (5), clearly $p|y$. We will prove $2|y$ with Ko Chao's elementary method.

If $2 \nmid y$, from (2), we have $x \equiv 1 \pmod{8}$. Put

$$\mathcal{A}(t) = (x^t+1)/(x+1), \quad t \geq 1 \text{ and } 2 \nmid t,$$

and so $\mathcal{A}(t) \equiv 1 \pmod{8}$. Let $1 < l < p$ be a positive odd integer. Then there exist an integer r , odd, $0 < r < l$, and $2k$ such that $p = 2kl + r$ or $p = 2kl - r$.

If $p = 2kl + r$, then

$$(6) \quad \mathcal{A}(p) = \frac{((x+1)\mathcal{A}(l)-1)^{2k}x^r+1}{x+1} \equiv \frac{x^r+1}{x+1} \equiv \mathcal{A}(r) \pmod{\mathcal{A}(l)},$$

since $x^l = (x+1)\mathcal{A}(l) - 1$. Now $(\mathcal{A}(p), \mathcal{A}(l)) = \mathcal{A}((p, l)) = \mathcal{A}(1) = 1$. Thus, (6) gives

$$(\mathcal{A}(p)/\mathcal{A}(l)) = (\mathcal{A}(r)/\mathcal{A}(l)).$$

If $p = 2kl - r$, then $l - r$ is even. Thus

$$(\mathcal{A}(p)/\mathcal{A}(l)) = (-x^{l-r}\mathcal{A}(r)/\mathcal{A}(l)) = (\mathcal{A}(r)/\mathcal{A}(l)),$$

since $\mathcal{A}(l) \equiv 1 \pmod{8}$ and

$$\mathcal{A}(p) = x^{l-r}\mathcal{A}(l(2k-1)) + \mathcal{A}(l) - x^{l-r}\mathcal{A}(r).$$

For l, r , we have

$$l = 2k_1r + \varepsilon_1r_1, \quad 0 < r_1 < r,$$

$$r = 2k_2r_1 + \varepsilon_2r_2, \quad 0 < r_2 < r_1,$$

...

$$r_{s-1} = 2k_{s+1}r_s + \varepsilon_{s+1}r_{s+1}, \quad 0 < r_{s+1} < r_s,$$

$$r_s = k_{s+2}r_{s+1},$$

where $\varepsilon_i = \pm 1$ ($i = 1, \dots, s+1$) and r_i ($i = 1, \dots, s+1$) are odd integers. Since $(l, p) = 1$, we have $r_{s+1} = 1$. Hence

$$\begin{aligned} \left(\frac{\mathcal{A}(p)}{\mathcal{A}(l)} \right) &= \left(\frac{\mathcal{A}(r)}{\mathcal{A}(l)} \right) = \left(\frac{\mathcal{A}(l)}{\mathcal{A}(r)} \right) = \left(\frac{\mathcal{A}(r_1)}{\mathcal{A}(r)} \right) = \left(\frac{\mathcal{A}(r)}{\mathcal{A}(r_1)} \right) = \left(\frac{\mathcal{A}(r_2)}{\mathcal{A}(r_1)} \right) \\ &= \dots = \left(\frac{\mathcal{A}(r_{s+1})}{\mathcal{A}(r_s)} \right) = \left(\frac{\mathcal{A}(1)}{\mathcal{A}(r_s)} \right) = \left(\frac{1}{\mathcal{A}(r_s)} \right) = 1. \end{aligned}$$

Now, from $(x^p + 1)/(x + 1) = py_2^2$, we have

$$(py_2)^2 \equiv p\mathcal{A}(p) \pmod{\mathcal{A}(l)}.$$

Thus

$$\left(\frac{p\mathcal{A}(p)}{\mathcal{A}(l)}\right) = \left(\frac{p}{\mathcal{A}(l)}\right) = \left(\frac{\mathcal{A}(l)}{p}\right) = \left(\frac{l}{p}\right) = 1,$$

since $x \equiv -1 \pmod{p}$ and so $\mathcal{A}(l) \equiv l \pmod{p}$. We have a contradiction if l is taken as an odd quadratic nonresidue of p . This proves the result. $\square$

REMARK 1. Using the above method, we can prove that the equations

$$x^p \pm y^p = 2z^2, \quad (x, y) = 1, p \text{ prime} > 3,$$

have no positive integer solution when $2 \nmid z, p \nmid z$.

LEMMA 2. Equation (3) has only positive integer solution $x = 3, y = 11$ (when $p = 5$).

PROOF. From (3), we have

$$(7) \quad (x^p - 1)/(x - 1) = a^2,$$

if $p \nmid y$. By the result of Ljunggren [5], the solution of (7) is $x = 3$ (when $p = 5$). Thus (3) has positive integer solution $x = 3, y = 11$ (when $p = 5$).

If $p \mid y$, then $2 \mid y$ by Remark 1. From (3), $(1 + \sqrt{-2}y)(1 - \sqrt{-2}y) = x^p$. With $(1 + \sqrt{-2}y, 1 - \sqrt{-2}y) = 1$, we have

$$(8) \quad 1 + \sqrt{-2}y = (a + b\sqrt{-2})^p, \quad x = a^2 + 2b^2,$$

where a, b are rational integers. Since $2 \mid y$, from (3), it follows that

$$(9) \quad x \equiv 1 \pmod{8}.$$

From (8) and (9), we have $2 \mid b$ and $b \neq 0$. Now, (8) gives

$$(10) \quad 1 = a^p + \binom{p}{2} a^{p-2} (b\sqrt{-2})^2 + \cdots + \binom{p}{p-1} a (b\sqrt{-2})^{p-1}.$$

Thus $a \mid 1$ and so $a = \pm 1$.

If $a = -1$, then (10) gives

$$-2 = \binom{p}{2} (b\sqrt{-2})^2 + \cdots + \binom{p}{p-1} (b\sqrt{-2})^{p-1},$$

and so $p \mid 2$ which is impossible.

If $a = 1$, then we have

$$(11) \quad 0 = \binom{p}{2} (b\sqrt{-2})^2 + \cdots + \binom{p}{p-1} (b\sqrt{-2})^{p-1}.$$

Since $2 \mid b$ and $b \neq 0$, let $2^{s_k} \parallel \binom{p}{2k} (b\sqrt{-2})^{2k}$ ($1 \leq k \leq (p-1)/2$), clearly $s_k > s_j$ ($k > j$). Thus (11) is impossible.

REMARK 2. Nagell [8] also proved Lemma 2, but the above proof is a new method.

3. Proof of the theorems.

PROOF OF THEOREM 1. Let $\Omega = u_0 + v_0\sqrt{D}$ be the fundamental solution of the equation $u^2 - Dv^2 = -1$, $\bar{\Omega} = u_0 - v_0\sqrt{D}$, $\Omega\bar{\Omega} = -1$, and let $\eta = U_0 + V_0\sqrt{D}$ be the fundamental solution of the equation $U^2 - DV^2 = 1$, $\eta = U_0 - V_0\sqrt{D}$, $\eta\bar{\eta} = 1$. Then, we have $\eta = \Omega^2$.

Suppose (x, y) is any positive integer solution of (1). Then

$$(12) \quad x^n = \frac{\eta^m + \bar{\eta}^m}{2} = \frac{\Omega^{2m} + \bar{\Omega}^{2m}}{2}, \quad m > 0.$$

Clearly, without loss of generality, we may assume that $n = 4$ or $n = p$ (p is odd prime).

(a) If $n = 4$, then (12) gives

$$x^4 = 2((\Omega^m + \bar{\Omega}^m)/2)^2 - (-1)^m,$$

and so $x = 1$, $m = 0$ which is impossible since $m > 0$.

(b) If $n = p$ (p is odd prime), then (12) gives

$$(13) \quad x^p = 2((\Omega^m + \bar{\Omega}^m)/2)^2 - (-1)^m,$$

(b.1) When $2|m$, let $m = 2s$, $s > 0$; then (13) gives

$$(14) \quad x^p + 1 = 2((\Omega^{2s} + \bar{\Omega}^{2s})/2)^2.$$

Suppose $p = 3$. Then by (14), we have (see [6])

$$(15) \quad x = (\Omega^{2s} + \bar{\Omega}^{2s})/2 = 1,$$

and

$$(16) \quad x = 23, \quad (\Omega^{2s} + \bar{\Omega}^{2s})/2 = 78.$$

Clearly (15) is impossible since $s > 0$, and (16) is also impossible since

$$\frac{\Omega^{2s} + \bar{\Omega}^{2s}}{2} = 2 \left(\frac{\Omega^s + \bar{\Omega}^s}{2} \right)^2 - (-1)^s$$

is odd.

Thus $p > 3$. For (14), we have $2p|(\Omega^{2s} + \bar{\Omega}^{2s})/2$ by Lemma 1. However, $2|(\Omega^{2s} + \bar{\Omega}^{2s})/2$ is impossible.

(b.2) When $2 \nmid m$, we have

$$(17) \quad x^p - 1 = 2((\Omega^m + \bar{\Omega}^m)/2)^2,$$

and so $x = 1$, $(\Omega^m + \bar{\Omega}^m)/2 = 0$ when $p = 3$ (see [6]). If $p > 3$, then (17) gives $x = 3$, $(\Omega^m + \bar{\Omega}^m)/2 = 11$ (when $p = 5$) by Lemma 2. Thus (1) has only positive integer solution $x = 3$, $y = 22$ (when $n = 5$, $D = 122$).

This completes the proof of Theorem 1. $\square$

PROOF OF THEOREM 2. If

$$x^n + y\sqrt{D} = \eta^{4m}, \quad n > 2, m > 0,$$

then we have

$$(18) \quad x^n = \frac{\eta^{4m} + \bar{\eta}^{4m}}{2} = 2 \left(\frac{\eta^{2m} + \bar{\eta}^{2m}}{2} \right)^2 - 1.$$

By Lemma 1, (18) is impossible since $2 \nmid (\eta^{2m} + \bar{\eta}^{2m})/2$ and $m > 0$. $\square$

4. Applications. As applications of the result, we will discuss other problems in number theory in the following paragraphs.

4.1. In 1939, Erdős [7] conjectured that the equation

$$(19) \quad \binom{n}{m} = y^k, \quad n > m > 1, k \geq 3,$$

has no integer solution. In 1951, Erdős himself proved that the conjecture is right when $m > 3$, leaving the cases $m = 2$ and $m = 3$ unsolved. Now, we can deduce the following corollary from Theorem 1 and Lemma 1.

COROLLARY 1. *The equation $\binom{n}{2} = y^{2k}$ has no positive integer solution n, y with $n > 2$ and $k > 1$.*

PROOF. From $\binom{n}{2} = n(n-1)/2 = y^{2k}$, we have

$$n-1 = 2y_1^{2k}, \quad n = y_2^{2k}, \quad y = y_1 y_2,$$

or

$$n-1 = y_2^{2k}, \quad n = y_1^{2k}, \quad y = y_1 y_2.$$

Hence

$$(20) \quad y_2^{2k} \mp 1 = 2y_1^{2k}.$$

If $2|k$, then (20) clearly gives $|y_1 y_2| \leq 1$; on the other hand, $n > 2$ and $\binom{n}{2} = y^{2k}$ imply $|y| = |y_1 y_2| > 1$. here we have a contradiction. If $2 \nmid k$, $k > 1$, we may conclude from Theorem 1 and Lemma 1 that (20) is impossible. $\square$

4.2. For the Pell sequence

$$(21) \quad x_0 = 1, \quad x_1 = a, \quad x_{n+2} = 2ax_{n+1} - x_n,$$

where a is an integer > 1 , we have

COROLLARY 2. *The equation $x_{4n} = y^m$ has no positive integer solution n, y when $m > 2$.*

PROOF. From (21), we have $x_n = (\alpha^n + \bar{\alpha}^n)/2$, $n \geq 0$, where $\alpha = a + \sqrt{a^2 - 1}$ and $\bar{\alpha} = a - \sqrt{a^2 - 1}$ are roots of the trinomial $z^2 - 2az + 1$. Let $a^2 - 1 = \mathcal{D}b^2$, where $\mathcal{D} > 0$ is squarefree and b is positive integer. Then

$$\alpha = a + b\sqrt{\mathcal{D}}, \quad \bar{\alpha} = a - b\sqrt{\mathcal{D}} \quad \text{and} \quad \alpha\bar{\alpha} = 1.$$

Thus $y_n = (\alpha^n - \bar{\alpha}^n)/2\sqrt{\mathcal{D}}$ satisfies

$$(22) \quad x_n^2 - \mathcal{D}y_n^2 = 1.$$

By Theorem 2, (22) is impossible when $4|n$, $x_n = y^m$ and $m > 2$. $\square$

Clearly, if $a = 2u^2 + 1$ ($u > 0$), then $\mathcal{D}b^2 = a^2 - 1 = 4u^2(u^2 + 1)$. Thus $2u|b$. let $b = 2uv$, we have $u^2 + 1 = \mathcal{D}v^2$. Hence, using Theorem 1, we have

COROLLARY 3. *For the Pell sequence*

$$x_0 = 1, \quad x_1 = 2u^2 + 1, \quad x_{n+2} = 2(2u^2 + 1)x_{n+1} - x_n,$$

where u is positive integer, x_n is never an m th power if $m > 2$ and except $x_1 = 2 \cdot 11^2 + 1 = 3^5$. $\square$

ACKNOWLEDGMENT. The author would like to thank Professor Sun Chi for his helpful suggestions.

REFERENCES

1. Ko Chao and Sun Chi, *On the diophantine equation $x^4 - Dy^2 = 1$* . II, Chinese Ann. Math. Ser. A **1** (1980), 83–88.
2. Cao Zhenfu, *On the diophantine equations $x^2 + 1 = 2y^2$ and $x^2 - 1 = 2Dz^2$* , J. Math. (Wuhan) **3** (1983), 227–235.
3. L. J. Mordell, *Diophantine equations*, Academic Press, London and New York, 1969.
4. Ko Chao, *On the diophantine equation $x^2 = y^n + 1, xy \neq 0$* , Sichun Daxue Xuebao **1** (1962), 1–6.
5. W. Ljunggren, *Some theorems on indeterminate equations of the form $(x^n - 1)/(x - 1) = y^q$* , Norsk Mat. Tidsskr. **25** (1943), 17–20.
6. R. W. van der Waall, *On the diophantine equations $x^2 + x + 1 = 3v^2, x^3 - 1 = 2y^2, x^3 + 1 = 2y^2$* , Simon Stevin **46** (1972/73), 39–51.
7. P. Erdős, *On a diophantine equation*, J. London Math. Soc. **26** (1951), 176–178.
8. W. Ljunggren, *On the Diophantine equation $Cx^2 + D = 2y^n$* , Math. Scand. **18** (1966), 69–86.

DEPARTMENT OF MATHEMATICS, HARBIN INSTITUTE OF TECHNOLOGY, HARBIN,
CHINA