A LOCAL STRONG UV^∞ -PROPERTY OF THE HOMEOMORPHISM GROUPS OF R^∞ - (Q^∞ -) MANIFOLDS

VO THANH LIEM

ABSTRACT. We will show in this note that if M is an R^∞ - (or Q^∞ -) manifold having the homotopy type of a finite complex, then the homeomorphism group of M , endowed with the compact-open topology, has the local strong UV^∞ -property with respect to the classes of pseudo CW complexes $\mathcal{CW}(\mathcal{C})$ and $\mathcal{CW}(\mathcal{M})$.

Let $E^\infty = \text{dirlim } E^n$ where E^n is either the n -Euclidean space or the n -fold product of the Hilbert cube Q . By an E^∞ -manifold, we mean a paracompact space which is locally homeomorphic to E^∞ . Given a space X , let $\text{Homeo}(X)$ denote the group of homeomorphisms of X endowed with the compact-open topology. Let A be a subset of $S \times X$ (or $X \times S$), a map $f: A \rightarrow S \times Y$ (or $f: A \rightarrow Y \times S$) is said to be S -fiber preserving (S -f.p.) if $\pi_S f = \pi_S$ where π_S denotes the projection. Given an S -f.p. map $f: S \times X \rightarrow S \times Y$, for each $s \in S$, let $f_s: X \rightarrow Y$ denote the map defined by $f(x, s) = (s, f_s(x))$ for each $x \in X$. By an S -f.p. isotopy (an S -f.p. invertible isotopy, resp.) from $S \times X$ onto $S \times Y$, we mean an $(I \times S)$ -f.p. map $f: I \times S \times X \rightarrow I \times S \times Y$ such that $f_{(t,s)}$ is a homeomorphism for each $(t, s) \in I \times S$ (f is an $(I \times S)$ -f.p. homeomorphism, resp.). Given an open cover $\mathcal{U}$ of Y , a homotopy $H: I \times X \rightarrow Y$ is said to be an $\mathcal{U}$ -homotopy if for each $x \in X$ there is an $U \in \mathcal{U}$ such that $H(I \times x) \subset U$. Two maps $f, g: X \rightarrow Y$ are said to be $\mathcal{U}$ -close if for each $x \in X$, there is an $U \in \mathcal{U}$ such that $\{f(x), g(x)\} \subset U$. Following [L], let $\mathcal{CW}(\mathcal{C})$ and $\mathcal{CW}(\mathcal{M})$ denote the classes of pseudo CW complexes generated by the class $\mathcal{C}$ of Hausdorff compact spaces and the class $\mathcal{M}$ of metric spaces, respectively. If A is a subset of X , let i_A denote the inclusion $A \hookrightarrow X$, and id_X the identity of X .

The homeomorphism group $\text{Homeo}(X)$ of a space X is said to have the *local strong UV^∞ -property with respect to a class $\mathcal{X}$* provided that given an open neighborhood U of an $h \in \text{Homeo}(X)$, there is an open neighborhood V of h in U such that if $\Lambda \in \mathcal{X}$, Γ is a closed G_δ -set in Λ , and $g: \Lambda \rightarrow V$ is a continuous map with $g(\Gamma) = \{h\}$, then g is homotopic rel Γ in V to the constant map $H: V \rightarrow \{h\}$.

If M is an R^∞ -manifold, Hale provided in [H] that $\text{Homeo}(M)$ is Lindelöf, homogeneous, and paracompact, but not compactly generated. In this note, we

Received by the editors April 12, 1985 and, in revised form, September 16, 1985.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 57S05.

Key words and phrases. k -space, invertible isotopy.

©1986 American Mathematical Society
 0002-9939/86 \$1.00 + \$.25 per page

intend to study the local contractibility of the groups of homeomorphisms of E^∞ -manifolds.

THEOREM 1. *If M is an E^∞ -manifold having the homotopy type of a finite complex, then $\text{Homeo}(M)$ has the local strong UV^∞ -property with respect to $\mathcal{CW}(\mathcal{C}) \cup \mathcal{CW}(\mathcal{M})$.*

Note that if $M = \tilde{D} \times E^\infty$ where $\tilde{D}$ is the open disk with infinitely many handles attached (refer to [R, p. 275]), then $\text{Homeo}(M)$ is not locally path-connected.

Let us introduce some notations that will be used in the following proofs. All spaces are Hausdorff. M is a given E^∞ -manifold having the homotopy type of a compact subpolyhedron A of an R^m . Let N be a regular neighborhood of A in R^m [Hd] and identify M with $N \times E^\infty$ [He, Theorem C] and N with $N \times 0$ in $N \times E^\infty$. Referring to [H-T], write $M = \text{dirlim } M_j$ where $M_j = N \times B_{j+1}$ and B_s denotes the s -fold product of $[-s, s]$ (Q , resp.) if $E = R$ (if $E = Q = \prod_1^\infty [0, 1]$, resp.). Given a positive integer s and a sequence $\{\varepsilon_s, \varepsilon_{s+1}, \dots\}$ of positive numbers, we write $E_{\{\varepsilon_s, \varepsilon_{s+1}, \dots\}}^\infty = \{x \in E^\infty \mid x_1 = \dots = x_{s-1} = 0 \text{ and } d(x_j, 0) < \varepsilon_j \text{ for } j = s, s+1, \dots\}$.

Given a compact subset K of M and an open set V of M , following [D] let $[K, V]$ denote the set $\{f \in \text{Homeo}(M) \mid f(K) \subset V\}$; this is a member of a subbasis for the compact-open topology on $\text{Homeo}(M)$. We will work with the category $\mathcal{CG}$ of compactly generated spaces [G]. For $X, Y \in \mathcal{CG}$, by $X \times Y$ we mean the $\mathcal{CG}$ -product of X and Y . Recall that $\mathcal{CW}(\mathcal{C}) \cup \mathcal{CW}(\mathcal{M})$ is a subclass of $\mathcal{CG}$ [L]; moreover, that if $\Lambda \in \mathcal{CW}(\mathcal{C}) \cup \mathcal{CW}(\mathcal{M})$, then so does $I \times \Lambda$.

LEMMA 2. *Let Γ be a closed subset of $\Lambda \in \mathcal{CW}(\mathcal{C}) \cup \mathcal{CW}(\mathcal{M})$ and q a nonnegative integer. Let $f, g: \Lambda \times M \rightarrow \Lambda \times M$ be Λ -f.p. maps such that $f|_{\Gamma \times M} = g|_{\Gamma \times M}$ and such that there is a Λ -f.p. homotopy ϕ from $f|_{\Lambda \times M_q}$ to $g|_{\Lambda \times M_q}$ ($\text{rel } \Gamma \times M_q$). Then, ϕ can be extended to a Λ -f.p. homotopy ψ from f to g ($\text{rel } \Gamma \times M$). Moreover, if g is a Λ -f.p. homotopy equivalence, then $\hat{\psi}: I \times \Lambda \times M \rightarrow I \times \Lambda \times M$, defined by $\hat{\psi}(t, \lambda, z) = (t, \psi(t, \lambda, z))$, is an $(I \times \Lambda)$ -f.p. homotopy equivalence.*

PROOF. Let $A = (I \times \Lambda \times M_q) \cup (I \times \Gamma \times M) \cup (\{0, 1\} \times \Lambda \times M)$ and $\bar{\phi}: A \rightarrow \Lambda \times M$ be the extension of ϕ defined by

$$\bar{\phi}(t, \lambda, x) = \begin{cases} \phi(t, \lambda, x) & \text{if } x \in M_q, \\ f(\lambda, x) (= g(\lambda, x)) & \text{if } \lambda \in \Gamma, \\ f(\lambda, x) & \text{if } t = 0, \\ g(\lambda, x) & \text{if } t = 1. \end{cases}$$

Since M_q is the strong deformation retract of M , we have a homotopy $\theta: M \times I \rightarrow M$ such that $\theta_0 = \text{id}$, $\theta_1(M) = M_q$ and $\theta_t|_{M_q} = \text{id}$ for each $t \in I$. Then, $\bar{\theta} = \text{id} \times \theta: I \times \Lambda \times M \times I \rightarrow I \times \Lambda \times M$ is an $(I \times \Lambda)$ -f.p. homotopy such that $\bar{\theta}_0 = \text{id}$, $\bar{\theta}(A \times I) \subset A$, and $\bar{\theta}_1: I \times \Lambda \times M \rightarrow I \times \Lambda \times M_q$ is a $(I \times \Lambda)$ -f.p. retraction. Observe that $\bar{\phi} = \bar{\phi}\bar{\theta}_0|_A$ is Λ -f.p. homotopic to $\bar{\phi}\bar{\theta}_1|_A = \phi\bar{\theta}_1|_A$ and $\phi\bar{\theta}_1|_A$ has the

extension $\phi\bar{\theta}_1: I \times \Lambda \times M \rightarrow \Lambda \times M$. By [L, Lemma 1.1], $\bar{\phi}$ has also a Λ -f.p. extension $\psi: I \times \Lambda \times M \rightarrow \Lambda \times M$. This ψ is a wanted extension of ϕ .

Finally, it is easy to prove that $\hat{\psi}$ is $(I \times \Lambda)$ -f.p. homotopic to $\text{id}_I \times g: I \times \Lambda \times M \rightarrow I \times \Lambda \times M$. Therefore, $\hat{\psi}$ is an $(I \times \Lambda)$ -f.p. homotopy equivalence if g is a Λ -f.p. homotopy equivalence.

PROOF OF THEOREM 1. Since it is straightforward to verify that the translation map is a homeomorphism of $\text{Homeo}(M)$ (or refer to [H]), we only need to show that $\text{Homeo}(M)$ has the local UV^∞ -property at id_M . Let U be a given neighborhood of id_M in $\text{Homeo}(M)$ of the form $\bigcap \{[K_i, \Omega_i] \mid i = 1, \dots, p\}$ where K_i is a compact subset of the open subset Ω_i in M for each i .

Definition of V . Without loss of generality, we can assume that $K_1 \cup \dots \cup K_p \subset M_0$ [Ha, Lemma 2.4]. Then, since M_1 is locally convex, there is a finite cover $\mathcal{A}$ of M_0 consisting of convex open sets in M_1 such that

(α) For $E = R$, if $A \in \mathcal{A}$ with $A \cap M_0 \neq \emptyset$, then $A \subset N \times \text{Int } B_2$.

(β) If $A \in \mathcal{A}$ with $A \cap K_i \neq \emptyset$, then its closure $\bar{A} \subset \Omega_i$.

Note that any two $\mathcal{A}$ -close maps into M_1 are canonically $\mathcal{A}$ -homotopic. Let $\delta > 0$ be a Lebesgue number for the open cover $\{A \cap M_0 \mid A \in \mathcal{A}\}$ of M_0 , and let $\mathcal{F}$ be a finite closed cover of M_0 each of whose members has diameter less than δ . Since the union $\bigcup \{\bar{A} \mid A \in \mathcal{A} \text{ and } \bar{A} \subset \Omega_i\}$ is compact for each $i = 1, \dots, p$, there is a sequence of positive numbers $\varepsilon_2, \varepsilon_3, \dots$ such that for each $i = 1, \dots, p$ if $\bar{A} \subset \Omega_i$, then $A \times E_{\{\varepsilon_2, \varepsilon_3, \dots\}}^\infty \subset \Omega_i$. Now, for each $C \in \mathcal{F}$, since $\text{diam}(C) < \delta$, there is an $A_C \in \mathcal{A}$ such that $C \subset A_C$. Define

$$V = \bigcap \left\{ \left[C, A_C \times E_{\{\varepsilon_2, \varepsilon_3, \dots\}}^\infty \right] \mid C \in \mathcal{F} \right\}.$$

(For each $C \in \mathcal{F}$, $A_C \times E_{\{\varepsilon_2, \varepsilon_3, \dots\}}^\infty$ is an open set in M by use of (α).) Observe that V is an open neighborhood of id_M in U : Let $f \in V$. Fix an i ($i = 1, \dots, p$) and let $x \in K_i$. Then, there is a $C \in \mathcal{F}$ such that $x \in C$. So, $A_C \cap K_i \neq \emptyset$; consequently, it follows from (β) that $C \subset A_C \subset \bar{A}_C \subset \Omega_i$; then, $f(x) \in A_C \times E_{\{\varepsilon_2, \varepsilon_3, \dots\}}^\infty \subset \Omega_i$. Therefore, $f(K_i) \subset \Omega_i$ for each i ; so, $f \in U$. Moreover, that if $f \in \text{Homeo}(M)$, $g \in V$ and $f|_{M_0} = g|_{M_0}$, then $f \in V$.

The local strong UV^∞ -property. Let $\Lambda \in \mathcal{CW}(\mathcal{C}) \cup \mathcal{CW}(\mathcal{M})$ and Γ a closed G_δ -set in Λ . Let $g: \Lambda \rightarrow V$ be a continuous map with $g(\Gamma) = \{\text{id}_M\}$. We will prove that g is homotopic rel Γ in V to the constant map Id with $\text{Id}(\Lambda) = \{\text{id}_M\}$.

Let $f: \Lambda \times M \rightarrow \Lambda \times M$ be the Λ -f.p. map associated to g , defined by $f_\lambda(z) = g(\lambda)(z)$ for all $(\lambda, z) \in \Lambda \times M$, where $\Lambda \times M$ is the $\mathcal{CT}$ -product space as used in [L]. Then, f is continuous by [G, Theorem 8.17]. By [D, Theorem XII.3.1(1)] for the $\mathcal{CT}$ -product, we only need to prove that there is a Λ -f.p. isotopy (rel $\Gamma \times M$) $F: I \times \Lambda \times M \rightarrow I \times \Lambda \times M$ with $F_0 = f$, $F_1 = \text{id}_{\Lambda \times M}$ and $F_{(t, \lambda)} \in V$ for each $(t, \lambda) \in I \times \Lambda$. We will define by induction a sequence $\{F^n \mid n = 1, 2, \dots\}$ of Λ -f.p. invertible isotopies of $\Lambda \times M$ such that

- (0) $F_1^1 = \text{id}_M$ and $F_{(t, \lambda)}^1 \in V$ for each $(t, \lambda) \in I \times \Lambda$,
- (1) $_n$ $F_1^n = F_0^{n-1}$ ($n > 1$),
- (2) $_n$ $F_0^n \mid \Lambda \times M_{n-1} = f \mid \Lambda \times M_{n-1}$ ($n > 0$),
- (3) $_n$ $F_t^n \mid \Lambda \times M_{n-2} = f \mid \Lambda \times M_{n-2}$ for each $t \in I$, $n > 1$, and
- (4) $_n$ $F_t^n \mid \Gamma \times M = \text{id}_{\Gamma \times M}$.

Next, define $F: I \times \Lambda \times M \rightarrow I \times \Lambda \times M$ by

$$F(t, \lambda, z) = \begin{cases} (0, f(\lambda, z)) & \text{if } t = 0, \\ (t, F^n(2^n t - 1, \lambda, z)) & \text{if } 1/2^n \leq t \leq 1/2^{n-1}. \end{cases}$$

Then, F will be an $(I \times \Lambda)$ -f.p. isotopy. It is well defined by $(1)_n$ and continuous by $(2)_n$, $(3)_n$ and $I \times \Lambda \times M = \text{dirlim}(I \times \Lambda \times M_n)$ [L, Lemma 0.3]. It also is clear that $F_0 = f$ and that $F_1 = F_1^1 = \text{id}_{\Lambda \times M}$ by (0). Moreover, each $F_{(t, \lambda)} \in V$ by use of (0) and $(3)_n$. Therefore, the proof will be complete. (Note that F is not necessary an invertible isotopy.)

First, let $H: I \times \Lambda \times M_0 \rightarrow \Lambda \times M$ be the straight-line homotopy from $f|_{\Lambda \times M_0}$ to $i_{\Lambda \times M_0}$. Observe that H is a Λ -f.p. homotopy $\text{rel } \Gamma \times M_0$ with $H(t \times \lambda \times C) \subset A_C \times E_{\{\epsilon_2, \epsilon_3, \dots\}}^\infty$ for each $C \in \mathcal{F}$. Let $\hat{H}: I \times \Lambda \times M_0 \rightarrow I \times \Lambda \times M$ be the $(I \times \Lambda)$ -f.p. map defined by $\hat{H}(t, \lambda, x) = (t, H(t, \lambda, x))$. By [L, Lemma 1.2], we can assume that $\hat{H}$ is an $(I \times \Lambda)$ -f.p. embedding. Moreover, following the proof of [L, Lemma 1.2], to adjust only the $E_{\{\epsilon_2, \epsilon_3, \dots\}}^\infty$ -component of $\hat{H}(\text{rel } \partial I \times \Lambda \times M_0)$, we can assume that $\pi_{\Lambda \times M} \hat{H}$ is still a Λ -f.p. homotopy $\text{rel } \Gamma \times M_0$ with

$$\pi_M \hat{H}(t \times \lambda \times C) \subset A_C \times E_{\{\epsilon_2, \epsilon_3, \dots\}}^\infty$$

for each $C \in \mathcal{F}$ (*). By Lemma 2, $\hat{H}$ has an $(I \times \Lambda)$ -f.p. homotopy-equivalence extension $\hat{\psi}^1: I \times \Lambda \times M \rightarrow I \times \Lambda \times M$ such that

- (a)₁ $\hat{\psi}_1^1 = \text{id}_{\Lambda \times M}$,
- (b)₁ $\hat{\psi}_0^1 = f$, and
- (c)₁ $\hat{\psi}_t^1|_{\Gamma \times M} = i_{\Gamma \times M}$ for each $t \in I$.

Therefore, by [L, Lemma 3.3], there is an $(I \times \Lambda)$ -f.p. homeomorphism F^1 of $I \times \Lambda \times M$ such that F^1 is an extension of $\hat{\psi}^1|_{(I \times \Lambda \times M_0) \cup (I \times \Gamma \times M) \cup (1 \times \Lambda \times M)}$. Now, the properties $(2)_1$ and $(4)_1$ follow from $(b)_1$ and $(c)_1$, respectively. To verify (0), it is clear that $F_1^1 = \text{id}_{\Lambda \times M}$ by $(a)_1$, and that $F_{(t, \lambda)}^1 \in V$ by (*) since $\hat{\psi}^1$ is an extension of $\hat{H}$.

Second, consider f and F_0^1 . Since $f|_{\Lambda \times M_0} = F_0^1|_{\Lambda \times M_0}$, by use of Lemma 2, we can obtain a Λ -f.p. homotopy $\phi: f|_{\Lambda \times M_1} = F_0^1|_{\Lambda \times M_1} \text{ rel } (\Gamma \times M_1) \cup (\Lambda \times M_0)$. By [L, Lemma 1.2], we can assume that $\hat{\phi}: I \times \Lambda \times M_1 \rightarrow I \times \Lambda \times M$, defined by $\hat{\phi}(t, \lambda, z) = (t, \phi(t, \lambda, z))$, is an $(I \times \Lambda)$ -f.p. embedding. By Lemma 2, $\hat{\phi}$ has an $(I \times \Lambda)$ -f.p. homotopy equivalence extension $\hat{\psi}^2: I \times \Lambda \times M \rightarrow I \times \Lambda \times M$ such that

- (a)₂ $\hat{\psi}_1^2 = F_0^1$,
- (b)₂ $\hat{\psi}_0^2 = f$, and
- (c)₂ $\hat{\psi}_t^2|_{\Gamma \times M} = i_{\Gamma \times M}$.

Then, by [L, Lemma 3.3], there is an $(I \times \Lambda)$ -f.p. homeomorphism F^2 of $I \times \Lambda \times M$ such that F^2 is an extension of $\hat{\psi}^2|_{(I \times \Lambda \times M_1) \cup (I \times \Gamma \times M) \cup (1 \times \Lambda \times M)}$. The properties $(1)_2$, $(2)_2$ and $(4)_2$ follow from $(a)_2$, $(b)_2$ and $(c)_2$, respectively. The property $(3)_2$ holds true since $\hat{\psi}^2$ is an extension of $\hat{\phi}$.

Finally, in a similar manner, we can define a wanted $(I \times \Lambda)$ -f.p. homeomorphism F^n of $I \times \Lambda \times M$ when an appropriate invertible isotopy F^{n-1} has already been defined. Therefore, the proof is complete. $\square$

The author wishes to thank the referee for his suggestion to simplify the proof of Lemma 2.

REFERENCES

- [D] J. Dugundji, *Topology*, Allyn and Bacon, Boston, Mass., 1973.
- [G] B. Gray, *Homotopy theory*, Academic Press, New York, 1975.
- [H] M. Hale, Doctoral Thesis, Vanderbilt University, 1981.
- [Ha] V. L. Hansen, *Some theorems on direct limits of expanding sequences of manifolds*, Math. Scand. **29** (1971), 5–36.
- [He] R. E. Heisey, *Manifolds modelled on the direct limit of lines*, Pacific J. Math. **102** (1982), 47–54.
- [H-T] R. E. Heisey and H. Toruńczyk, *On the topology of direct limits of ANR's*, Pacific J. Math. **93** (1981), 307–312.
- [Hd] J. F. P. Hudson, *Piecewise linear topology*, Math. Lecture Note Ser., Benjamin, New York, 1969.
- [L] V. T. Liem, *On R^∞ (Q^∞)-manifold bundles over CW complexes*, Trans. Amer. Math. Soc. (to appear).
- [R] T. B. Rushing, *Topological embedding*, Academic Press, New York, 1973.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ALABAMA, UNIVERSITY, ALABAMA 35486