ON A QUESTION OF FAITH IN COMMUTATIVE ENDOMORPHISM RINGS

JOHN CLARK

ABSTRACT. Given a commutative ring R , let $Q(R)$ denote its maximal ring of quotients and, for any ideal I of R , let $\text{End}(I)$ denote the ring of R -endomorphisms of I . It is known that if $Q(R)$ is a self-injective ring then $\text{End}(I)$ is commutative for each ideal I of R . Carl Faith has asked if the converse holds. It does if R is either Noetherian or has no nontrivial nilpotent elements but here we produce an example to show that it does not hold in general.

Introduction. Throughout this paper R is a commutative ring with identity and $Q(R)$ will denote its maximal ring of quotients. The definition and fundamental properties of $Q(R)$ are to be found in, for example, the books of Lambek [7] and Stenström [9]. In particular, Theorem 4.1 on p. 279 of [9] shows that $Q(R)$ is a self-injective ring if and only if for every ideal I of R and every R -homomorphism $\phi: I \rightarrow R$ there exists a faithful ideal J of R and an R -homomorphism $\psi: J \rightarrow R$ such that $I \subseteq J$ and ψ extends ϕ .

For every ideal I of R we let $\text{End}(I)$ denote the ring of R -endomorphisms of I . In Proposition 1.2 of [3], Cox showed that $\text{End}(J)$ is commutative whenever J is a faithful ideal. It then follows easily from the previous paragraph that if $Q(R)$ is a self-injective ring then $\text{End}(I)$ is commutative for every ideal I of R . Carl Faith has asked, [4, p. 199, problem 10] and [5, p. 98, problem 6], if the converse holds. Alamelu, [1 and 2], and Cox [3] have shown that it does with some restrictions on R , in particular, if R is either Noetherian or has no nontrivial nilpotent elements. We produce below an example of a ring R which gives a negative answer to Faith's general question.

The example. Let A be a discrete valuation ring with maximal ideal At and quotient field K . Moreover, assume that A is not complete, for example A can be taken to be countable. Let M denote the A -module K/A and for each $n \in \mathbb{N}$ let M_n denote the A -submodule At^{-n}/A of M where $At^{-n} = \{at^{-n} : a \in A\} = \{ut^k : u \text{ is a unit in } A, k \in \mathbb{Z}, k \geq -n\}$. Then M is the direct union $\bigcup_{n \in \mathbb{N}} M_n$ and any proper nonzero A -submodule of M is M_n for some $n \in \mathbb{N}$.

For each $n \in \mathbb{N}$ let E_n denote the A -endomorphism ring of M_n and let E denote the A -endomorphism ring of M . Any f in E_n is determined by $f(t^{-n} + A)$, which is of the form $a(t^{-n} + A)$ for some a in A . This produces a ring isomorphism $\phi: E_n \rightarrow A/At^n$ given by $\phi(f) = a + At^n$.

Received by the editors April 30, 1985 and, in revised form, October 2, 1985.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 13B30, 13C11; Secondary 16A65.

Key words and phrases. Endomorphism ring, maximal quotient ring, self-injective, trivial extension.

If x is an arbitrary nonzero element of some M_n we can write $x = ut^{m-n} + A$ where u is a unit in A and $m < n$. Then if $g: M_n \rightarrow M$ is an A -homomorphism with $g(x) = vt^{-k} + A$, where v is a unit and $k \geq 0$, we have

$$0 = g(t^{n-m}x) = t^{n-m}g(x) = vt^{n-m-k} + A$$

and so $-k \geq m - n$. Hence $g(x) \in M_{n-m} \subseteq M_n$. This argument allows us to define, for any $m, n \in \mathbb{N}$ with $m \leq n$, the restriction functions $\phi_{mn}: E_m \rightarrow E_n$ by $\phi_{mn}(g) = g|_{M_n}$ and $\phi_n: E \rightarrow E_n$ defined similarly. This gives a projective system $(E_m, \phi_{mn})_{m,n \in \mathbb{N}}$ of rings and ring homomorphisms and it is readily checked that

$$E = \text{proj lim } E_m.$$

Since, from above, each E_n is isomorphic to the commutative ring A/At^n , it follows that E is a commutative ring (see, for example, Proposition 1.9 on p. 53 of Lafon [6]).

We now let R be the trivial extension of A by M . In other words, $R = A \oplus M$, made into a ring by defining addition componentwise and multiplication by

$$(a_1 + m_1)(a_2 + m_2) = a_1a_2 + a_1m_2 + a_2m_1$$

for all $a_1, a_2 \in A$, $m_1, m_2 \in M$. This procedure was called *the principle of idealization* by Nagata [8] since the ring R contains the ring A and M is an ideal of R , with $M^2 = 0$. We determine the ideals of R .

First, if S is any A -submodule of M , then, by our multiplication in R , S is an ideal of R . Moreover, any ideal of R contained in M is such a submodule. Thus the ideals of R contained in M are precisely 0 , M and the M_n . Furthermore, each M_n is just the principal ideal of R generated by $t^{-n} + A$. On the other hand, if r is an element of R not in M , say $r = ut^n + m$, where u is a unit in A , $n \geq 0$ and $m \in M$, then by the divisibility of the A -module $M = K/A$ we have $t^n M = M$. Using this, it is not difficult to show that the principal ideal Rr is $At^n \oplus M$ and coincides with Rt^n . In particular, we have that all ideals of R not contained in M are principal. Thus all the ideals of R except M are principal.

Now from the definition of R it is easily shown that the R -endomorphism ring of M coincides with its A -endomorphism ring, in other words $\text{End}_R(M) = E$. In particular, from above, the R -ideal M has a commutative endomorphism ring. Since all the other ideals of R are principal, they also have commutative endomorphism rings. Thus to establish our example it remains to show that $Q(R)$ is not self-injective.

In fact, $Q(R) = R$. To see this we first note that for any ideal I of R contained in M we have $MI = 0$, while if J is a proper ideal of R not contained in M then $J = At^n \oplus M$ for some $n > 0$ and so $M_n J = 0$. Thus the only faithful ideal in R is R itself. That $Q(R) = R$ now follows from Proposition 4 on p. 39 of Lambek [7].

Now let $\hat{A}$ denote the completion of A and choose an element b belonging to $\hat{A}$ but not to A . Then b can be represented by a formal power series

$$b = u_0 + u_1 + \cdots + u_n t^n + \cdots$$

where the u_i are elements of A coming from a fixed set of representatives for A modulo At and, since b is not in A , u_i is nonzero for infinitely many i . Then multiplication by b gives an R -endomorphism h of the ideal M which differs from any multiplication of M by an element from A . However, for any R -endomorphism

$k: R \rightarrow R$, if $k(1) = a + m$ where $a \in A$, $m \in M$, then the restriction of k to M is just multiplication by a . This shows that our homomorphism $h: M \rightarrow R$ does not extend to R . Hence R , and so $Q(R)$, is not self-injective.

ACKNOWLEDGEMENT. It is a pleasure to acknowledge the referee for suggestions that improved the presentation.

REFERENCES

1. S. Alamelu, *On commutativity of endomorphism rings of ideals*, Proc. Amer. Math. Soc. **37** (1973), 29–31.
2. ———, *Commutativity of endomorphism rings of ideals. II*, Proc. Amer. Math. Soc. **55** (1976), 271–274.
3. S. H. Cox, Jr., *Commutative endomorphism rings*, Pacific J. Math. **45** (1975), 87–91.
4. C. Faith, *Injective quotient rings of commutative rings*, Module Theory (Proc. Special Session, Amer. Math. Soc., Univ. Washington, Seattle, Wash. 1977), Lecture Notes in Math., vol. 700, Springer-Verlag, Berlin, 1979, pp. 151–203.
5. ———, *Injective modules and injective quotient rings*, Lecture Notes in Pure and Appl. Math, vol. 72, Dekker, New York, 1982.
6. J.-P. Lafon, *Les formalismes fondamentaux de l'algèbre commutative*, Hermann, Paris, 1974.
7. J. Lambek, *Lectures on rings and modules*, Blaisdell, Waltham, Mass., 1966.
8. M. Nagata, *Local rings*, Interscience, New York and London, 1962.
9. B. Stenström, *Rings of quotients*, Springer-Verlag, Berlin and New York, 1975.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF OTAGO, P.O. BOX 56, DUNEDIN, NEW ZEALAND