A WEAKLY INFINITE-DIMENSIONAL SPACE WHOSE PRODUCT WITH THE IRRATIONALS IS STRONGLY INFINITE-DIMENSIONAL

ELŻBIETA POL

ABSTRACT. We give an example of a weakly infinite-dimensional space X such that the product $X \times B$ of X and a subspace B of the irrationals is strongly infinite-dimensional; under the assumption of the Continuum Hypothesis, B can be the irrationals. This example answers a question of Addis and Gresham [AG].

1. Terminology and notation. All spaces under discussion are metrizable and separable. Our terminology follows [AP and E]. We denote by I the real interval $[0, 1]$, by C the usual Cantor set in I and by I^ω the Hilbert cube. We denote the space of the irrational numbers from I by P and the rational numbers from I by Q .

A space X is weakly infinite-dimensional [AP, Chapter 10, §§4–7] if for every sequence $\{(A_1, B_1), (A_2, B_2), \dots\}$ of pairs of closed disjoint subsets of X there are partitions L_i in X between A_i and B_i such that $\bigcap_{i=1}^\infty L_i = \emptyset$. Otherwise, X is strongly infinite-dimensional.

A space X is called a C -space if for every sequence $\mathcal{G}_1, \mathcal{G}_2, \dots$ of open covers of X there exists a sequence $\mathcal{U}_1, \mathcal{U}_2, \dots$ of families of open subsets of X such that, for $i = 1, 2, \dots$,

- (i) the members of $\mathcal{U}_i$ are pairwise disjoint,
- (ii) each member of $\mathcal{U}_i$ is contained in a member of $\mathcal{G}_i$,
- (ii) the union $\bigcup_{i=1}^\infty \mathcal{U}_i$ covers X .

The notion of C -space was introduced by W. Haver in [H] for metric space and by D. Addis and J. Gresham in [AG] for general topological spaces.

LEMMA 1 [AG]. *Every C -space is weakly infinite-dimensional.*

2. Results. The aim of this note is to construct the following examples.

EXAMPLE 1. *There exists a weakly infinite-dimensional space X such that the product $X \times B$ of X with a certain subspace B of the space of irrationals is strongly infinite-dimensional. Moreover, X is a C -space while $X \times B$ is not a C -space.*

EXAMPLE 2. *Under the assumption of the Continuum Hypothesis there exists a weakly infinite-dimensional space X such that the product $X \times P$ of X with the space of irrationals P is strongly infinite-dimensional. Moreover, X is a C -space while $X \times P$ is not a C -space.*

Received by the editors June 18, 1985.

1980 *Mathematics Subject Classification.* Primary 54F45, 54B10.

Key words and phrases. Weakly infinite-dimensional spaces, products, property C .

©1986 American Mathematical Society
 0002-9939/86 \$1.00 + \$.25 per page

Example 2 gives, in particular, an answer to a question formulated by Addis and Gresham in [AG] as to whether the product of a C -space and the space of irrationals is a C -space.

The first example of two metrizable separable weakly infinite-dimensional spaces X_1 and X_2 (being also C -spaces) whose product $X_1 \times X_2$ is strongly infinite-dimensional was given by R. Pol in [P2] (see [EP, Example 8.20] for the proof that X_i are C -spaces).

3. Constructions. Our constructions are based on the following lemma, which follows an idea of Michael [M].

LEMMA 2. *Let $f: X \rightarrow Y$ be a mapping of a space X into a space Y and let $A = f^{-1}(B)$ for some $B \subset Y$. If the subset $\text{Graph}(f|A) = \{(x, f(x)): x \in A\}$ of $X \times Y$ is strongly infinite-dimensional, then the product $X \times B$ is strongly infinite-dimensional (and thus is not a C -space).*

The proof of this lemma follows from the fact that a closed subspace of a weakly infinite-dimensional space is weakly infinite-dimensional and that $X \times B$ contains $\text{Graph}(f|A)$ as a closed subset.

Construction of Example 1. Let T be the weakly infinite-dimensional compactum, which is not countable dimensional defined by R. Pol in [P1]. The space T has the following structure: $T = Y \cup Z$, where Y is a completely metrizable totally disconnected space, which is strongly infinite-dimensional, and Z is the union of countably many 0-dimensional sets $Z_1, Z_2, \dots$. Moreover, we can assume (see [P1, Comment B]) that T is embedded in the product $C \times I^\omega$ of the Cantor set C and the Hilbert cube I^ω , in such a way that Y is the graph of a certain first-Baire-class function $f: C \rightarrow I^\omega$, i.e., $Y = \{(t, f(t)): t \in C\} \subset C \times I^\omega$.

Split the Cantor set C into two disjoint Bernstein sets B_1 and B_2 (see [K, §40]), i.e., all compact subsets of B_i are countable. Since Y is strongly infinite-dimensional, either $Y_1 = \{(t, f(t)): t \in B_1\}$ or $Y_2 = \{(t, f(t)): t \in B_2\}$ is strongly infinite-dimensional. Suppose that Y_2 is strongly infinite-dimensional and let X be the following subspace of $C \times C \times I^\omega$: $X = (B_1 \times T) \cup \{(t, t, f(t)): t \in B_2\} \subset C \times T$. Denote $X_1 = (B_1 \times T)$, $X_2 = \{(t, t, f(t)): t \in B_2\}$. The set X_2 is obviously homeomorphic to Y_2 , hence it is strongly infinite dimensional. We will show that

(1) X is a C -space.

Suppose that $\mathcal{G}_1, \mathcal{G}_2, \dots$ is a sequence of open covers of X . Fix an $i \geq 3$. Since the set $B_1 \times Z_{i-2}$ is 0-dimensional, there exists a family $\mathcal{W}_i$ of pairwise disjoint sets open in $C \times T$ such that $B_1 \times Z_{i-2} \subset \bigcup \mathcal{W}_i = W_i$ and for each $W \in \mathcal{W}_i$ the set $W \cap X$ is contained in some member of $\mathcal{G}_i$. Let $F = (B_1 \times T) \setminus \bigcup_{i=3}^\infty W_i$; we will show that $\dim F = 0$. Since $\bigcup_{i=3}^\infty W_i \supset B_1 \times Z$, F is a closed subset of $C \times T$ contained in $B_1 \times Y$. Consider the projection $p_1: B_1 \times T \rightarrow B_1$. Since T is compact, p_1 , as well as $p = p_1/F: F \rightarrow B_1$, is a closed mapping. Moreover, for each $t \in B_1$ the set $p^{-1}(t) = (\{t\} \times T) \cap F$ is a compact set contained in $\{t\} \times Y$. It follows that $\dim p^{-1}(t) = 0$, because Y is totally disconnected. Thus, by the Hure-

wicz Theorem (see [E, Theorem 1.12.4])

$$\dim F \leq \sup\{\dim p^{-1}(t): t \in B_1\} + \dim B_1 = 0.$$

Now, let $\mathcal{W}_2$ be a family of pairwise disjoint open subsets of $C \times T$ such that $F \subset \bigcup \mathcal{W}_2 = W_2$ and for each $W \in \mathcal{W}_2$ there exists $U \in \mathcal{G}_2$ such that $W \cap X \subset U$. Put $K = X \setminus \bigcup_{i=2}^\infty W_i$; since $\bigcup_{i=2}^\infty W_i \supset B_1 \times T$, then $K \subset X_2$. We shall show that the set K is countable. Let $p_2: C \times T \rightarrow C$ be the projection. Since p_2 is a closed mapping, the set $L = p_2((C \times T) \setminus \bigcup_{i=2}^\infty W_i)$ is closed in C . Moreover, the set L is contained in B_2 , hence it is countable. It follows that the set $p_2(K) \subset L$ is countable, which implies that K is countable, because $p_2|_{X_2}: X_2 \rightarrow B_2$ is a one-to-one mapping. Let $\mathcal{U}_1$ be a family of pairwise disjoint open subsets of X such that $K \subset \bigcup \mathcal{U}_1$ and each member of $\mathcal{U}_1$ is contained in some member of $\mathcal{G}_1$. Then the family $\mathcal{U}_1, \mathcal{U}_2, \dots$, where $\mathcal{U}_i = \{W \cap X: W \in \mathcal{W}_i\}$ for $i \geq 2$, satisfies conditions (i), (ii) and (iii).

Finally, let $f: X \rightarrow C$ be the projection. Then $f^{-1}(B_2) = X_2$ and the $\text{Graph}(f|_{X_2})$ is homeomorphic to X_2 in a natural way, hence it is strongly infinite-dimensional. It follows by Lemma 2 that the product $X \times B_2$ is strongly infinite-dimensional. Since B_2 is a 0-dimensional metrizable separable space, it is homeomorphic to a subset B of the irrationals P .

Construction of Example 2. In this example we use, among others, some ideas of [P2].

Let $Z = \prod_{i=0}^\infty I_i$, where $I_i = I$ for $i = 0, 1, 2, \dots$, be the Hilbert cube and the mappings $p_0: Z \rightarrow I_0$ and $p_n: \prod_{i=1}^\infty I_i \rightarrow \prod_{i=1}^n I_i$ be appropriate projections. Arrange all rational numbers in I_0 into a sequence $q_1, q_2, \dots$. Denote by T a compactum obtained from Z by attaching to each compactum $\{q_n\} \times \prod_{i=1}^\infty I_i$ the n -cube $\prod_{i=1}^n I_i$ by the map p_n , i.e., T is a quotient space defined by an upper semicontinuous decomposition consisting of singletons $\{(t, x)\}$, where $t \in P$ and $x \in \prod_{i=1}^\infty I_i$, and the sets $\{q_n\} \times p_n^{-1}(y)$, where $y \in \prod_{i=1}^n I_i$. Let $\pi: Z \rightarrow T$ be the natural quotient mapping and $X_1 = \pi(Q \times \prod_{i=1}^\infty I_i)$.

Using the Continuum Hypothesis, define a transfinite sequence $G_1 \supset G_2 \supset \dots \supset G_\alpha \supset \dots$, $\alpha < \omega_1$, of G_δ -subsets of T containing X_1 and such that for every G_δ -set G in T containing X_1 there exists α with $G_\alpha \subset G$ (ω_1 denotes the first uncountable ordinal). Again by CH, we can arrange all continua in Z intersecting $\{0\} \times \prod_{i=1}^\infty I_i$ and $\{1\} \times \prod_{i=1}^\infty I_i$ into a sequence $K_1, K_2, \dots, K_\alpha, \dots$, $\alpha < \omega_1$.

By transfinite induction we will choose, for each $\alpha < \omega_1$, a point $y_\alpha \in K_\alpha \cap \pi^{-1}(G_\alpha) \cap p_0^{-1}(P)$ such that $y_\alpha \notin \{y_\beta: \beta < \alpha\}$. Suppose that $\alpha = 0$ or that we have already defined all points y_β for $\beta < \alpha$. Since $\pi^{-1}(G_\alpha)$ is a G_δ -set in Z containing $Q \times \prod_{i=1}^\infty I_i$, $M = p_0(Z \setminus \pi^{-1}(G_\alpha))$ is an F_σ -set in I_0 , contained in P (notice that the projection p_0 is closed). Thus the set $P \setminus M$ is of second category in I_0 . Hence there exists $p_\alpha \in (P \setminus M) \setminus \{p_0(y_\beta): \beta < \alpha\}$. Since $\{p_\alpha\} \times \prod_{i=1}^\infty I_i \subset \pi^{-1}(G_\alpha)$ and the continuum K_α intersects $\{p_\alpha\} \times \prod_{i=1}^\infty I_i$, there exists $y_\alpha \in K_\alpha \cap \pi^{-1}(G_\alpha) \cap p_0^{-1}(p_\alpha)$. If we put $x_\alpha = \pi(y_\alpha)$ for every $\alpha < \omega_1$, then

$$x_\alpha \in G_\alpha \cap \pi(K_\alpha) \setminus (X_1 \cup \{x_\beta: \beta < \alpha\}).$$

Claim. The subspace $X = X_1 \cup \{x_\alpha: \alpha < \omega_1\}$ of T has the desired properties. We will show that

(2) X is a C -space,

and

(3) the subspace $X_2 = \{x_\alpha: \alpha < \omega_1\}$ of X is strongly infinite dimensional.

To show (2) suppose that $\mathcal{G}_1, \mathcal{G}_2, \dots$ is a sequence of open covers of X . Decompose the set X_1 into a sequence $Z_1, Z_2, \dots$ of 0-dimensional sets. For $i = 2, 3, \dots$ choose a family $\mathcal{W}_i$ of disjoint open subsets of T such that $Z_{i-1} \subset W_i = \bigcup \mathcal{W}_i$ and for each $U \in \mathcal{W}_i$ there exists $G \in \mathcal{G}_i$ satisfying $U \cap X \subset G$ (see [E, Lemma 1.7.3]). Then $X_1 \subset \bigcup_{i=2}^\infty W_i$ and there exists α such that $G_\alpha \subset \bigcup_{i=2}^\infty W_i$. The set $X \setminus \bigcup_{i=2}^\infty W_i \subset X \setminus G_\alpha$ is contained in $\{x_\beta: \beta < \alpha\}$, hence it is countable. Thus there exists a family $\mathcal{U}_1$ of open subsets of X satisfying conditions (i) and (ii) for $i = 1$ and such that $U_1 = \bigcup \mathcal{U}_1 \supset X \setminus \bigcup_{i=2}^\infty W_i$. The families $\mathcal{U}_1, \mathcal{U}_2, \dots$, where $\mathcal{U}_i = \{W \cap X: W \in \mathcal{W}_i\}$ for $i \geq 2$, satisfy conditions (i), (ii), and (iii).

To prove (3) we will show that the subspace $\pi^{-1}(X_2)$ of the Hilbert cube, which is homeomorphic to X_2 , is strongly infinite dimensional. For $n = 1, 2, \dots$, let $A_n = \{\{x_i\} \in \prod_{i=0}^\infty I_i: x_n = 0\}$ and $B_n = \{\{x_i\} \in \prod_{i=0}^\infty I_i: x_n = 1\}$ be opposite faces of the Hilbert cube Z and let U_i and V_i be open subsets of Z such that $A_i \subset U_i$, $B_i \subset V_i$, and $\bar{U}_i \cap \bar{V}_i = \emptyset$. For $i = 1, 2, \dots$, let L_i be an arbitrary partition between $\bar{U}_i \cap \pi^{-1}(X_2)$ and $\bar{V}_i \cap \pi^{-1}(X_2)$ in $\pi^{-1}(X_2)$. The partition L_i can be extended (see [E, Lemma 1.2.9]) to a partition L'_i between A_i and B_i in Z satisfying $L'_i \cap \pi^{-1}(X_2) \subset L_i$. Since $\bigcap_{i=1}^\infty L'_i$ contains a continuum connecting $\{0\} \times \prod_{i=1}^\infty I_i$ and $\{1\} \times \prod_{i=1}^\infty I_i$ (see [RSW, Lemma 5.2]), say K_α , we have $y_\alpha \in \bigcap_{i=1}^\infty L'_i \cap \pi^{-1}(X_2)$. Therefore $\bigcap_{i=1}^\infty L_i \neq \emptyset$. This ends the proof of (3).

Finally, let $f: X \rightarrow I_0$ be the projection. Then $f^{-1}(P) = X_2$. Moreover, $\text{Graph}(f|X_2)$ is homeomorphic to X_2 , hence it is strongly infinite-dimensional. Thus, by Lemma 2, the product $X \times P$ is strongly infinite-dimensional.

REFERENCES

- [AG] D. F. Addis and J. H. Gresham, *A class of infinite-dimensional spaces. Part I: Dimension theory and Alexandroff's problem*, Fund. Math. **101** (1978), 195–205.
- [AP] P. S. Alexandroff and B. A. Pasynkov, *Introduction to dimension theory*, Nauka, Moskva, 1973. (Russian)
- [E] R. Engelking, *Dimension theory*, PWN, Warszawa, 1978.
- [EP] R. Engelking and E. Pol, *Countable-dimensional spaces: A survey*, Dissertationes Math. **216** (1983), 1–41.
- [H] W. E. Haver, *A covering property for metric spaces*, Lecture Notes in Math., vol. 375, Springer-Verlag, Berlin and New York, 1974, pp. 108–113.
- [K] K. Kuratowski, *Topology*, vols. I, II, PWN, Warszawa, 1966, 1968.
- [M] E. Michael, *The product of a normal space and a metric space need not be normal*, Bull. Amer. Math. Soc. **69** (1963), 375–376.
- [P1] R. Pol, *A weakly infinite-dimensional compactum which is not countable-dimensional*, Proc. Amer. Math. Soc. **82** (1981), 634–636.
- [P2] ———, *A remark on A -weakly infinite-dimensional spaces*, Topology Appl. **13** (1982), 97–101.
- [RSW] L. R. Rubin, R. M. Shori, and J. J. Walsh, *New dimension-theory techniques for constructing infinite-dimensional examples*, General Topology Appl. **10** (1979), 93–103.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WARSAW, PALAC KULTURY I NAUKI, 00-901 WARSZAWA, POLAND