A JC -ALGEBRA WHICH IS NOT THE RANGE OF A POSITIVE PROJECTION ON A C^* -ALGEBRA

A. GUYAN ROBERTSON

ABSTRACT. An example is given of a JC -algebra which is not the range of any positive projection on the selfadjoint part of its generated C^* -algebra.

JC -algebras arise naturally in the study of positive projections on operator algebras [1, Theorem 1.4]. Effros and Størmer showed in [1, Theorem 2.1] that if A is a simple JC -algebra and B is the C^* -algebra generated by A , then there exists a positive projection P on B such that $P(B_{sa}) = A$. Furthermore they expressed the belief that this result may fail for general JC -algebras [1, Remark 2.7]. The purpose of this note is to provide an explicit example showing that this is indeed the case.

The example is constructed as follows. Let F be the CAR -algebra, acting irreducibly on a Hilbert space H . By [2, Theorem 6.2.2], F is the C^* -algebra generated by an infinite-dimensional spin factor V . If K denotes the algebra of compact operators on H , then $F \cap K = \{0\}$ [3, Theorem 6.5.7]. Let $A = K_{sa} + V$ and let B be the C^* -algebra generated by A . Then A is a JC -algebra and $B = K + F$. These statements follow easily from [3, Corollary 1.5.8] and its Jordan-algebraic analogue [2, 3.4].

THEOREM. *There is no positive projection P on B such that $P(B_{sa}) = A$.*

PROOF. Suppose that such a P exists. We show that this implies that V is *reversible* in the sense that it is closed under symmetric products of the form $a_1 a_2 \cdots a_n + a_n \cdots a_2 a_1$ [2, 2.3.2]. This will contradict [2, Theorem 6.2.5], which asserts that V is not reversible.

Given $a_1, \dots, a_n \in V$, let $a = a_1 a_2 \cdots a_n + a_n \cdots a_2 a_1$. If $b \in K_{sa}$ then $a \circ b = \frac{1}{2}(ab + ba) \in K_{sa}$. Therefore $a \circ b = P(a \circ b) = P(a) \circ b$, where the last equality follows from [1, Lemma 1.1] together with the fact that the range of P is assumed to be a Jordan algebra. Now the identity operator is a strong limit of elements of K_{sa} . Hence $a = P(a) \in A$. It follows that $a = k + x$, where $k \in K_{sa}$ and $x \in V$. Since it is obvious from its definition that $a \in F$, we have

$$a - x = k \in F \cap K = \{0\}.$$

Therefore $a = x \in V$. This shows that V is reversible, contradicting [2, Theorem 6.2.5].

REMARK. The argument above actually shows that A is not the range of a positive projection on any reversible JC -algebra containing A as a JC -subalgebra.

Received by the editors July 16, 1985.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 46L05; Secondary 17C65.

©1986 American Mathematical Society
 0002-9939/86 \$1.00 + \$.25 per page

REFERENCES

1. E. G. Effros and E. Størmer, *Positive projections and Jordan structure in operator algebras*, Math. Scand. **45** (1979), 127–138.
2. H. Hanche-Olsen and E. Størmer, *Jordan operator algebras*, Pitman, London, 1984.
3. G. K. Pedersen, *C^* -algebras and their automorphism groups*, Academic Press, London, 1979.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF EDINBURGH, EDINBURGH EH9 3JZ,
SCOTLAND