APPROXIMATION BY RATIONAL FUNCTIONS

RONALD A. DEVORE

ABSTRACT. Making use of the Hardy-Littlewood maximal function, we give a new proof of the following theorem of Pekarski: If f' is in $L \log L$ on a finite interval, then f can be approximated in the uniform norm by rational functions of degree n to an error $O(1/n)$ on that interval.

It is well known that approximation by rational functions of degree n can produce a dramatically smaller error than that for polynomials of degree n . The best example of this is Newman's theorem [3] which shows that the function $f(x) = |x|$ can be approximated on $[-1, 1]$ by rational functions of degree n to an error $O(\exp(-c\sqrt{n}))$, whereas for polynomials of degree n the error is known to be larger than c/n . Other authors have shown that such improvement also occurs for certain classes of functions. For example, V. Popov [5] showed that if $f' \in L_p[0, 1]$, with $p > 1$, then $r_n(f) = O(n^{-1})$ where $r_n(f)$ is the error in approximating f by rational functions R of degree at most n in the *uniform norm*:

$$r_n(f) := \inf_{\deg(R)=n} \|f - R\|_\infty[0, 1].$$

To obtain this order of approximation for polynomials requires roughly speaking that $f' \in L_\infty$. A striking limiting version of Popov's result was given by A. A. Pekarski [4], who showed that the same conclusion holds when $f' \in L \log L$, i.e. if $|f'| \log(1 + |f'|)$ is integrable.

The Popov and Pekarski proofs of these theorems are quite technical, and it was the purpose of [2] to introduce an elementary technique using maximal functions and partitions of unity for rational functions in order to give a simpler proof of Popov's results. The point of this note is to show that a modification of the technique in [2], albeit a little tricky, will also prove Pekarski's theorem.

The idea in [2] is to partition $[0, 1]$ into a set I of disjoint intervals I and construct associated rational functions ψ_I which form a partition of unity: $\sum_{I \in I} \psi_I \equiv 1$. Our rational approximation R is then given by

$$(1) \quad R(x) := \sum_{I \in I} f(x_I) \psi_I(x)$$

with x_I the center of I . Of course, the intervals I depend on f .

The rational functions ψ_I are constructed using a standard method for partitions of unity. Namely, $\psi_I := \phi_I / \Phi$ with $\Phi := \sum \phi_I$. In the case of Popov's theorem, the ϕ_I depend only on the interval I and all can be taken of degree 4. The intervals

Received by the editors October 3, 1985.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 41A20, 41A25, 41A63, 42B25.

Supported by the National Science Foundation Grant DMS 8320562.

©1986 American Mathematical Society
0002-9939/86 \$1.00 + \$.25 per page

I are determined by using the Hardy-Littlewood maximal functions M which is defined for $g \in L_1$ by

$$Mg(x) := \sup_{J \ni x} \frac{1}{|J|} \int_J |g|,$$

where the sup is taken over all intervals $J \subset [0, 1]$ which contain x .

To prove the Pekarski theorem, we will need to let the degree of ϕ_I depend on f . The desired properties of ϕ_I are given in the following lemma.

LEMMA 1. *For each even integer $m \geq 8$, and each interval I there is a nonnegative rational function ϕ_I of degree at most $6m$ with the following properties:*

- (i) $\phi_I(x) \geq 1$, $x \in I$,
- (ii) $\phi_I(x) \leq 8 \cdot 2^{-\sqrt{\lambda}/4}$, if $2^{-m/\lambda}|I| \leq \text{dist}(x, I) \leq 1/2$ and $0 < \lambda < m$,
- (iii) $\phi_I(x) \leq 4(a^2 + 1)^{-m}$, if $\text{dist}(x, I) \geq a|I|$ and $a > 0$.

We postpone the proof of this lemma until the end of the paper. We now use this result to prove the following.

THEOREM. *There is an absolute constant $c > 0$ such that for $n = 1, 2, \dots$*

$$r_n(f) \leq c \|M(f')\|_1 n^{-1}, \quad n = 1, 2, \dots,$$

whenever $M(f')$ is in $L_1[0, 1]$.

REMARK: It is well known (see e.g. [1]) that $g \in L \log L$ is equivalent to $M(g) \in L_1$ and therefore this theorem is equivalent to Pekarski's.

PROOF. It is enough to consider functions f with $\|M(f')\|_1 = 1$. It follows that $\|f'\|_1 \leq 1$ and hence there is a collection $\mathcal{I}$ of at most n intervals I which are a disjoint partition of $[0, 1]$ and satisfy

$$(2) \quad \frac{1}{n} \leq \int_I |f'| \leq \frac{2}{n}, \quad I \in \mathcal{I}.$$

For each $I \in \mathcal{I}$, we let m_I be the smallest integer which is both larger than 7 and also larger than $4n \int_I M(f')$. If ϕ_I is the function of Lemma 1 for the interval I and for $m = m_I$, we let $\Phi := \sum_{I \in \mathcal{I}} \phi_I$. By Lemma 1, $\Phi \geq 1$, on $[0, 1]$ and hence the functions ψ_I satisfy

$$(3) \quad \psi_I(x) \leq \phi_I(x), \quad 0 \leq x \leq 1.$$

We now take R as in (1) with x_I the center of I . Since $\sum m_I \leq 16n$, R has degree $\leq 96n$. To estimate $|f(x) - R(x)|$, we let I_0 denote the interval of $\mathcal{I}$ which contains x ; I_1 the interval of $\mathcal{I}$ immediately to the right of I_0 ; I_{-1} the interval immediately to the left of I_0 ; and so on. We have

$$(4) \quad f(x) - R(x) = \sum_{I \in \mathcal{I}} (f(x) - f(x_I)) \psi_I(x) =: \sum_{-1} + \sum_0 + \sum_1$$

Where $\sum_{-1}$ denotes the sum over those $I = I_k$ with $k < -1$, $\sum_1$ the sum over those $I = I_k$ with $k > 1$ and $\sum_0$ the sum of the terms $k = -1, 0, 1$. Clearly, $|f(x) - f(x_{I_k})| \leq 2(|k| + 1)/n$. Since the ψ_I are nonnegative and add up to one, we have

$$(5) \quad \sum_0 \leq 12/n.$$

The estimates for $\sum_{-1}$ and $\sum_1$ are the same and therefore we estimate only $\sum_1$. For this, we fix $k > 1$ and estimate the term in $\sum_1$ corresponding to $I = I_k$. We have

$$(6) \quad e_k := |f(x) - f(x_I)|\psi_I(x) \leq \frac{2(k+1)}{n}\psi_I(x) \leq \frac{4k}{n}\phi_I(x).$$

We write $\text{dist}(x, I) =: a|I|$, with $a \geq 0$, and we consider three cases.

Case $a \geq \sqrt{k}$. Then since $m \geq 8$, by (iii) of Lemma 1, we have $\psi_I(x) \leq \phi_I(x) \leq 4k^{-4}$ and consequently

$$(7) \quad e_k \leq 16k^{-3}n^{-1}.$$

Case $1/2 \leq a \leq \sqrt{k}$. The smallest interval J which contains x and I has length $(a+1)|I|$ and on I ,

$$M(f') \geq \frac{1}{|J|} \int_J |f'| \geq \frac{k}{n(a+1)|I|}$$

and therefore $m \geq 4n \int_I M(f') \geq 4k/(a+1) \geq \sqrt{k}$. This gives by (iii) of Lemma 1,

$$(8) \quad e_k \leq \frac{4k}{n}\phi_I(x) \leq \frac{4k}{n}(a^2+1)^{-m} \leq \frac{16k}{n}(5/4)^{-\sqrt{k}}.$$

Case $0 < a < 1/2$. We write $a =: 2^{-m/\lambda}$ with $0 < \lambda < m$. Similar to the second case, for $u \in I$, we have $M(f')(u) \geq (k-1)/n(u-x)$. Therefore,

$$m \geq 4n \int_I M(f') \geq 4(k-1) \int_{2^{-m/\lambda}|I|}^{|I|} \frac{du}{u} \geq 2k \left(\frac{m}{\lambda} \right) \log 2.$$

This shows that $\lambda \geq 2k \log 2 \geq k$. Hence by (ii) of Lemma 1, we have

$$(9) \quad e_k \leq \frac{4k}{n}\phi_I(x) \leq \frac{32k}{n}2^{-\sqrt{k}/4}.$$

The estimates (7)–(9) serve to show that $\sum_1 = \sum e_k \leq cn^{-1}$, with c an absolute constant. This combined with (5) and the corresponding estimate for $\sum_{-1}$ when placed in (4) proves the theorem.

We turn now to the proof of Lemma 1. For this, we shall use the following:

LEMMA 2. *For each even integer $m \geq 8$ there is a rational function R of degree $\leq 2m$ with the following properties:*

- (i) $R(x) \geq 1$, $x \in [-1, 0]$,
- (ii) $0 \leq R(x) \leq 2$, for $-\infty < x < \infty$,
- (iii) $|R(x)| \leq 2 \cdot 2^{-m/4j}$, if $2^{-(j+1)^2/m} \leq x \leq 1/2$, with $\sqrt{m} - 1 \leq j < m$.

PROOF. With $a := 2^{-1/m}$ and $a_k := a^{k^2}$, we define $p(x) := \prod_1^m (x + a_k)$. We first estimate $\pi(x) := p(-x)/p(x) = \prod_1^m (-x + a_k)/(x + a_k)$ when $x \geq 0$. Since each term in π has absolute value at most 1, we have

$$(10) \quad |\pi(x)| \leq 1, \quad x \geq 0.$$

When $a_m \leq x \leq 1/2$, we take j so that $a_{j+1} \leq x \leq a_j$; so $\sqrt{m} - 1 \leq j < m$. Then,

$$|\pi(x)| \leq \pi_1(x) := \prod_1^j \frac{a_k - x}{a_k + x}.$$

We now use the inequality $(1-t)/(1+t) \leq e^{-2t}$, which is valid for $0 \leq t \leq 1$. This gives

$$(11) \quad |\pi(x)| \leq \prod_1^j \frac{1-x/a_k}{1+x/a_k} \leq \exp \left(-2 \sum_1^j \frac{a_{j+1}}{a_k} \right) =: \exp(-2\sigma(j)).$$

Since $(j+1)^2 - k^2 \leq (j-k+1)(2j+1)$, we have with $b := a^{2j+1}$,

$$\sigma(j) \geq \sum_1^j b^\nu = b \frac{1-b^j}{1-b}.$$

But, since $\sqrt{m}-1 \leq j < m$, $b \geq 1/4$; $1-b^j \geq 1/2$; also $1-e^{-t} \leq t$, for $0 < t \leq 1$. Hence,

$$\sigma(j) \geq \frac{m}{8(2j+1)\log 2}.$$

Since $2\log 2 \leq 1/\log 2$, using our last estimate for $\sigma(j)$ in (11) gives

$$(12) \quad |\pi(x)| \leq \exp \left(\frac{-m \log 2}{2(2j+1)} \right) \leq 2^{-m/8j}, \quad a_{j+1} \leq x \leq a_j, \text{ for } \sqrt{m}-1 \leq j < m.$$

We can now take $R(x) := 2\pi^2(x)/(1+\pi^2(x))$. Since $\pi(-x) = 1/\pi(x)$ and $R(-x) = 2/(1+\pi^2(x))$, (i) follows from (10). The estimate (ii) is obvious, while (iii) follows immediately from (12).

PROOF OF LEMMA 1. It is enough to consider $I = [-1, 0]$ since the lemma then follows for any other interval by a change of scale. We let

$$T(x) := ((x+1/2)^2 + 3/4)^{-m}$$

and R be as in Lemma 2. We can then take $\phi(x) := R(x)R(-1-x)T(x)$. Since $T(x) \geq 1$, $x \in I$, (i) follows from (i) of Lemma 2. Since $T(x) \leq 1$, $x \notin I$, (ii) follows when $\lambda \leq 4$ from Lemma 2(ii). For the other values of λ , we choose j so that $j^2 < m^2/\lambda < (j+1)^2$, and then (ii) follows from Lemma 2(ii), (iii). Finally, if $\text{dist}(x, I) \geq a$, then $(x-1/2)^2 + 3/4 \geq (a+1/2)^2 + 3/4 \geq a^2 + 1$ and therefore (iii) follows from (ii) of Lemma 2.

REFERENCES

1. C. Bennett and R. Sharpley, *Weak type inequalities for H_p and BMO*, Harmonic Analysis in Euclidean Spaces, Proc. Sympos. Pure Math., vol. 35, Amer. Math. Soc., Providence, R.I., 1979, pp. 201-229.
2. R. DeVore, *Maximal functions and their application to rational approximation*, Approximation Theory, CMS Conf. Proc., vol. 3, Amer. Math. Soc., Providence, R.I., 1983, pp. 143-155.
3. D. J. Newman, *Rational approximation to $|x|$* , Michigan Math. J. **11** (1964), 11-14.
4. A. A. Pekarski, *Rational approximation of absolutely continuous functions with derivatives in an Orlicz space*, Math. Sb. **45** (1983), 121-137.
5. V. Popov, *On the connection between rational uniform approximation and polynomial L_p approximation of functions*, Quantitative Approximation, Academic Press, New York, 1980, pp. 119-129.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTH CAROLINA, COLUMBIA, SOUTH CAROLINA 29208