SOME NEW EXAMPLES OF NONORIENTABLE MINIMAL SURFACES

M. ELISA G. G. DE OLIVEIRA

ABSTRACT. The classical Henneberg's minimal surface (1875, [3, 4, 11]) was the unique nonorientable example known until 1981, when Meeks [6] exhibited the first example of a nonorientable, regular, complete, minimal surface of finite total curvature -6π .

In this paper, we study the nonorientable, regular, complete minimal surfaces of finite total curvature and give some examples of punctured projective planes regularly and minimally immersed in $\mathbf{R}^3$ and $\mathbf{R}^n$.

1. Nonorientable minimal surfaces in $\mathbf{R}^n$. We consider surfaces S in $\mathbf{R}^n$ defined by maps $X: M \rightarrow \mathbf{R}^n$, where M is a two-dimensional manifold.

For the study of a nonorientable, connected surface S we take the double surface $\tilde{S}$ given by $\tilde{X}: \tilde{M} \rightarrow \mathbf{R}^n$, where $\Pi: \tilde{M} \rightarrow M$ is the oriented two-sheeted covering of M , and $\tilde{X} = X \circ \Pi$. We have an involution $\tilde{I}: \tilde{M} \rightarrow \tilde{M}$ without fixed points and a conformal structure on $\tilde{M}$ such that $\tilde{I}$ is antiholomorphic.

We have a representation theorem for the nonorientable minimal surfaces:

THEOREM 1.1. *Let S be a nonorientable regular connected minimal surface in $\mathbf{R}^n$. The double surface $\tilde{S}$ is a minimal surface in $\mathbf{R}^n$ defined by $\tilde{X}: \tilde{M} \rightarrow \mathbf{R}^n$,*

$$(1.1) \quad \tilde{X}(p) = \operatorname{Re} \int_{p_0}^p \tilde{\phi}(\zeta) d\zeta, \quad p_0, p \in \tilde{M},$$

with $\alpha = \tilde{\phi}(\zeta)d\zeta$ such that $\tilde{I}^*\alpha = \bar{\alpha}$ (that is, $\tilde{I}^*\alpha_k = \bar{\alpha}_k$, $1 \leq k \leq n$).

Conversely, if $\tilde{S}$ is a regular orientable connected minimal surface in $\mathbf{R}^n$ given by (1.1) and if there is an antiholomorphic involution $\tilde{I}: \tilde{M} \rightarrow \tilde{M}$ without fixed points such that $\tilde{I}^*\alpha = \bar{\alpha}$, then $\tilde{S}$ is the double surface of a regular nonorientable minimal surface in $\mathbf{R}^n$.

The consequences of the condition $\tilde{I}^*\alpha = \bar{\alpha}$ with respect to the various forms of representation for minimal surfaces in $\mathbf{R}^n$ are

COROLLARY 1.2 (MEEKS [6]). *Let f and g be the functions of Weierstrass's representation of an orientable, regular, connected minimal surface $\tilde{S}$ in $\mathbf{R}^3$, that is*

$$\tilde{X}(p) = \operatorname{Re} \int_{p_0}^p \frac{f(\zeta)}{2} (1 - g^2(\zeta), i(1 + g^2(\zeta)), 2g(\zeta)) d\zeta, \quad p_0, p \in \tilde{M}.$$

The surface $\tilde{S}$ is the double surface of a nonorientable, regular minimal surface in $\mathbf{R}^3$ if and only if

$$(1.2) \quad \begin{cases} \text{(i)} & g(\tilde{I}(p)) = -1/\overline{g(p)}, & p \in \tilde{M}, \text{ and} \\ \text{(ii)} & \tilde{I}^*(\omega) = -g^2\omega, & \omega = f(z) dz, \end{cases}$$

for some antiholomorphic involution $\tilde{I}: \tilde{M} \rightarrow \tilde{M}$ without fixed points.

Received by the editors October 31, 1985.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 53A10; Secondary 30A68.

©1986 American Mathematical Society
0002-9939/86 \$1.00 + \$.25 per page

COROLLARY 1.3. *Let $\tilde{S}$ be an orientable, regular, connected minimal surface in $\mathbf{R}^4$ given by $\tilde{X}: \tilde{M} \rightarrow \mathbf{R}^4$,*

$$\tilde{X}(p) = \operatorname{Re} \int_{p_0}^p (1 + g_1 g_2, i(1 - g_1 g_2), g_1 - g_2, -i(g_1 + g_2)) \omega, \quad p_0, p \in \tilde{M}.$$

Then, $\tilde{S}$ is the double surface of a nonorientable minimal surface in $\mathbf{R}^4$ if and only if, for some antiholomorphic involution $\tilde{I}$ on $\tilde{M}$ without fixed points,

$$(1.3) \quad \begin{cases} \text{(i)} & g_k(\tilde{I}(p)) = -1/\overline{g_k(p)}, & k = 1, 2, \text{ and} \\ \text{(ii)} & \tilde{I}^* \omega = \overline{g_1 g_2 \omega}, & \omega = f(z) dz, \end{cases}$$

If $C(\tilde{S})$ is the total curvature of $\tilde{S}$, we can define the total curvature of S by

$$(1.4) \quad C(S) = C(\tilde{S})/2.$$

In what follows, the double surface $\tilde{S}$ will always be connected, complete and of finite total curvature $C(\tilde{S}) = 2\pi\tilde{m}$. From Chern and Osserman's theorem [1] we have $\tilde{M}$ conformally equivalent to a compact Riemann surface of genus $\tilde{\gamma}$ punctured at $\tilde{r} = 2r$ points $\{p_1, p_2, \dots, p_r, q_1, \dots, q_r\}$, with $\tilde{I}(p_j) = q_j$, $\tilde{I}(p_j)$ being the extension of $\tilde{I}$ to p_j , $1 \leq j \leq r$. The function $\tilde{\phi}$ of (1.1) has poles of order m_j at p_j and q_j , and

$$(1.5) \quad \tilde{m} - 2 \sum_{j=1}^r m_j = 2\tilde{\gamma} - 2.$$

The Chern and Osserman inequality for nonorientable minimal surfaces is

$$(1.6) \quad C(S) \leq 2\pi(\chi(M) - r),$$

with $\chi(M)$ the Euler characteristic of M .

A nonorientable version of Gackstatter's theorem [2] gives an estimate for the dimension of a nonorientable, complete, minimal surface S with total curvature $C(S) = -2\pi m$, r ends and genus γ :

$$(1.7) \quad \operatorname{Dim}(S) \leq 2m - 2\gamma - r + 3.$$

We have

PROPOSITION 1.4. *The total curvature of a nonorientable, regular, complete minimal surface is at most -4π .*

PROOF. From (1.7) and the fact that $\operatorname{Dim} S \geq 3$, $2m \geq 4$.

2. Genus one nonorientable complete minimal surfaces in $\mathbf{R}^n$. A characterization of the complete minimal surfaces of finite total curvature and genus

one can be set up combining Theorem 1.1 and Hoffman and Osserman's theorem [5]:

THEOREM 2.1. *Let $\tilde{M}$ be the complex plane minus $2r-1$ points $\{0, z_1, \dots, z_{r-1}, -1/\bar{z}_1, \dots, -1/\bar{z}_{r-1}\}$. Let the complex vector $\phi(\zeta)$ be of the form*

$$\phi(\zeta) = F(\zeta)(p_1(\zeta), \dots, p_n(\zeta)),$$

with

$$F(\zeta) = \left(\zeta^{\nu_0} \prod_{k=1}^{r-1} (\zeta - z_k)^{\nu_k} \left(\zeta + \frac{1}{\bar{z}_k} \right)^{\nu_k} \right)^{-1},$$

$p_j(\zeta)$ satisfying

- (i) $p_j(\zeta)$ is a polynomial,
- (ii) the maximum degree of p_j is $2m$,
- (iii) the p_j 's have no common factor,
- (iv) $\sum_{j=1}^n p_j^2 \equiv 0$,
- (v) $(-1)^{m+1} \bar{\zeta}^{2m} p_j(-1/\bar{\zeta}) = \overline{p_j(\zeta)}$, $1 \leq j \leq n$.

Suppose that the ν_k 's satisfy $\sum_{k=0}^{r-1} \nu_k = m+1$, $\nu_k \geq 2$, $0 \leq k \leq r-1$ and $\text{Re} \int_{\gamma} \phi = 0$, for any simple closed curve γ in $\tilde{M}$. Then $\tilde{X}(p) = \text{Re} \int_{p_0}^p \phi(\zeta) d\zeta$ defines a double minimal surface associated to a nonorientable regular complete minimal surface in $\mathbf{R}^n$ of genus one, r ends and total curvature $-2\pi m$.

Conversely, every genus one nonorientable minimal surface in $\mathbf{R}^n$ admits a representation as mentioned above.

PROOF. The double surface associated to a genus one nonorientable minimal surface is conformally equivalent to the complex plane $\hat{\mathbf{C}}$ minus $2r$ points with the involution $\tilde{I}: \mathbf{C} \rightarrow \mathbf{C}$ given by $\tilde{I}(z) = -1/\bar{z}$, that is,

$$\tilde{M} = \mathbf{C} - \{0, z_1, \dots, z_{r-1}, -1/\bar{z}_1, \dots, -1/\bar{z}_{r-1}\}.$$

From Hoffman and Osserman's result, $\phi = F \cdot (p_1, \dots, p_n)$, p_j polynomials satisfying (i)–(iv).

The map $\tilde{X}: \tilde{M} \rightarrow \mathbf{R}^n$ defines a double surface if and only if $\tilde{I}^*(\phi_j(\zeta) d\zeta) = \overline{\phi_j(\zeta) d\zeta}$, $1 \leq j \leq n$; from this fact and (1.5), we have $\sum_{k=0}^{r-1} \nu_k = m+1$ and (v). $\square$

We study initially surfaces whose total curvature is the upper bound given by Proposition 1.4.

THEOREM 2.2. *Let S be a nonorientable, regular, complete minimal surface of finite total curvature -4π . Then S is a minimal immersion of the projective plane minus one point and lies fully in $\mathbf{R}^4$; any two such minimal surfaces are similar.*

PROOF. From (1.5), and (1.6) and (1.7), using $\chi(M) = 2 - \gamma - r$, we have $\gamma = 1$, $r = 1$ and $m_1 = 3$. The double surface has genus zero and two ends; thus, by Theorem 2.1, it can be given by $\tilde{X}: \mathbf{C} - \{0\} \rightarrow \mathbf{R}^n$,

$$\tilde{X}_j(p) = \text{Re} \int_{p_0}^p \phi_j(\zeta) d\zeta, \quad p_0, p \in \tilde{M} = \mathbf{C} - \{0\}, \quad \phi_j(\zeta) = p_j(\zeta)/\zeta^3,$$

p_j satisfying conditions (i)–(v).

Setting $p_j(\zeta) = a_j\zeta^4 + b_j\zeta^3 + c_j\zeta^2 + d_j\zeta + e_j$, we can verify that $a_j = -e_j$, $b_j = \bar{d}_j$, $c_j = 0$ and that $A_1 = (\operatorname{Re} a_1, \dots, \operatorname{Re} a_n)$, $A_2 = (\operatorname{Im} a_1, \dots, \operatorname{Im} a_n)$, $B_1 = (\operatorname{Re} b_1, \dots, \operatorname{Re} b_n)$ and $B_2 = (\operatorname{Im} b_1, \dots, \operatorname{Im} b_n)$ are real orthogonal vectors of the same length.

Thus, $A = (1, i, 0, 0)$ and $B = (0, 0, 1, i)$ give a solution in $\mathbf{R}^4$; two solutions $\tilde{X}$ and $\tilde{Y}$ are such that $\tilde{Y} = \lambda M \tilde{X} + X_0$, with $\lambda \in \mathbf{R}$ and M a real orthogonal matrix.

Immediately, we have Meeks' theorem:

COROLLARY 2.3. *The total curvature of a nonorientable, regular, complete minimal surface in $\mathbf{R}^3$ is at most -6π .*

The total curvature of an orientable minimal surface in $\mathbf{R}^3$ is of the form $C(\tilde{S}) = -4\pi \cdot \text{degree}(g)$, with g the meromorphic function of the Weierstrass representation. Thus, for nonorientable minimal surfaces,

$$(2.1) \quad C(S) = -2\pi \cdot \text{degree}(g),$$

with $g: \tilde{M} \rightarrow \mathbf{C}$ the function of the Weierstrass representation of the double surface $\tilde{S}$, which is a rational function when it has finite degree and the genus of $\tilde{M}$ is zero.

The next result is very important for the construction of examples:

PROPOSITION 2.4. *Let g be a rational complex function satisfying $g(-1/\bar{z}) = -1/\overline{g(z)}$. Then, the degree of g is odd and g is of the form*

$$g(z) = cz^\alpha \frac{\prod_{j=1}^m (z - a_j)}{\prod_{j=1}^m (z + 1/\bar{a}_j)}, \quad \text{with } |c| \prod_{j=1}^m |a_j| = 1.$$

PROOF. Let g be a rational function, and let

$$g(z) = cz^\alpha \frac{\prod_{j=1}^m (z - a_j)}{\prod_{k=1}^n (z - b_k)}, \quad a_j \neq b_k \text{ for any } j, k, a_j \neq 0, b_k \neq 0,$$

where the factors appear with multiplicities. From the condition

$$g(-1/\bar{z}) = -1/\overline{g(z)},$$

with some calculations, we complete the proof. $\square$

PROPOSITION 2.5. *The total curvature of a nonorientable regular complete minimal surface in $\mathbf{R}^3$ of genus one is of the form $C(S) = -2\pi m$, m odd, $m \geq 3$.*

PROOF. This follows immediately from Proposition 2.4. $\square$

Meeks [6] has proved that there exists a unique nonorientable regular complete minimal surface in $\mathbf{R}^3$ of total curvature -6π . We extend Meeks' surface in the sense of

THEOREM 2.6. *There exists a nonorientable regular complete minimal surface in $\mathbf{R}^3$ of genus one, one end and total curvature $-2\pi m$, for any m odd, $m \geq 3$.*

PROOF. With straightforward calculations, one can show that

$$f(z) = i(z+1)^2/z^{m+1} \quad \text{and} \quad g(z) = z^{m-1}(z-1)/(z+1)$$

are the functions of the Weierstrass representation of a double minimal surface $\tilde{S}$; that is, the map

$$\tilde{X}(p) = \operatorname{Re} \int_{p_0}^p \frac{f}{2} (1 - g^2, i(1 + g^2), 2g) d\zeta, \quad p_0, p \in \mathbf{C} - \{0\},$$

satisfies (i)–(v) of Theorem 2.1. The surfaces obtained are regular and complete. $\square$

REMARKS. 1. If $n = 3$ the surface of Theorem 2.6 is exactly Meeks' surface.

2. The surface is an infinite Moebius band with $(m-1)/2$ twists and the image of $|z| = 1$ is a circle centered at $(0, -2/(m-1), 0)$ with radius $2/(m-1)$ covered $(m-1)$ times.

3. The normals to the surface along $|z| = 1$ have the third component $N_3|_{|z|=1} = \operatorname{Re}(z)|_{|z|=1}$.

The Chern-Osserman inequality for nonorientable minimal surfaces of total curvature -6π gives us the relation $\gamma + 2r \leq 5$; hence, if $\gamma = 1$, we may have $r = 1$ or $r = 2$. Meeks has proved that the case $r = 2$, $C = -6\pi$ cannot occur. Thus, we have

PROPOSITION 2.7. *The total curvature of a nonorientable regular complete minimal surface in $\mathbf{R}^3$ of genus one and two ends is at most -10π .*

For $C = -10\pi$, we have

THEOREM 2.8. *There exists a nonorientable regular complete minimal surface in $\mathbf{R}^3$ of genus one, two ends and total curvature -10π .*

PROOF. The double surface associated to such a surface will be an orientable surface of genus zero, four ends with total curvature -20π and, by Theorem 2.1, it may be given by

$$\tilde{X}: \mathbf{C} - \{0, 1, -1\} \rightarrow \mathbf{R}^3, \quad \tilde{I}: \mathbf{C} \rightarrow \mathbf{C}, \quad \tilde{I}(z) = -1/\bar{z}.$$

Let $f(z) = a(\bar{b}^2 z^2 - 1)^2/z^2(z-1)^4(z+1)^4$ and $g(z) = \alpha \bar{b}^2 z^3(z^2 - b^2)/(\bar{b}^2 z^2 - 1)$; f and g are the functions of the Weierstrass representation of a double minimal surface if and only if

$$(2.2) \quad |\alpha| |b|^2 = 1 \quad \text{and} \quad a = -\bar{a}\bar{\alpha}b^4 \quad \text{or} \quad \alpha\bar{\alpha}\bar{b}^2 = -\bar{a}\bar{\alpha}b^2.$$

The completeness and regularity can easily be verified.

To choose α , a and b such that

$$\operatorname{Re} \int_{\gamma} \frac{f(z)}{2} (1 - g^2(z), i(1 + g^2(z)), 2g(z)) dz = \bar{0}$$

for any closed curve γ in $\mathbf{C} - \{0, 1, -1\}$, we observe that, by Cauchy's integral formula, for simple closed curves γ_0 and γ_1 around $z = 0$ and $z = 1$, respectively,

$$\int_{\gamma_0} \frac{p_j(\zeta) d\zeta}{\zeta^2(\zeta-1)^4(\zeta+1)^4} = 2\pi i p'_j(0),$$

and

$$\int_{\gamma_1} \frac{p_j(\zeta) d\zeta}{\zeta^2(\zeta-1)^4(\zeta+1)^4} = \frac{2\pi i}{3!} \cdot \frac{1}{16} [p_j'''(1) - 12p_j''(1) + 57p_j'(1) - 105p_j(1)].$$

Calling $R_j = p_j'''(1) - 12p_j''(1) + 57p_j'(1) - 105p_j(1)$, $j = 1, 2, 3$, we have

$$\begin{aligned} R_1 &= -105(a - \bar{a}) + 30(\bar{a}b^2 - \bar{a}b^2) + 3(\bar{a}b^4 - \bar{a}b^4), \\ R_2 &= i[-105(a + \bar{a}) + 30(\bar{a}b^2 + \bar{a}b^2) + 3(\bar{a}b^4 + \bar{a}b^4)], \\ R_3 &= 0. \end{aligned}$$

Thus, $\operatorname{Re}[2\pi i R_j] = 0$, $j = 1, 2$ if $-105a + 30a\bar{b}^2 + 3a\bar{b}^4 = 0$.

For each b , $\alpha = -1/\bar{b}^2$, $a = i$ satisfy (2.2) and

$$\tilde{X}(p) = \operatorname{Re} \int_{p_0}^p \frac{f}{2} (1 - g^2, i(1 + g^2), 2g) d\zeta$$

gives the double surface. $\square$

We investigate the problem of the regular minimal surfaces in $\mathbf{R}^3$ of finite total curvature, genus one and three ends, and, we have

THEOREM 2.9. *There exists a nonorientable regular complete minimal surface in $\mathbf{R}^3$ of genus one, three ends and total curvature -14π .*

PROOF. We construct the double surface $\tilde{S}$ defined by $\tilde{X}: \mathbf{C} - \{0, 1, -1, i, -i\} \rightarrow \mathbf{R}^3$ with the functions f and g of the Weierstrass representation (1.2) given by $g(z) = a\bar{b}^4 z^3(z^4 - b^4)/(\bar{b}^4 z^4 - 1)$ and $f(z) = \alpha(\bar{b}^4 z^4 - 1)^2/z^2(z^2 - 1)^3(z^2 + 1)^3$. The functions f and g satisfy (1.2) if and only if

$$(2.3) \quad |a||b|^4 = 1, \quad \alpha = \bar{\alpha}\bar{a}^2\bar{b}^8.$$

With some calculations, we can verify that the real parts of the integrals

$$\int_{\gamma} \frac{p_j(z) dz}{z^2(z^4 - 1)^3},$$

where γ is a simple closed curve, are null for any j , $j = 1, 2, 3$, provided b satisfies the following condition: $3b^8 + 10b^4 - 45 = 0$. Thus, $b^4 \in \mathbf{R}$.

Choosing $a = 1/b^4$ and $\alpha = 1$ the conditions (2.3) are verified. The completeness and regularity can be easily checked. $\square$

In $\mathbf{R}^4$, exploring the properties of the functions g_1, g_2 of the representation (1.3), we have

THEOREM 2.10. *For any integer m , $m \geq 2$, there exists a nonorientable regular complete minimal surface S in $\mathbf{R}^4$ of genus one, one end and total curvature $-2\pi m$.*

PROOF. The double surface $\tilde{S}$ associated to S has total curvature $-4\pi m = -2\pi\tilde{m}$, $\tilde{m} = 2m$, genus zero and two ends. We set the functions f, g_1, g_2 in the representation (1.3) by $f(z) = z^{m-1}$, $g_1(z) = \varepsilon_1/z^{k_1}$, $g_2(z) = \varepsilon_2/z^{k_2}$, with $\tilde{m} = 2m = k_1 + k_2$, k_1, k_2 odd, $k_1, k_2 \geq 1$, and $\tilde{M} = \mathbf{C} - \{0\}$.

The functions f, g_1 and g_2 satisfy (i) and (ii) of Corollary 1.3, if and only if

$$(2.4) \quad |\varepsilon_1| = |\varepsilon_2| = 1, \quad \varepsilon_1\varepsilon_2 = (-1)^{m-1}.$$

To verify that $\operatorname{Re} \int_{\gamma} \phi(\zeta) d\zeta = \vec{0}$, for any closed curve in $\mathbf{C} - \{0\}$, we observe that

$$\begin{aligned} \phi_1(\zeta) &= f(1 + g_1g_2) = z^{m-1} + \varepsilon_1\varepsilon_2/z^{m+1}, \\ \phi_2(\zeta) &= if(1 - g_1g_2) = i(z^{m-1} - \varepsilon_1\varepsilon_2/z^{m+1}), \\ \phi_3(\zeta) &= f(g_1 - g_2) = (\varepsilon_1z^{k_2} - \varepsilon_2z^{k_1})/z^{m+1}, \\ \phi_4(\zeta) &= -if(g_1 + g_2) = -i(\varepsilon_1z^{k_2} + \varepsilon_2z^{k_1})/z^{m+1}, \end{aligned}$$

and rewriting $k_1 = m - n$, $k_2 = m + n$, $n \geq 1$, we have $\int_\gamma \phi_j = 0$, $j = 1, \dots, 4$. Thus, the double surface $\tilde{S}$ is defined by

$$\tilde{X}(p) = \operatorname{Re} \int_{p_0}^p \phi(\zeta) d\zeta, \quad p_0, p \in \mathbf{C} - \{0\}. \quad \square$$

For $\gamma = 1$, $r = 1$, the dimension of S satisfies $\operatorname{Dim}(S) \leq 2m$. We will show that this upper bound is sharp.

THEOREM 2.11. *For any integer m , $m \geq 2$, there exists a nonorientable regular complete minimal surface S in $\mathbf{R}^{2m}$ of genus one, one end and total curvature $-2\pi m$ that lies fully in $\mathbf{R}^{2m}$.*

PROOF. The double surface $\tilde{S}$ associated to S , by Theorem 2.1, is given by $\tilde{X}: \mathbf{C} - \{0\} \rightarrow \mathbf{R}^n$, $\tilde{X}(p) = \operatorname{Re} \int_{p_0}^p \phi(\zeta) d\zeta$, with $\phi_j(\zeta) = p_j(\zeta)/\zeta^{m+1}$, $p_j(\zeta)$ satisfying (i)–(v).

Particularly, condition (v) is equivalent to

$$(2.5) \quad (-1)^{m+1} \bar{\zeta}^{2m} p_j(-1/\bar{\zeta}) = \overline{p_j(\zeta)}, \quad 1 \leq j \leq n.$$

Taking $p_j(\zeta) = a_0^j + a_1^j \zeta + \dots + a_{2m-1}^j \zeta^{2m-1} + a_{2m}^j \zeta^{2m}$ in (2.5) we obtain

$$(-1)^{m+1} (a_0^j \bar{\zeta}^{2m} - a_1^j \bar{\zeta}^{2m-1} + \dots + a_{2m}^j) = (\bar{a}_{2m}^j \bar{\zeta}^{2m} + \bar{a}_{2m-1}^j \bar{\zeta}^{2m-1} + \dots + \bar{a}_0^j).$$

Then, there are two possibilities:

- (a) $(m+1)$ even: $a_0^j = \bar{a}_{2m}^j$, $-a_1^j = \bar{a}_{2m-1}^j$, $\dots$, $a_{m-1}^j = \bar{a}_{m+1}^j$, $-a_m^j = \bar{a}_0^j$,
- (b) $(m+1)$ odd: $a_0^j = -\bar{a}_{2m}^j$, $a_1^j = \bar{a}_{2m-1}^j$, $\dots$, $a_{m-1}^j = \bar{a}_{m+1}^j$, $a_m^j = -\bar{a}_0^j$.

The integrals $\int_\gamma \phi_j(\zeta) d\zeta$, γ a closed curve in $\mathbf{C} - \{0\}$, should have null real part; therefore, $\operatorname{Re}\{2\pi i \cdot a_m^j\} = 0$, $1 \leq j \leq n$, that is $\operatorname{Im} a_m^j = 0$, $1 \leq j \leq n$. Comparing with the last condition in (a) or (b), we have $a_m^j = 0$, $1 \leq j \leq n$.

To determine solutions in $\mathbf{R}^n$, we should find complex vectors $A_k = (a_k^1, \dots, a_k^n)$, $0 \leq k \leq m-1$, such that the corresponding polynomials satisfy $\sum_{j=1}^n p_j^2(\zeta) \equiv 0$.

It is easy to verify that $A_0 = (1, i, 0, \dots, 0)$, $A_1 = (0, 0, 1, i, 0, \dots, 0)$, $\dots$, $A_{m-1} = (0, \dots, 0, 1, i)$ give a solution in $\mathbf{C}^{2m}$ if m is even and $A_0 = (1, i, 0, \dots, 0)$, $A_2 = (0, 0, 1, i, 0, \dots, 0)$, $\dots$, $A_{m-2} = (0, 0, \dots, 0, \sqrt{2}, i\sqrt{2}, 0, 0)$, $A_{m-1} = (0, 0, \dots, 0, 1, i)$ give a solution if m is odd. $\square$

With these results, we have partial answers for the problem of existence of nonorientable, regular, minimal surfaces in $\mathbf{R}^3$ and $\mathbf{R}^n$ posed by Nitsche [7, 1965]. We do not know, for instance, an example of a genus two nonorientable minimal surface in $\mathbf{R}^3$, that is, a minimal Klein bottle in $\mathbf{R}^3$.

REFERENCES

1. S. S. Chern and R. Osserman, *Complete minimal surfaces in Euclidean n -space*, J. Analyse Math. **19** (1967), 15–30.
2. F. Gackstatter, *Über die Dimension einer minimalfläche und zur Ungleichung von St. Cohn-Vossen*, Arch. Rational Mech. Anal. **61** (1976), 141–152.
3. L. Henneberg, *Über solche minimalfläche, welche eine vorgeschriebene ebene curve zur geodätischen line haben*, Doctoral Dissertation, Eidgenössisches Polytechnikum, Zurich, 1875.
4. —, *Über diejenige minimalfläche, welche die Neil'sche Parabel zur ebenen geodätischen line hat*, Vierteljahr Natuforsch, Ges. Zürich **21** (1876), 66–70.

5. D. A. Hoffman and R. Osserman, *The geometry of the generalized Gauss map*, Mem. Amer. Math. Soc. No. 28 (1980).
6. W. H. Meeks III, *The classification of complete minimal surfaces with total curvature greater than -8π* , Duke Math. J. **48** (1981), 523–535.
7. J. C. C. Nitsche, *On new results in the theory of minimal surfaces*, Bull. Amer. Math. Soc. **71** (1965), 195–270.
8. M. E. G. Oliveira, *Superfícies mínimas não-orientáveis no R^n* , Tese de Doutorado-IME-USP, 1984.
9. R. Osserman, *A survey of minimal surfaces*, Van Nostrand Reinhold, New York, 1969.
10. ———, *Global properties of minimal surfaces in E^3 and E^n* , Ann. of Math. (2) **80** (1964), 340–364.
11. M. Spivak, *A comprehensive introduction to differential geometry*, Vol. 3, Publish or Perish, Boston, Mass., 1975.

INSTITUTO DE MATEMÁTICA E ESTATÍSTICA, UNIVERSIDADE DE SÃO PAULO, CAIXA
POSTAL 20.570 (AGÊNCIA IGUATEMI), SÃO PAULO, BRAZIL