GENERAL POSITION THEOREMS FOR GENERALIZED MANIFOLDS

J. L. BRYANT

ABSTRACT. It is an open question as to whether a generalized n -manifold, $n \geq 5$, that satisfies the disjoint disks property is a topological manifold. In this paper it is shown that any such space X satisfies general position properties for maps of polyhedra into X .

The program to establish a topological characterization of euclidean space has been stalled, at least temporarily, by the discovery that Quinn's obstruction to resolving a generalized manifold may not be categorically zero as first asserted in [6]. (See [7].) Thus Cannon's conjecture [3] that a generalized n -manifold X , $n \geq 5$, is a topological manifold if and only if X satisfies the disjoint disks property remains open. Quinn's results [6, 7], however, do produce an obstruction $\sigma(X)$ that vanishes if and only if X is a topological manifold, although it is not known at this time whether the obstruction can be nonzero.

In this paper we show (Theorem 3) that any generalized n -manifold X , $n \geq 5$, having the disjoint disks property also satisfies general position properties for maps of arbitrary polyhedra into X . Thus, in a dimension theory sense X behaves like a topological manifold. Theorems of this type have also been obtained by John Walsh [9].

DEFINITIONS. A *generalized n -manifold*, n -gm, is a euclidean neighborhood retract X that is also a *homology n -manifold*; that is, $H_*(X, X - x) \cong H_*(\mathbf{R}^n, \mathbf{R}^n - 0)$ for all $x \in X$. (All homology is understood to have integer coefficients.) The space X is said to satisfy the *disjoint k -disks property*, DD^kP (or, simply, the disjoint disks property, DDP, when $k = 2$) if for every pair of maps $f_1, f_2: D^k \rightarrow X$ of the unit k -disk D^k of $\mathbf{R}^k$ into X and $\varepsilon > 0$, there are maps $f'_1, f'_2: D^k \rightarrow X$ such that $d(f'_i, f_i) < \varepsilon$ and $f'_1(D^k) \cap f'_2(D^k) = \emptyset$. If $f, g: A \rightarrow X$ and $\varepsilon > 0$, then $f \simeq_\varepsilon g$ means f is ε -homotopic to g . A subset A of X is said to be 1-LCC in X (1-locally coconnected in X) if for every $a \in A$ and neighborhood U of a in X there is a neighborhood V of a in X such that the inclusion induced homomorphism $\pi_1(V - A) \rightarrow \pi_1(U - A)$ is zero. A map (or embedding) $f: A \rightarrow X$ is said to be 1-LCC provided $f(A)$ is 1-LCC in X .

Main results. In [3] Cannon shows that if X is an n -gm, $n \geq 5$, having the DDP, then an arbitrary map of a 2-disk (hence, a 2-dimensional polyhedron) into

Received by the editors September 24, 1985.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 57P05, 57N75; Secondary 57Q65.

The author's research was partially funded by ONR grant #N00014-84-K-0761.

©1986 American Mathematical Society
0002-9939/86 \$1.00 + \$.25 per page

X can be approximated by 1-LCC embeddings. Our first result is

THEOREM 1. *If X is an n -gm, $n \geq 5$, satisfying the DDP, P is a k -dimensional polyhedron, $2k + 1 \leq n$, and $f: P \rightarrow X$, then f can be approximated by 1-LCC embeddings.*

We show that an n -gm X with the DDP is never “ghastly” [4].

THEOREM 2. *Suppose X is an n -gm, $n \geq 5$, having the DDP, P is a k -dimensional polyhedron, and $f: P \rightarrow X$ is a map. Then f can be approximated by a map $f': P \rightarrow X$ such that*

- (1) $\dim f'(P) \leq k$, and
- (2) $f'(P)$ is 1-LCC if $k \leq n - 3$.

Finally, we obtain the main general position property.

THEOREM 3. *Suppose X is an n -gm, $n \geq 5$, with the DDP, P and Q are polyhedra of dimensions p and q , respectively, and $f: P \rightarrow X$ and $g: Q \rightarrow X$ are maps. Then f and g can be approximated by maps f' and g' such that*

- (1) $\dim[f'(P) \cap g'(Q)] \leq p + q - n$, and
- (2) $f'(P) \cap g'(Q)$ is 1-LCC in X if $p + q - n \leq n - 3$.

Proofs of the main results. We start with a well-known fact (proved, for example, in [2, 8]).

LEMMA 0. *A locally compact, finite-dimensional, separable metric space X admits a 1-LCC embedding in some euclidean space.*

LEMMA 1. *Suppose X is an n -dimensional, 1-LCC closed subset of $\mathbf{R}^m$, $m - n \geq 3$. Then there are F_σ subsets $X_0 \subset X_1 \subset \cdots \subset X_n = X$ such that*

- (1) $\dim X_i \leq i$, and
- (2) $\dim(X_i - X_j) \leq i - j - 1$.

PROOF. Let $L_k \subset \mathbf{R}^m$ denote the Menger space:

$$L_k = \{x \in \mathbf{R}^m \mid \text{at least } m - k \text{ coordinates of } x \text{ are rational}\}.$$

Then $\dim L_k = k$, L_k is a countable union of k -dimensional hyperplanes, and $\dim(\mathbf{R}^m - L_k) = m - k - 1$ [5]. Also $L_0 \subset L_1 \subset \cdots \subset L_m = \mathbf{R}^m$ and $\dim(L_r - L_s) = r - s - 1$. Since X is 1-LCC in $\mathbf{R}^m$, we can assume that $L_k \cap X = \emptyset$ for $k \leq m - n - 1$ [2, 8].

Set $X_i = X \cap L_{m-n+i}$. Then

$$\begin{aligned} \dim X_i &= \dim(X \cap L_{m-n+i}) \\ &= \dim(X \cap (L_{m-n+i} - L_{m-n-1})) \\ &\leq (m - n + i) - (m - n - 1) - 1 = i, \end{aligned}$$

and

$$\begin{aligned} \dim(X_i - X_j) &= \dim(X \cap (L_{m-n+i} - L_{m-n+j})) \\ &\leq \dim(L_{m-n+i} - L_{m-n+j}) = i - j - 1. \end{aligned}$$

Throughout the rest of this paper X is an n -gm satisfying the DDP, 1-LCC embedded in $\mathbf{R}^m$ for some $m \geq n + 3$.

LEMMA 2. *There are F_σ subsets $X_0 \subset X_1 \subset \cdots \subset X_n = X$ such that*

- (1) $\dim X_i \leq i$,
- (2) $\dim(X_i - X_j) \leq i - j - 1$, and
- (3) X_i is 1-LCC in X for $i \leq n - 3$.

PROOF. Let $\{f_i\}_{i=1}^\infty$ be a countable dense subset of $\text{Map}(D^2, X)$, the space of maps (with the compact-open topology), consisting of 1-LCC embeddings [3]. In the proof of Lemma 1, require that $L_k \cap f_i(D^2) = \emptyset$ for $k \leq m - 3$ and all i (using the fact that the 1-LCC property is transitive). Then $X_i \cap f_j(D^2) = \emptyset$ for $i \leq n - 3$. Thus X_i is 1-LCC in X for $i \leq n - 3$.

LEMMA 3. *Let $X_0 \subset X_1 \subset \cdots \subset X_n = X$ be F_σ subsets of X satisfying the conclusion of Lemma 2. Then $\pi_k(X, X - X_i) = 0$ for $k < n - i$.*

PROOF. By local duality in X [1, 10], $H_k(X, X - X_i) = 0$ for $k < n - i$. If $i = n - 1$ or $n - 2$, then $\pi_k(X, X - X_i) = H_k(X, X - X_i) = 0$. If $i \leq n - 3$, then X_i is 1-LCC in X so that $\pi_2(X, X - X_i) = 0$. Thus by the Hurewicz isomorphism theorem, $\pi_k(X, X - X_i) = 0$ for $i \leq n - 3$ and $k < n - i$. (We may assume that X , and hence $X - X_i$, $i \leq n - 3$, are simply connected by passing to the universal cover of X .)

LEMMA 4. *Suppose $f: D^k \rightarrow X$ is a map and $\varepsilon > 0$. Then $f \simeq_\varepsilon f'$ such that $f'(D^k) \cap X_{n-k-1} = \emptyset$ and f' is 1-LCC, if $k \leq n - 3$.*

PROOF. To get $f'(D^k) \cap X_{n-k-1} = \emptyset$, apply Lemma 3 locally, using induction on the skeleta of a fine triangulation of D^k . Similarly, if D is a 1-LCC 2-cell in X , then for a relative neighborhood U of $\text{Int } D$, $(\text{rel } \text{Bd } D)$, $\pi_2(U, U - \text{Int } D) = 0$ and $H_i(U, U - \text{Int } D) = 0$ for $i \leq n - 3$. Thus for $r \leq n - 3$ any map $g: (D^r, \text{Bd } D^r) \rightarrow (U, U - \text{Int } D)$ is homotopic rel $\text{Bd } D^r$ to a map $g': D^r \rightarrow U - \text{Int } D$. Applying this argument inductively on the skeleta of a fine triangulation of D^k , we can get $f \simeq_\varepsilon f'$ such that $f'(D^k) \cap D = \emptyset$ for any 1-LCC 2-cell D in X . Thus we may simultaneously obtain $f \simeq_\varepsilon f'$ so that f' is 1-LCC and $f'(D^k) \cap X_{n-k-1} = \emptyset$.

COROLLARY. *If $f: D^k \rightarrow X$ and $\varepsilon > 0$, then $f \simeq_\varepsilon f'$ where f' is 1-LCC, if $k \leq n - 3$, and $\dim f'(D^k) \leq k$.*

The proof of Theorem 2 follows immediately.

LEMMA 5. *X satisfies DD^kP for $k < n/2$.*

PROOF. Given $f, g: D^k \rightarrow X$, apply the Corollary to get $f \simeq_\varepsilon f'$ where f' is 1-LCC and $\dim f'(D^k) \leq k$. Then for any open set U of X , $\pi_i(U, U - f'(D^k)) = 0$ for $i \leq n - k - 1$. If $k < n/2$, then $k \leq n - k - 1$ and, hence, by a now familiar argument, for any $\varepsilon > 0$, $g \simeq_\varepsilon g'$ where $g'(D^k) \cap f'(D^k) = \emptyset$.

The proof of Theorem 1 now follows easily as in [3].

PROOF OF THEOREM 3. It is clearly sufficient to prove the theorem when P is a p -simplex and Q is a q -simplex. Suppose $f: P \rightarrow X$ and $g: Q \rightarrow X$ are maps. Let $X_0 \subset X_1 \subset \cdots \subset X_n = X$ be F_σ subsets of X obtained from Lemma 2. Then f can be approximated by $f': P \rightarrow X$ such that $f'(P) \cap X_i \subset X_i - X_{n-p-1}$, by applying Lemma 3.

Observe that $f'(P) \cap X_{2n-p-q-1}$ is a σ -compact subset of $X_{2n-p-q-1} - X_{n-p-1}$, which has dimension $n - q - 1$. We wish to approximate g by $g': Q \rightarrow X$ so that $g'(Q) \cap f'(P) \cap X_{2n-p-q-1} = \emptyset$, for then it would follow that $g'(Q) \cap f'(P) \subset X - X_{2n-p-q-1}$, which has dimension $p + q - n$, as desired.

Recall that $L_r = \{x \in \mathbf{R}^m \mid \text{at most } m - r \text{ coordinates of } x \text{ are rational}\} = \bigcup_{j=1}^{\infty} E_j$, where $E_1 \subset E_2 \subset \dots$ and each E_j is a finite union of r -dimensional hyperplanes [5]. Thus each $X_i = \bigcup_{j=1}^{\infty} Y_{ij}$ where Y_{ij} is a closed subset of X of dimension $\leq i$ and Y_{ij} is 1-LCC in X if $i \leq n - 3$. For $i = 2n - p - q - 1$, $Y_{ij} \cap f'(P)$ is a compact set of dimension $\leq n - q - 1$.

Assume first that $i = 2n - p - q - 1 \leq n - 3$. Then $H_k(X, X - (Y_{ij} \cap f'(P))) = 0$ for $0 \leq k \leq q$ and so $\pi_k(X, X - (Y_{ij} \cap f'(P))) = 0$ for $0 \leq k \leq q$; hence, we can choose $g': Q \rightarrow X$ approximating g so that $g'(Q) \cap (Y_{ij} \cap f'(P)) = \emptyset$ for all j . That is, $g'(Q) \cap f'(P) \cap X_{2n-p-q-1} = \emptyset$.

Suppose next that $i = 2n - p - q - 1 = n - 2$, $p \leq q$ (without loss of generality), and $n \geq 6$. Then $n - p \geq 3$ and we can assume further that $f'(P)$ and, hence, $f'(P) \cap Y_{ij}$ are 1-LCC in X . From this we see that $\pi_k(X, X - (f'(P) \cap Y_{ij})) = 0$ for $0 \leq k \leq q$ and so we can find g' such that $g'(Q) \cap f'(P) \cap Y_{ij} = \emptyset$.

Suppose $2n - p - q - 1 = n - 1$, $n \geq 5$, and $p \leq q$. Then again $n - p \geq 3$ and the above argument works.

Finally, if $2n - p - q - 1 = n - 2$ and $n = 5$, then the only case in which the above argument does not work is $p = q = 3$.

Suppose then that $n = 5$ and $p = q = 3$. Let $\{f_i\}_{i=1}^{\infty}$ be a countable dense subset of $\text{Map}(D^2, X)$ consisting of 1-LCC embeddings. Given $f: P \rightarrow X$ and $g: Q \rightarrow X$, approximate f by f' such that $\dim(f'(P) \cap f_i(D^2)) \leq 0$ for all $i = 1, 2, \dots$ and $\dim f'(P) \leq 3$. Then $f'(P)$ and $f'(P) \cap f_i(D^2)$, $i = 1, 2, \dots$, are 1-LCC in $\mathbf{R}^m$ since X is. Thus for any $\varepsilon > 0$ it is possible to find an ε -homeomorphism $h: \mathbf{R}^m \rightarrow \mathbf{R}^m$ such that $h(f'(P) \cap f_i(D^2)) \cap L_{m-2} = \emptyset$, $i = 1, 2, \dots$, $h(X) \cap L_k = \emptyset$, for $k \leq m - 4$, and $h(f'(P)) \cap L_{m-2}$ is a 1-dimensional σ -compact set that is 1-LCC in X .

Now replace $X_i = X \cap L_{m-5+i}$ by $Y_i = X \cap h^{-1}(L_{m-5+i})$. Then $f'(P) \cap Y_3$ is a 1-dimensional σ -compact set that is 1-LCC in X . Hence, g can be approximated by $g': Q \rightarrow X$ such that $g'(Q) \cap f'(P) \cap Y_3 = \emptyset$. This means that $g'(Q) \cap f'(P) \subset X - Y_3$, which has dimension 1.

REFERENCES

1. G. E. Bredon, *Sheaf theory*, McGraw-Hill, New York, 1967.
2. J. Bryant, *On embeddings of compacta in euclidean space*, Proc. Amer. Math. Soc. **23** (1969), 46-51.
3. J. W. Cannon, *Shrinking cell-like decompositions of manifolds. Codimension three*, Ann. of Math. (2) **110** (1979), 83-112.
4. R. J. Daverman and J. Walsh, *A ghastly generalized n -manifold*, Illinois J. Math. **25** (1981), 555-576.
5. W. Hurewicz and H. Wallman, *Dimension theory*, Princeton Univ. Press, Princeton, N. J., 1948.
6. F. Quinn, *Resolutions of homology manifolds and the topological characterization of manifolds*, Invent. Math. **72** (1983), 267-284.
7. —, *An obstruction to the resolutions of homology manifolds*, preprint.
8. M. A. Stan'ko, *The embedding of compacta in euclidean space*, Mat. Sb. **83** (125) (1970), 234-255 (=Math. USSR-Sb. **12** (1970), 234-255).
9. J. Walsh, *General position properties of generalized manifolds: a primer*, Proc. Idaho State Topology Conf., 1985.
10. R. L. Wilder, *Topology of manifolds*, Amer. Math. Soc. Colloq. Publ., vol. 32, Amer. Math. Soc., Providence, R.I., 1949.