ON A THEOREM OF HUNEKE CONCERNING MULTIPLICITIES

LIAM O'CARROLL

ABSTRACT. We consider a theorem by Huneke on multiplicities, and show that the extension surmised by Huneke is a stronger form of the Syzygy Conjecture.

Craig Huneke has proved the following result.

THEOREM 1 [9]. *Let $(A, \underline{m})$ be a complete local ring containing a field, let n be a positive integer and let $e(A)$ be the multiplicity of A . Suppose that A satisfies Serre's condition S_n and that $e(A) \leq n$; if $n = 1$ suppose further that A is equidimensional. Then A is Cohen-Macaulay.*

All rings considered in this note are commutative, Noetherian and have an identity element, and all modules are finitely generated. For a discussion of the theory of (complete) local rings and of multiplicities, see [11, 12, 13, 14]. A finitely generated module M over a Noetherian ring A is said to satisfy S_n if

$$\text{depth } M_P \geq \inf(n, \text{height } P) \quad \text{for all } P \in \text{Spec } A.$$

As remarked on [11, p. 125], the condition S_1 is equivalent to A having no embedded associated primes. We use the term 'equidimensional' in the sense of Grothendieck's E.G.A., i.e. a local ring is said to be *equidimensional* when all its minimal primes have the same co-height. Recall from [7, (5.10.9)] that a catenary (and so a complete) local ring which satisfies S_2 is equidimensional. A recent handy reference for much of the material in this note is the monograph [5], where a special case of Huneke's result is discussed on pp. 71–72 and on p. 75. As the latter shows, and as has been remarked by Huneke himself, in the case $n = 1$ it is essential in Theorem 1 to add the hypothesis of equidimensionality.

From his theorem Huneke deduces results by Nagata [12, (40.6)] and by Ikeda, in the case where the rings involved contain a field. (As regards the former result, see the interesting aside on [6, p. 41].) Since these are known to hold without this restriction, Huneke surmises that the following is probably true:

- (1) *Theorem 1 is valid without the assumption that A contain a field.*

Now recall the Syzygy Conjecture [4, p. 143] for regular local rings, which runs as follows:

- (2) *if R is a regular local ring, and if K is the j th syzygy of an R -module with rank $K < j$, then K is a free R -module.*

Received by the editors November 4, 1985.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 13H15.

Key words and phrases. Multiplicity, local ring, syzygy.

©1987 American Mathematical Society
 0002-9939/87 \$1.00 + \$.25 per page

For a discussion of the links between j th syzygies and the condition S_j , and for a proof that (2) holds when R contains a field, see [5, Chapter 3] and especially [loc. cit., Theorem 3.8 and Corollary 3.16]. It is also known that (2) holds when $\dim R \leq 4$ [4, p. 147].

Our final conjecture is the following:

- (3) *Let $(R, \underline{m})$ be a regular local ring, let $(B, \underline{m}')$ be a local ring which dominates [12, p. 14] R and suppose that B is a finitely generated R -module. Let n be a positive integer. If B satisfies S_n and if the multiplicity $e_R(B) \leq n$ (where, if $n = 1$, B is also supposed to be equidimensional), then B is Cohen-Macaulay.*

REMARKS. Note that B is catenary, being the homomorphic image of a polynomial ring over the regular ring R ; note also that if $m \geq n$ then $S_m \Rightarrow S_n$. Hence, by [7, (5.10.9)], whatever the value of n in (3), B is equidimensional. Since B is an integral extension of R , any two primes in B having the same restriction to R must have the same co-height in B (see [10, Theorem 47]), and hence the same height in B . It follows from [7, (5.7.11)] that B satisfies S_n as a B -module if and only if it satisfies S_n as an R -module.

THEOREM 2. *Conjectures (2) and (3) are equivalent, and are implied by Conjecture (1).*

PROOF. First suppose that (3) holds, and let R and K be as in the statement of (2). Consider the ring $B = R \ltimes K$ formed by Nagata's process of idealization [12, p. 2]. Then B is a local ring which dominates R and B is a finitely generated R -module. Since $B = R \oplus K$ as an R -module, $\text{rank}_R B \leq j$. But as remarked in [9] (and as can easily be deduced from [12, §23])

$$e_R(B) = \text{rank}_R B,$$

where $e_R(B)$ is the multiplicity of B as an R -module. By Auslander and Bridger's theorem [1, (4.25)] (see [5, Theorem 3.8]), K satisfies S_j . Now as R -modules, $B_P = R_P \oplus K_P$ for all $P \in \text{Spec } R$. By the definition of depth via the Ext-functor say (e.g. see [5, p. 3]), for all $P \in \text{Spec } R$,

$$\text{depth}_R B_P = \min\{\text{depth}_R R_P, \text{depth}_R K_P\} = \min\{j, \text{height } P\},$$

since R is regular. Hence B satisfies S_j as an R -module, and so as a B -module, and $e_R(B) \leq j$. By (3) therefore, B is Cohen-Macaulay. Thus B is a free module, by [12, (25.16)], and therefore K is also free, by [2, p. 147, Ex. 4] (or [5, p. 25, Ex. 10]).

Conversely, suppose that (2) holds, and let R and B be as in (3). Now B satisfies S_n as an R -module, and by the remarks above, $\text{rank}_R B \leq n$. Let $C = B/R$ and consider the exact sequence

$$0 \rightarrow R \rightarrow B \rightarrow C \rightarrow 0.$$

Then $\text{rank}_R C \leq n - 1$.

Fix $P \in \text{Spec } R$. Then we have the exact sequence

$$0 \rightarrow R_P \rightarrow B_P \rightarrow C_P \rightarrow 0.$$

First suppose that $\text{height } P \leq n$. Then B_P is a Cohen-Macaulay R_P -module, and B_P is a semi-local integral over-ring of R_P . It easily follows that B_P is itself a Cohen-Macaulay ring. Hence as before, by [12, (25.16)] and [2, p. 147, Ex. 4], C_P is a free R_P -module and so, by the definition of depth via the Ext-functor (say) and by the regularity of R , $\text{depth}_R C_P$ equals $\text{height } P$.

On the other hand, suppose that $\text{height } P > n$. It follows from the 'Depth Lemma' [5, p. 13], since R is regular, that $\text{depth}_R C_P \geq \text{depth}_R B_P$ or $\text{depth}_R C_P = \text{height } P - 1$. Whatever the case therefore, $\text{depth}_R C_P \geq n$ since B satisfies S_n as an R -module. Hence C also satisfies S_n as an R -module, and $\text{rank}_R C \leq n - 1$. By (2) therefore, C is free. Hence B is free, and so is Cohen-Macaulay by [12, (25.16)] again.

Finally consider Conjecture (1). Assume that it is valid. Then Conjecture (3) holds in the case where R is a complete regular local ring, for it easily follows that in this case B is also complete in the $\underline{m}'$ -adic (as well as the $\underline{m}$ -adic) topology; by [14, Corollary 1, p. 299] and [12, (23.6)] $e_R(B) \geq e(B)$. By Theorem 2 (and its proof), Conjecture (2) then holds in the case where R is complete. But by the faithful flatness of the completion functor, this is no restriction on Conjecture (2), which therefore holds as stated.

COROLLARY 1. *Conjecture (3) holds when $\dim R (= \dim B) \leq 4$.*

PROOF. See the remarks after the statement of Conjectures (2) and (3).

If one wishes to dispense with the restriction that R be a regular ring in Conjecture (3) and merely assume that B is a complete local ring, then it easily follows (as on [3, p. 59], say) that B is an equidimensional catenary ring which is a finitely generated torsion-free module over a complete local ring S which is either regular or a hypersurface (i.e. a regular local ring modulo a principal ideal); moreover, S contains a system of parameters for B . It is not clear (to me) what else can usefully be said about this more general situation. The example given on pp. 27–28 of [8] is of interest in this regard, as it shows that the analogue of [12, (25.16)] for (complete) hypersurfaces (rather than regular local rings) is no longer valid.

NOTE ADDED IN PROOF. I would like to thank S. Ikeda for pointing out an error in the original version of this paper, and for very valuable discussions.

REFERENCES

1. M. Auslander and M. Bridger, *Stable module theory*, Mem. Amer. Math. Soc. **94** (1969).
2. N. Bourbaki, *Commutative algebra*, Addison-Wesley, Reading, Mass., 1972.
3. W. Bruns, *On the number of elements independent with respect to an ideal*, J. London Math. Soc. (2) **22** (1980), 57–62.
4. W. Bruns, E. G. Evans and P. Griffith, *Syzygies, ideals of height two, and vector bundles*, J. Algebra **67** (1980), 143–162.
5. E. G. Evans and P. Griffith, *Syzygies*, London Math. Soc. Lecture Notes Ser., vol. 106, Cambridge Univ. Press, Cambridge and New York, 1985.
6. W. Fulton, *Introduction to intersection theory in algebraic geometry*, CBMS Regional Conf. Ser. in Math., no. 54, Amer. Math. Soc., Providence, R. I., 1984.
7. A. Grothendieck, *Éléments de géométrie algébrique*, Inst. Hautes Études Sci. Publ. Math. **24** (1965).
8. M. Hochster, *Contracted ideals from integral extensions of regular rings*, Nagoya Math. J. **51** (1973), 25–43.
9. C. Huneke, *A remark concerning multiplicities*, Proc. Amer. Math. Soc. **85** (1982), 331–332.

10. I. Kaplansky, *Commutative rings*, Allyn and Bacon, Boston, Mass., 1970.
11. H. Matsumura, *Commutative algebra*, Benjamin, New York, 1970.
12. M. Nagata, *Local rings*, Wiley-Interscience, New York, 1962.
13. D. G. Northcott, *Lessons on rings, modules and multiplicities*, Cambridge Univ. Press, London, 1968.
14. O. Zariski and P. Samuel, *Commutative algebra*. Vol. II, Springer-Verlag, New York, 1960.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF EDINBURGH, JAMES CLARK MAXWELL BUILDING, KING'S BUILDINGS, MAYFIELD ROAD, EDINBURGH EH9 3JZ, SCOTLAND