AN UPPER BOUND FOR THE PERMANENT OF A 3-DIMENSIONAL $(0, 1)$ -MATRIX

STEPHEN J. DOW AND PETER M. GIBSON

ABSTRACT. Let $A = (a_{ijk})$ be a 3-dimensional matrix of order n . The permanent of A is defined by

$$\text{per } A = \sum_{\sigma, \tau \in S_n} \prod_{i=1}^n a_{i\sigma(i)\tau(i)},$$

where S_n is the symmetric group on $\{1, 2, \dots, n\}$. Suppose that A is a $(0, 1)$ -matrix and that $r_i = \sum_{j,k=1}^n a_{ijk}$ for $i = 1, 2, \dots, n$. In this paper it is shown that $\text{per } A \leq \prod_{i=1}^n r_i!^{1/r_i}$. A similar bound is then obtained for a second function, the 2-permanent of a 3-dimensional matrix, that is another analogue of the permanent of an ordinary (2-dimensional) matrix.

1. Introduction. It was conjectured by Minc [3] and proved by Brègman [1] that if A is a $(0, 1)$ -matrix of order n with row sums $r_1, r_2, \dots, r_n$, then

$$\text{per } A \leq \prod_{i=1}^n r_i!^{1/r_i},$$

where by definition $0!^{1/0} = 0$. In this paper it is shown that an analogous result holds for 3-dimensional matrices. The *permanent* of a 3-dimensional matrix $A = (a_{ijk})$ of order n is defined by

$$(1) \quad \text{per } A = \sum_{\sigma, \tau \in S_n} \prod_{i=1}^n a_{i\sigma(i)\tau(i)}$$

where S_n is the symmetric group on $\{1, 2, \dots, n\}$. We now state our principal result.

THEOREM 1. *Let $A = (a_{ijk})$ be a 3-dimensional $(0, 1)$ -matrix of order n , and let $r_i = \sum_{j,k=1}^n a_{ijk}$ for $i = 1, 2, \dots, n$. Then*

$$\text{per } A \leq \prod_{i=1}^n r_i!^{1/r_i}.$$

The proof of Theorem 1, presented in §2, is an adaptation of Schrijver's proof [4] of the Minc-Brègman bound, with the extension to 3-dimensional matrices requiring a number of additional lemmas which may be of some independent interest.

In equation (1) and Theorem 1 the planes of A play the role of the rows and columns of an ordinary (2-dimensional) matrix, where a *plane* of a 3-dimensional

Received by the editors December 2, 1985.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 15A15; Secondary 05A20.

©1987 American Mathematical Society
 0002-9939/87 \$1.00 + \$.25 per page

matrix $A = (a_{ijk})$ is a submatrix obtained by fixing one of the indices i, j, k and letting the other two vary. We could also have taken the analogue of rows and columns to be the *lines* of A , obtained by fixing two of these indices. In §3 we give a second definition of the permanent of a 3-dimensional matrix, called the 2-permanent, based on this observation. Using a result from [2] which relates these two types of permanents we obtain an upper bound for the 2-permanent of a 3-dimensional $(0, 1)$ -matrix similar to the one given in Theorem 1, with plane sums replaced by line sums.

2. Proof of the principal result. We write N , Z , and R^+ to denote, respectively, the set of natural numbers, the set of integers, and the set of positive real numbers. Also, for any real number x , we use the notation $\lfloor x \rfloor = \max\{k \in Z : k \leq x\}$ and $\lceil x \rceil = \min\{k \in Z : x \leq k\}$. The first lemma appears in [4].

LEMMA 1. *If $t = \sum_{i=1}^n t_i$, $t_i \in R^+$, then $t^t \leq n^t \prod_{i=1}^n t_i^{t_i}$.*

LEMMA 2. *For all $n \in N$, $n!^{1/n}(n+2)!^{1/(n+2)} \leq (n+1)!^{2/(n+1)}$.*

PROOF. It is well known that if $x_1, x_2, \dots, x_k \in R^+$, then $\prod_{i=1}^k x_i \leq \bar{x}^k$ where $\bar{x} = (\sum_{i=1}^k x_i)/k$. Let $x_i = i$ for $i = 1, 2, \dots, n$, $x_i = i - n$ for $i = n+1, n+2, \dots, 2n$, and $x_i = n+2$ for $i = 2n+1, 2n+2, \dots, n^2+3n$. Then

$$\prod_{i=1}^{n^2+3n} x_i = n!^2(n+2)^{n^2+n}$$

and

$$\sum_{i=1}^{n^2+3n} x_i = (n^2+3n)(n+1).$$

Hence the inequality above becomes $n!^2(n+2)^{n^2+n} \leq (n+1)^{n^2+3n}$, which is equivalent to the desired inequality.

LEMMA 3. *Suppose that a function $f: N \rightarrow R^+$ has the property that*

$$f(x)f(x+2) \leq (f(x+1))^2$$

for all $x \in N$, and that $x_1, x_2, \dots, x_k \in N$. Let $\sum_{i=1}^k x_i = k\bar{x} = k_1\lfloor\bar{x}\rfloor + k_2\lceil\bar{x}\rceil$, where $k_1, k_2 \in Z$, $k_1, k_2 \geq 0$, and $k_1 + k_2 = k$. Then

$$\prod_{i=1}^k f(x_i) \leq f(\lfloor\bar{x}\rfloor)^{k_1} f(\lceil\bar{x}\rceil)^{k_2}.$$

PROOF. It follows from induction on n that

$$(2) \quad f(x)f(x+n) \leq f(x+1)f(x+n-1)$$

for all $x, n \in N$. To see this note that the inductive step is

$$f(x)f(x+n+1) = \frac{f(x)f(x+2)f(x+1)f(x+1+n)}{f(x+1)f(x+2)} \leq f(x+1)f(x+n).$$

By (2), if there exist p and q such that $x_p - x_q > 1$ and we let $x'_p = x_p - 1$, $x'_q = x_q + 1$, and $x'_i = x_i$ for $i \neq p, q$, then $\prod_{i=1}^k f(x_i) \leq \prod_{i=1}^k f(x'_i)$. By repeating this step the upper bound eventually becomes $f(\lfloor\bar{x}\rfloor)^{k_1} f(\lceil\bar{x}\rceil)^{k_2}$.

LEMMA 4. Let $f(x) = x!^{1/x}$ for $x \in N$. Suppose that $n, p, r \in N$, $n \geq 3$, and $p \leq (r-1)/(n-1)$. Then

$$f(r-p-1)^{r-pn+p-1} f(r-p)^{(p+1)n-p-r} \leq (r-1)! r^{n/r-1}.$$

PROOF. Let $k \in N$, $k \leq p$. Then $r-k-1 > 0$. Therefore, by Lemma 2, $f(r-k-1)f(r-k+1) \leq f(r-k)^2$. Since $k \leq (r-1)/(n-1)$, we can raise both sides of this inequality to the power $r-kn+k-1$ to obtain

$$f(r-k-1)^{r-kn+k-1} f(r-k+1)^{r-kn+k-1} \leq f(r-k)^{2(r-kn+k-1)}.$$

This inequality is equivalent to

$$(3) \quad \begin{aligned} f(r-k-1)^{r-kn+k-1} f(r-k)^{(k+1)n-k-r} \\ \leq f(r-k)^{r-(k-1)n+k-2} f(r-k+1)^{kn-k+1-r}. \end{aligned}$$

The desired inequality can now be obtained by successively applying (3) with $k = p, p-1, p-2, \dots, 1$.

LEMMA 5. Let $x_1, x_2, \dots, x_{n-1}$, $r \in N$ with $n > 2$. If $\sum_{i=1}^{n-1} x_i \leq (n-2)r+1$, then

$$\prod_{i=1}^{n-1} x_i!^{1/x_i} \leq (r-1)! r^{n/r-1}.$$

PROOF. Since $x!^{1/x}$ is increasing in x we may assume that $\sum_{i=1}^{n-1} x_i = (n-2)r+1$. Hence

$$\bar{x} = \frac{(n-2)r+1}{n-1} = r - \frac{r-1}{n-1}.$$

Let $p = \lfloor (r-1)/(n-1) \rfloor$. Then $\lfloor \bar{x} \rfloor = \lceil \bar{x} \rceil = r-p$ if x is an integer and $\lfloor \bar{x} \rfloor = r-p-1$, $\lceil \bar{x} \rceil = r-p$ otherwise. Since

$$(r-pn+p-1)(r-p-1) + ((p+1)n-p-r)(r-p) = (n-2)r+1$$

we can apply Lemma 3 with $f(x) = x!^{1/x}$, $k = n-1$, $k_1 = r-pn+p-1$, and $k_2 = (p+1)n-p-r$. (When $\lfloor \bar{x} \rfloor = \lceil \bar{x} \rceil = r-p$ we have $k_1 = 0$.) Therefore Lemmas 3 and 4 imply that

$$\prod_{i=1}^{n-1} x_i!^{1/x_i} \leq f(r-p-1)^{r-pn+p-1} f(r-p)^{(p+1)n-p-r} \leq (r-1)! r^{n/r-1}.$$

LEMMA 6. If $B = (b_{jk})$ is a (0, 1)-matrix of order n with $\sum_{j,k=1}^n b_{jk} = r$, then

$$\sum_{i=1}^{n-1} \sum_{\substack{j,k \\ j \neq i \\ k \neq i}} b_{jk} \leq (n-2)r+1.$$

PROOF.

$$\begin{aligned} \sum_{i=1}^{n-1} \sum_{\substack{j,k \\ j \neq i \\ k \neq i}} b_{jk} &= \sum_{i=1}^{n-1} \left(r - \sum_{j=1}^n b_{ji} - \sum_{k=1}^n b_{ik} + b_{ii} \right) \\ &= (n-3)r + \sum_{j=1}^n b_{jn} + \sum_{k=1}^n b_{nk} + \sum_{i=1}^{n-1} b_{ii} \\ &\leq (n-3)r + r + 1 = (n-2)r+1. \end{aligned}$$

We now show how Schrijver's proof [4] of the Minc-Brègman bound can be altered to prove Theorem 1.

PROOF OF THEOREM 1. The proof is by induction on n . Clearly the theorem holds for $n = 1$. Let $n > 1$. It suffices to assume that $r_i \in N$ for $i = 1, 2, \dots, n$. Let A_{ijk} denote the 3-dimensional submatrix of A of order $n - 1$ obtained by deleting the three planes through cell (i, j, k) . Let

$$S = \left\{ (\sigma, \tau) : \prod_{i=1}^n a_{i\sigma(i)\tau(i)} = 1 \right\}.$$

Then $|S| = \text{per } A$. Apply Lemma 1 with $t = \text{per } A = \sum_{j,k; a_{ijk}=1} \text{per } A_{ijk}$ for $i = 1, 2, \dots, n$, to obtain

$$\begin{aligned} (\text{per } A)^{n \text{ per } A} &= \prod_{i=1}^n (\text{per } A)^{\text{per } A} \\ &\leq \prod_{i=1}^n r_i^{\text{per } A} \prod_{\substack{j,k \\ a_{ijk}=1}} (\text{per } A_{ijk})^{\text{per } A_{ijk}}. \end{aligned}$$

For each i we have

$$\begin{aligned} \prod_{\substack{j,k \\ a_{ijk}=1}} (\text{per } A_{ijk})^{\text{per } A_{ijk}} &= \prod_{\substack{j,k \\ a_{ijk}=1}} (\text{per } A_{ijk})^{|\{(\sigma, \tau) \in S : \sigma(i)=j, \tau(i)=k\}|} \\ &= \prod_{(\sigma, \tau) \in S} \text{per } A_{i\sigma(i)\tau(i)}. \end{aligned}$$

By the induction hypothesis,

$$\text{per } A_{i\sigma(i)\tau(i)} \leq \prod_{\substack{p \\ p \neq i}} r(p, i)^{1/r(p, i)}$$

where

$$r(p, i) = \sum_{\substack{j,k \\ j \neq \sigma(i) \\ k \neq \tau(i)}} a_{pj k}.$$

Combining the inequalities above we have

$$\begin{aligned} (\text{per } A)^{n \text{ per } A} &\leq \prod_{i=1}^n r_i^{\text{per } A} \prod_{(\sigma, \tau) \in S} \prod_{\substack{p \\ p \neq i}} r(p, i)^{1/r(p, i)} \\ &= \prod_{(\sigma, \tau) \in S} \left(\prod_{i=1}^n r_i \right) \prod_{i=1}^n \prod_{\substack{p \\ p \neq i}} r(p, i)^{1/r(p, i)} \\ &= \prod_{(\sigma, \tau) \in S} \left(\prod_{i=1}^n r_i \right) \prod_{p=1}^n \prod_{\substack{i \\ i \neq p}} r(p, i)^{1/r(p, i)}. \end{aligned}$$

Fix σ, τ , and p and let $b_{jk} = a_{p\sigma(j)\tau(k)}$. Then $\sum_{j,k} b_{jk} = r_p$ and for each i ,

$$\sum_{\substack{j,k \\ j \neq i, k \neq i}} = r(p, i).$$

Hence, by Lemma 6,

$$\sum_{\substack{i \\ i \neq p}} r(p, i) \leq (n-2)r_p + 1.$$

Therefore, by Lemma 5, if $n > 2$, then

$$\prod_{\substack{i \\ i \neq p}} r(p, i)!^{1/r(p, i)} \leq (r_p - 1)! r_p!^{n/r_p - 1}.$$

Moreover, it is easy to see that this inequality also holds for $n = 2$. Therefore,

$$\begin{aligned} (\text{per } A)^{n \text{ per } A} &\leq \prod_{(\sigma, \tau) \in S} \left(\prod_{i=1}^n r_i \right) \prod_{p=1}^n (r_p - 1)! r_p!^{n/r_p - 1} \\ &= \prod_{(\sigma, \tau) \in S} \prod_{p=1}^n r_p!^{n/r_p} = \left(\prod_{i=1}^n r_i!^{1/r_i} \right)^{n \text{ per } A}, \end{aligned}$$

and the theorem follows.

For some values of the r_i it is easy to construct matrices A such that equality holds in Theorem 1. Let $A_m = (a_{ijk})$ denote the 3-dimensional $(0, 1)$ -matrix of order m with $a_{ijj} = 1$ for $i, j = 1, 2, \dots, m$, and $a_{ijk} = 0$ otherwise. Then $r_i = \sum_{j,k=1}^m a_{ijk} = m$ for $i = 1, 2, \dots, m$, and $\text{per } A_m = m!$. Now let $n_1, n_2, \dots, n_t$ be a partition of n , and let A be the direct sum of $A_{n_1}, A_{n_2}, \dots, A_{n_t}$. It follows that A is a 3-dimensional $(0, 1)$ -matrix of order n such that equality holds in Theorem 1.

3. The 2-permanent. Theorem 1 can be used to obtain another extension of the Minc-Brègman inequality to 3-dimensional matrices. Let $A = (a_{ijk})$ be a 3-dimensional matrix of order n . The planes of A are the submatrices obtained by fixing one of i, j, k , and the lines of A are the submatrices obtained by fixing two of i, j, k . Observe that the permanent (or 1-permanent) of A is equal to the summation of all products of n entries of A , no two entries from the same plane. Similarly, the 2-permanent of A is defined to be the summation of all products of n^2 entries of A , no two entries from the same line [2]. We have the following upper bound for the 2-permanent of a $(0, 1)$ -matrix.

THEOREM 2. *Let $A = (a_{ijk})$ be a 3-dimensional $(0, 1)$ -matrix of order n , and let $r_{ij} = \sum_{k=1}^n a_{ijk}$ for $i, j = 1, 2, \dots, n$. Then*

$$2\text{-per } A \leq \prod_{i,j=1}^n r_{ij}!^{1/r_{ij}}.$$

PROOF. As shown by the authors [2], there exists a 3-dimensional $(0, 1)$ -matrix $B = (b_{ijk})$ of order n^2 such that $\text{per } B = 2\text{-per } A$. Moreover, B can be chosen so that if $s_i = \sum_{j,k=1}^{n^2} b_{ijk}$ for $i = 1, 2, \dots, n^2$, then $\{s_i : i = 1, 2, \dots, n^2\} = \{r_{ij} : i, j = 1, 2, \dots, n\}$. Therefore, if Theorem 1 is applied to B the desired upper bound on 2-per A is obtained.

REFERENCES

1. L. M. Brègman, *Certain properties of nonnegative matrices and their permanents*, Dokl. Akad. Nauk SSSR **211** (1973), 27–30; English transl., Soviet Math. Dokl. **14** (1973), 945–949.
2. S. J. Dow and P. M. Gibson, *Permanents of d -dimensional matrices* (submitted for publication).
3. H. Minc, *Upper bounds for permanents of $(0, 1)$ -matrices*, Bull. Amer. Math. Soc. **69** (1963), 789–791.
4. A. Schrijver, *A short proof of Minc's conjecture*, J. Combin. Theory Ser. A **25** (1978), 80–83.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ALABAMA, HUNTSVILLE, ALABAMA
35899