A PROBABILITY DISTRIBUTION ASSOCIATED WITH THE HURWITZ ZETA FUNCTION

R. L. DUNCAN

ABSTRACT. A new discrete probability distribution is defined in terms of the Hurwitz zeta function and its relationship to the Poisson distribution is demonstrated.

If we define the Hurwitz zeta function by

$$\zeta(s, a) = \sum_{n=0}^{\infty} \frac{1}{(n+a)^s}$$

for $a > 0$ and $s > 1$, then it is easy to show that

$$f(x; a) = a\zeta(x+1, a+1), \quad x = 1, 2, 3, \dots,$$

is a probability density function (p.d.f) with factorial moment generating function

$$G(t) = a \sum_{n=1}^{\infty} \left(\frac{1}{n+a-t} - \frac{1}{n+a} \right)$$

for $|t| < 1+a$ and factorial moments

$$G^{(m)}(1) = am!\zeta(m+1, a).$$

It will now be shown how $f(x; a)$ arises in a natural way from the Poisson distribution. It is known [1, 13.12] that

$$\zeta(s, a) = \frac{1}{\Gamma(s)} \int_0^{\infty} \frac{y^{s-1} e^{-ay}}{1 - e^{-y}} dy.$$

If $g(x; y)$ and $h(y; a)$ are the p.d.f.s for the Poisson distribution with mean y and the exponential distribution with mean $1/a$, then

$$f(x; a) = \int_0^{\infty} g^*(x; y) h(y; a) dy,$$

where

$$g^*(x; y) = \frac{g(x; y)}{1 - g(0; y)}, \quad x = 1, 2, 3, \dots,$$

is the p.d.f. for a truncated Poisson distribution. Thus, by the same argument used in [2, 5.13B], $f(x; a)$ is the p.d.f. for a compound truncated Poisson distribution where the parameter in the truncated Poisson distribution has an exponential distribution with mean $1/a$.

Received by the editors November 22, 1985 and, in revised form, February 17, 1986 and May 21, 1986.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 62E10.

©1987 American Mathematical Society
0002-9939/87 \$1.00 + \$.25 per page

Now let a and b be positive and replace $h(y; a)$ by the p.d.f. for the gamma distribution defined by

$$h(y; a, b) = \frac{a^b}{\Gamma(b)} y^{b-1} e^{-ay}, \quad 0 < y < \infty.$$

Then it follows as before that the compound truncated Poisson distribution, where the truncated Poisson distribution is the same as above but whose parameter now has a gamma distribution, will have the p.d.f.

$$f(x; a, b) = a^b b(b+1) \cdots (b+x-1) \zeta(x+b, a+1)/x!, \quad x = 1, 2, 3, \dots,$$

and it can be shown that the factorial moments for this distribution are

$$G^{(m)}(1) = a^b b(b+1) \cdots (b+m-1) \zeta(m+b, a).$$

The corresponding problem for the compound Poisson distribution is considered in [2, 5.13B].

If $b/a = c = \text{constant}$ and x is a positive integer, then

$$\zeta(x+b, a+1) = \left(\frac{c}{b+c}\right)^{x+b} \sum_{n=0}^{\infty} \left(1 + \frac{cn}{b+c}\right)^{-x-b}$$

and

$$f(x; a, b) = \frac{c^x}{x!} \left(\frac{b}{b+c}\right)^b \left[\frac{b(b+1) \cdots (b+x-1)}{(b+c)^x} \right] \sum_{n=0}^{\infty} \left(1 + \frac{cn}{b+c}\right)^{-x-b}.$$

Hence

$$\lim_{b \rightarrow \infty} f(x; a, b) = \frac{c^x e^{-c}}{x!} \sum_{n=0}^{\infty} e^{-cn} = \frac{c^x e^{-c}}{x!(1-e^{-c})}.$$

Thus the truncated Poisson distribution with parameter b/a provides a simple approximation for $f(x; a, b)$ when a and b are large.

For $t < \log(1+a)$, the moment generating function is given by

$$\begin{aligned} M_X(t) &= a^b \sum_{n=1}^{\infty} (n+a)^{-b} \sum_{x=1}^{\infty} \frac{b(b+1) \cdots (b+x-1)}{x!} \left[\frac{e^t}{n+a} \right]^x \\ &= a^b \sum_{n=1}^{\infty} (n+a)^{-b} \left[\left(1 - \frac{e^t}{n+a}\right)^{-b} - 1 \right] \\ &= a^b \sum_{n=1}^{\infty} [(n+a-e^t)^{-b} - (n+a)^{-b}]. \end{aligned}$$

If $Y = (aX - b)/c$, where $c = \sqrt{b(a+1)}$, then

$$M_Y(t) = \sum_{n=1}^{\infty} \left\{ \left[\frac{a}{e^{t/c}(n+a-e^{at/c})} \right]^b - \left[\frac{a}{e^{t/c}(n+a)} \right]^b \right\}$$

and

$$\begin{aligned} \lim_{b \rightarrow \infty} M_Y(t) &= \lim_{b \rightarrow \infty} \left[a/e^{t/c}(1+a-e^{at/c}) \right]^b \\ &= \lim_{b \rightarrow \infty} [(1+t/c+t^2/2c^2+\cdots)(1-t/c-at^2/2c^2-\cdots)]^{-b} \\ &= \lim_{b \rightarrow \infty} [1-t^2/2b+o(1/b)]^{-b} = e^{t^2/2}. \end{aligned}$$

Hence Y has an approximate standard normal distribution when b is large.

If $Z = 2aX$, then

$$M_Z(t) = \sum_{n=1}^{\infty} \left[\left(\frac{a}{n+a-e^{2at}} \right)^b - \left(\frac{a}{n+a} \right)^b \right]$$

and

$$\lim_{a \rightarrow 0} M_Z(t) = \lim_{a \rightarrow 0} \left(\frac{a}{1+a-e^{2at}} \right)^b = (1-2t)^{-b}.$$

Thus Z has an approximate chi-square distribution with $2b$ degrees of freedom when a is small.

When $x = 1$, we have

$$\begin{aligned} f(1; a, b) &= a^b b \zeta(b+1, a+1) = a^b b \sum_{n=1}^{\infty} (n+a)^{-b-1} \\ &\geq a^b b \int_1^{\infty} (u+a)^{-b-1} du = \left(\frac{a}{a+1} \right)^b. \end{aligned}$$

Hence X has a distribution which is approximately degenerate at one when a is large or b is small.

Finally, it has been observed by the referee that there are two other discrete distributions [3, 10.3] that are related to the Riemann zeta function $\zeta(s, 1)$.

REFERENCES

1. E. T. Whittaker and G. N. Watson, *Modern analysis*, Cambridge Univ. Press, New York, 1927.
2. M. Fisz, *Probability theory and mathematical statistics*, Wiley, New York, 1963.
3. N. L. Johnson and S. Kotz, *Discrete distributions*, Houghton Mifflin, Boston, Mass., 1969.

GRADUATE STUDIES AND CONTINUING EDUCATION, PENNSYLVANIA STATE UNIVERSITY, KING OF PRUSSIA, PENNSYLVANIA 19406