ON THE p -ADIC HEIGHTS OF SOME ABELIAN VARIETIES

HIDEO IMAI

ABSTRACT. For an abelian variety defined over an algebraic number field, different definitions of p -adic heights have been given by several authors. In this note, we shall prove that the p -adic height defined by A. Néron and that by P. Schneider coincide.

1. In this section we shall recall briefly the construction of p -adic heights. Let K be a finite extension field of the p -adic rational number field $\mathbf{Q}_p$, where p is a prime number. Let $\mathcal{O}$ be the integer ring of K , $\mathfrak{p}$ the maximal ideal of $\mathcal{O}$, and let $k = \mathcal{O}/\mathfrak{p}$ be the residue field of $\mathcal{O}$. Let A be an abelian variety of dimension d defined over K , $\delta = \text{id}_A$ the identity map of A . Let 0 be the identity point of A and let $t_1, \dots, t_d$ be a $\mathfrak{p}$ -admissible system of local coordinates of A at 0 in the sense of A. Néron; namely, $t_1, \dots, t_d$ are K -rational functions on A which constitute a local coordinate system at 0 and their reductions modulo $\mathfrak{p}$ are k -rational functions on $A \times_{\mathcal{O}} k$ also constitute a local coordinate system at the identity point of $A \times_{\mathcal{O}} k$.

Let $A(K)$ be the group of K -rational points of A and let U be an open neighborhood of 0 in $A(K)$. A function $\Phi: U \rightarrow K$ is called $\mathcal{O}$ -analytic on U if for $a \in U$ we have

$$\Phi(a) = \phi(t_1(a), \dots, t_d(a)),$$

where $\phi = \phi(\tau_1, \dots, \tau_d)$ is a d -variable power series with coefficients in $\mathcal{O}$ which converges at $(t_1(a), \dots, t_d(a))$ for all $a \in U$.

A K -rational divisor Δ on A is called disjoint from $0 \bmod \mathfrak{p}$ if any component of the set obtained from the reduction mod $\mathfrak{p}$ of the support of Δ does not contain the identity point of $A \times_{\mathcal{O}} k$.

For a subgroup G of $A(K)$, let $\Lambda = \mathbf{Z}[G]$ be the group ring of G with coefficients in $\mathbf{Z}$, which is also the group of 0 -cycles with components in G . Let $I \subset \Lambda$ be the augmentation ideal of Λ ; i.e., the ideal generated by the cycles of the form $(a) - (0)$ with $a \in G$. Let $I^2 \subset \Lambda$ be the ideal generated by the cycles of the form $(a + b) - (a) - (b) + (0)$ with $a, b \in G$. As $(a + b) - (a) - (b) + (0) = ((a) - (0)) * ((b) - (0))$, where $*$ is the multiplication in Λ , I^2 coincides with the square of I ; $I^2 = \{a * b \mid a, b \in I\}$.

A divisor and a 0 -cycle are called disjoint if their supports are disjoint. For a divisor $\Delta = \text{div}(f)$ linearly equivalent to 0 , and a 0 -cycle $\mathfrak{a} \in I$ which are disjoint, we define a pairing $[\Delta, \mathfrak{a}]$ by $[\Delta, \mathfrak{a}] = \prod f(a_i)^{m_i}$ for $\mathfrak{a} = \sum m_i(a_i)$, as usual. For a divisor Δ algebraically equivalent to 0 and a 0 -cycle $\mathfrak{a} = \sum m_i(a_i)$, let $\Delta * \mathfrak{a} = \sum m_i \Delta_{a_i}$, where Δ_{a_i} is the translate of Δ by a_i , and let $\mathfrak{a}^- = \sum m_i(-a_i)$. If $\mathfrak{a} \in I$, then $\Delta * \mathfrak{a}$ is linearly equivalent to 0 . So for $\mathfrak{a} = \mathfrak{b} * \mathfrak{c} \in I^2$, Δ algebraically

Received by the editors August 7, 1985 and, in revised form, March 21, 1986.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 14K15.

equivalent to 0, the value $[\Delta * b^-, c]$ has a meaning. We define the pairing $[\Delta, a]$ by $[\Delta, a] = [\Delta * b^-, c]$. By a certain reciprocity law, this pairing is known to be well defined.

In [4], A. Néron has shown the following fact: Let G be an open subgroup of $A(K)$ such that $t_i(a) \in p\mathcal{O}$ for $a \in G$, $i = 1, \dots, d$, where $t_1, \dots, t_d$ are p -admissible system of local coordinates of A at 0. Let Δ be a K -rational divisor on A algebraically equivalent to 0 and disjoint from 0 mod p . Then for $a \in I = I_G$,

$$\theta'_{\Delta,0}(a) = \lim_{\nu \rightarrow \infty} [\Delta, p^\nu a - p^\nu \delta a]^{1/p^\nu}$$

converges and satisfies $\theta'_{\Delta,0}(0) = 1$, $\theta'_{\Delta,0}(a) = [\Delta, a]$ for $a \in I^2$. The function $\theta'_{\Delta,0}(a) = \theta'_{\Delta,0}((a) - (0))$ on $a \in G$ is an $\mathcal{O}$ -analytic function with values in $1 + p\mathcal{O}$ (cf. [4, Théorème and its proof]).

In general, for a K -rational divisor Δ not necessarily disjoint from 0 mod p , take a K -rational function f on A such that $\text{div}(f) + \Delta = \Delta'$ is disjoint from 0 mod p , and put $\theta'_{\Delta,0}(a) = \theta'_{\Delta',0}(a)f(a)^{-1}$.

REMARK. Note that for a divisor $\Delta = \text{div}(f)$ linearly equivalent to 0, Néron's p -adic theta function $\theta'_{\Delta,0}$ differs from the function f slightly (cf. [4, §4(b)]). So there are some ambiguities in the above definition of $\theta'_{\Delta,0}$ in the case of Δ not disjoint from 0 mod p . However from op. cit., for $a \in I$, $\theta'_{\Delta,0}(a)$ is defined up to units so this ambiguity disappears when we take the image of $\theta'_{\Delta,0}(a)$ by a homomorphism which is trivial on the unit group.

In §3 we shall prove the following Proposition.

PROPOSITION. *Let Ω be the completion of the algebraic closure of K , R the integer ring of Ω , and $\mathfrak{m}$ the maximal ideal of R . Let $\hat{A}(R) = \{a \in A(\Omega) | t_i(a) \in \mathfrak{m}, i = 1, \dots, d\}$. Assume that A has ordinary good reduction mod p . Let Δ be a K -rational divisor on A algebraically equivalent to 0 and disjoint from 0 mod p . Then the limit*

$$\theta'_{\Delta,0}(a) = \lim_{\nu \rightarrow \infty} [\Delta, p^\nu a - p^\nu \delta a]^{1/p^\nu}$$

converges for $a = (a) - (0)$ with $a \in \hat{A}(R)$, and $\theta'_{\Delta,0}(a) = \theta'_{\Delta,0}((a) - (0))$ defines an $\mathcal{O}$ -analytic function on $\hat{A}(R)$ with values in $1 + \mathfrak{m}$, satisfying $\theta'_{\Delta,0}(a) = [\Delta, a]$ for $a \in I^2 = I_{\hat{A}(R)}^2$. Moreover, $\theta'_{\Delta,0}$ is the unique $\mathcal{O}$ -analytic function on $\hat{A}(R)$ which satisfies $\theta'_{\Delta,0}(a) = [\Delta, a]$ for $a \in I^2$.

A. Néron has extended his p -adic theta function $\theta'_{\Delta,0}$ to $A(K)$ in the following way. Let Δ be a K -rational divisor on A algebraically equivalent to 0 and let $a = (a) - (0)$ be a 0-cycle disjoint from Δ with $a \in A(K)$. Let m be the exponent of the finite group $A(K)/G$ (G is an open subgroup of $A(K)$ such that $t_i(a) \in p\mathcal{O}$ for $a \in G$, $i = 1, \dots, d$). Then define the extension $\tilde{\theta}'_{\Delta,0}$ of $\theta'_{\Delta,0}$ by

$$[\Delta, ma - m\delta a] = \tilde{\theta}'_{\Delta,0}(a)^m \theta'_{\Delta,0}(m\delta a)^{-1}.$$

So $\tilde{\theta}'_{\Delta,0}(a) = \tilde{\theta}'_{\Delta,0}((a) - (0))$ is defined up to m th root of unity (with value in the algebraic closure of K).

Now let K be an algebraic number field of finite degree. Let A be an abelian variety defined over K . For a nonarchimedean prime v of K , let K_v be the completion of K at v , $\mathcal{O}_v$ the integer ring of K_v , $\mathfrak{p}_v$ the maximal ideal of $\mathcal{O}_v$, and let

$k_v = \mathcal{O}_v/\mathfrak{p}_v$ be the residue field at v . Let $\mathbf{A}_K^\times$ be the group of finite ideles of K ; i.e., $\mathbf{A}_K^\times$ is the restricted direct product $\prod' K_v^\times$ where the product ranges through all of the nonarchimedean primes of K and (x_v) is in $\mathbf{A}_K^\times$ if $x_v \in \mathcal{O}_v^\times$ for all but a finite number of primes v . We have a canonical homomorphism $K^\times \rightarrow \mathbf{A}_K^\times$ and we identify $K^\times$ with its image. Let p be a fixed prime number such that A has ordinary good reduction at all primes of K over p . Let $\rho: \mathbf{A}_K^\times \rightarrow \mathbf{Z}_p$ be a nontrivial continuous homomorphism which is trivial on $K^\times$ corresponding to a $\mathbf{Z}_p$ -extension of K by the class field theory. For a nonarchimedean prime v , let ρ_v be the component of ρ at v .

Let Δ be a K -rational divisor on A algebraically equivalent to 0 and $a \in A(K)$, which are disjoint. For a nonarchimedean prime v , we can construct the v -adic theta function $\tilde{\theta}'_{\Delta,0,v}(a)$ substituting A, K_v instead of A, K in the above construction.

Then A. Néron's p -adic height corresponding to the homomorphism ρ is defined by

$$h_{\Delta,\rho}(a) = \sum \rho_v(\tilde{\theta}'_{\Delta,0,v}(a)),$$

where the summation ranges through all of the nonarchimedean primes of K , and as $\tilde{\theta}'_{\Delta,0,v}(a)^m \in K_v^\times$ for some m , we put

$$\rho_v(\tilde{\theta}'_{\Delta,0,v}(a)) = \frac{1}{m} \rho_v(\tilde{\theta}'_{\Delta,0,v}(a)^m).$$

It can be shown that this function is well defined and depends only on the linear equivalence class of Δ .

Let $A' = \text{Pic}^0(A)$ be the dual abelian variety. Then we obtain a pairing from $A(K) \times A'(K)$ to $\mathbf{Q}_p$ by putting

$$\langle a, \text{cl}(\Delta) \rangle_N = h_{\Delta,\rho}(a),$$

here $\text{cl}(\Delta)$ is the image of Δ in A' .

Next we recall the construction of P. Schneider's p -adic height pairing (as P. Schneider's p -adic height pairing coincides with Mazur and Tate's p -adic height pairing (cf. [3, §4.4]), we recall the latter).

Let K be an algebraic number field of finite degree and A an abelian variety defined over K . Let $\mathcal{O}$ be the integer ring of K and let $\mathcal{A}$ be the Néron model of A over $\mathcal{O}$. Let $\mathcal{A}^0$ be the connected component of the identity of $\mathcal{A}$ and let A' be the dual abelian variety of A . For a K -rational divisor Δ of A algebraically equivalent to 0 there corresponds an extension X of $\mathcal{A}^0$ by the multiplicative group $\mathbf{G}_m$ over $\mathcal{O}$ (cf. [1, §1]);

$$(1) \quad 0 \rightarrow \mathbf{G}_m \rightarrow X \xrightarrow{\pi} \mathcal{A}^0 \rightarrow 0.$$

Here we note that for an extension field L of K , the group of L -rational points $X(L)$ of X may be described as follows (cf. [3, p. 210]): $X(L)$ is the set of pairs $(\mathfrak{a}, c)$ where $\mathfrak{a}$ is a 0-cycle of degree 0 with L -rational components and $c \in L^\times$; $c(\mathfrak{a}, 1) = (\mathfrak{a}, c)$, $(\mathfrak{a}_1, c_1)(\mathfrak{a}_2, c_2) = (\mathfrak{a}_1 + \mathfrak{a}_2, c_1 c_2)$, when $(\mathfrak{a}, c)$ maps to $0 \in A(L)$ (i.e., when $\mathfrak{a} \in I^2$ is Néron's notation recalled as before), then $(\mathfrak{a}, c) = (0, [\Delta, \mathfrak{a}]c)$. For $a \in A(L)$, the map $\sigma_\Delta: a \mapsto ((a) - (0), 1)$ defines a K -rational section of π , and σ_Δ is regular on $A(L) \setminus |\Delta|$.

Let p be a fixed prime number such that A has ordinary good reduction at the primes of K over p . Let $\rho: \mathbf{A}_K^\times \rightarrow \mathbf{Z}_p$ be a nontrivial continuous homomorphism

which is trivial on $K^\times$ corresponding to a $\mathbf{Z}_p$ -extension of K , where $\mathbf{A}_K^\times$ is the group of finite ideles of K as before. For a nonarchimedean prime v of K , let ρ_v be the component of ρ at v .

By considering the K_v -rational points of (1) we obtain the following exact sequence:

$$0 \rightarrow K_v^\times \rightarrow X(K_v) \rightarrow A(K_v) \rightarrow 0$$

(note that $H^1(R, \mathbf{G}_m) = 0$ for a local ring R). In [3, Theorem (1.5)], Mazur and Tate have constructed a canonical ρ_v -splitting $\psi_{\rho,v}: X(K_v) \rightarrow \mathbf{Q}_p$ for the above exact sequence (in our case, canonical ρ_v -splitting is the restriction to the fiber at $\text{cl}(\Delta) \in A'(K)$ of their canonical ρ_v -splitting of the biextension of $A(K_v) \times A'(K_v)$ by $K_v^\times$). We recall that a homomorphism $\psi_{\rho,v}: X(K_v) \rightarrow \mathbf{Q}_p$ is called a ρ_v -splitting if it satisfies

$$\psi_{\rho,v}(cx) = \rho_v(c) + \psi_{\rho,v}(x) \quad \text{for } c \in K_v^\times, x \in X(K_v).$$

(The characterizing property of canonicalness will be explained later.)

Now let $a \in A(K)$ and take a point $x \in X(K)$ mapping to a . Mazur and Tate have defined their p -adic height pairing by

$$\langle a, \text{cl}(\Delta) \rangle_S = \sum \psi_{\rho,v}(x),$$

where the summation ranges through the nonarchimedean primes of K . Here we note that the sum on the right is well defined and independent of the choice of x and depends only on the linear equivalence class of Δ .

2. The result of this note is the following Theorem.

THEOREM. *Let K be an algebraic number field of finite degree and A an abelian variety defined over K . Let p be a prime number such that A has ordinary good reduction at all primes of K over p . Let $\rho: \mathbf{A}_K^\times \rightarrow \mathbf{Z}_p$ be a nontrivial continuous homomorphism trivial on $K^\times$, corresponding to a $\mathbf{Z}_p$ -extension of K . Then for $a \in A(K)$ and Δ a K -rational divisor on A algebraically equivalent to 0, we have*

$$\langle a, \text{cl}(\Delta) \rangle_N = \langle a, \text{cl}(\Delta) \rangle_S.$$

PROOF. Let the notations be the same as in §1. Let v be a nonarchimedean prime of K . We take a divisor from the linear equivalence class of Δ , which is disjoint from 0 mod v for all primes v over p (for simplicity, we write this divisor by Δ). We compare the local components of Néron's and Schneider's p -adic height pairings.

First we treat the case where v is a prime not above p . In this case ρ_v is trivial on $\mathcal{O}_v^\times$ as ρ is continuous. Let q be the residue characteristic of v , and let G be an open subgroup of $\mathcal{A}^0(\mathcal{O}_v)$ such that $t_i(a) \in q\mathcal{O}_v$ for $a \in G$, $i = 1, \dots, d$; here $t_1, \dots, t_d$ are v -admissible systems of local coordinates of A at 0.

If $\Delta = \text{div}(f)$ is linearly equivalent to 0, then from [3, (2.2.2)], we have $\psi_{\rho,v}((a, 1)) = \rho_v(f(a))$ for $a \in I = I_G$, whenever the right-hand side is defined. On the other hand, from the Remark before the Proposition in §1 (cf. also [4, §4(b)]), $\rho_v(\theta'_{\Delta,0}(a)) = \rho_v(f(a))$.

In general let Δ be a divisor algebraically equivalent to 0. Take a K_v -rational function on A such that $\text{div}(f) + \Delta = \Delta'$ is disjoint from 0 mod v . We consider the objects of §1 defined with respect to Δ' . Now in the case we are considering, the

characterizing property of canonicalness of ρ_v -splitting is that $\psi_{\rho,v} = 0$ on $X(\mathcal{O}_v)$, where X is the extension of $\mathcal{A}^0$ by $\mathbf{G}_m$ corresponding to Δ' . Let $m = [A(K_v) : G]$, where G is an open subgroup of $\mathcal{A}^0(\mathcal{O}_v)$ as above. Then we have for $\mathfrak{a} = (a) - (0)$, $a \in A(K_v)$.

$$\begin{aligned}\psi_{\rho,v}((\mathfrak{a}, 1)) &= \frac{1}{m} \psi_{\rho,v}((m\mathfrak{a}, 1)) \\ &= \frac{1}{m} \psi_{\rho,v}((0, [\Delta', m\mathfrak{a} - m\delta\mathfrak{a}])) + \frac{1}{m} \psi_{\rho,v}((m\delta\mathfrak{a}, 1)) \\ &= \frac{1}{m} \rho_v([\Delta', m\mathfrak{a} - m\delta\mathfrak{a}]),\end{aligned}$$

as $m\delta\mathfrak{a}$ has support in $\mathcal{A}^0(\mathcal{O}_v)$.

On the other hand from the equation defining the v -adic theta function $\tilde{\theta}'_{\Delta',0,v}$, we have

$$\frac{1}{m} \rho_v([\Delta', m\mathfrak{a} - m\delta\mathfrak{a}]) = \rho_v(\tilde{\theta}'_{\Delta',0,v}(\mathfrak{a})),$$

as $\theta'_{\Delta',0,v}(m\delta\mathfrak{a}) \in 1 + q\mathcal{O}_v$.

From the multiplicativity of the local pairings with respect to Δ , we see that in this case the local components of Néron's and Schneider's p -adic height pairings coincide.

In the case when v is a prime over p , we consider the fiber at v of the exact sequence (1). Taking the formal completion along the maximal torus of the special fiber (we denote this completion by superscript t) we obtain the following exact sequence (cf. [3, §5])

$$0 \rightarrow \mathbf{G}_m^t \rightarrow X^t \rightarrow A^t \rightarrow 0.$$

Mazur and Tate have proved that the above exact sequence has a unique splitting $\psi: X^t \rightarrow \mathbf{G}_m^t$. The characterizing property of canonicalness of ρ_v -splitting is that $\psi_{\rho,v} = \rho_v \circ \psi$ on $X^t(\mathcal{O}_v)$. Now take the formal completion along the 0-section of the special fiber of the exact sequence (1) (we denote this completion by $\hat{X}$ etc.); we obtain the following exact sequence:

$$0 \rightarrow \hat{\mathbf{G}}_m \rightarrow \hat{X} \rightarrow \hat{A} \rightarrow 0.$$

Let Ω_v be the completion of the algebraic closure of K_v , R_v the integer ring of Ω_v , and let $\mathfrak{m}_v$ be the maximal ideal of R_v . Considering the R_v -valued points we obtain the following exact sequence:

$$(2) \quad 0 \rightarrow 1 + \mathfrak{m}_v \rightarrow \hat{X}(R_v) \rightarrow \hat{A}(R_v) \rightarrow 0.$$

From Mazur and Tate's splitting $\psi: X^t \rightarrow \mathbf{G}_m^t$, we obtain a splitting $\psi': \hat{X}(R_v) \rightarrow 1 + \mathfrak{m}_v$ of the above exact sequence. Now Δ is disjoint from $0 \bmod v$. From the Proposition in §1, for $\mathfrak{a} = (a) - (0)$ with $a \in \hat{A}(R_v)$,

$$\theta'_{\Delta,0,v}(\mathfrak{a}) = \lim_{\nu \rightarrow \infty} [\Delta, p^\nu \mathfrak{a} - p^\nu \delta \mathfrak{a}]^{1/p^\nu}$$

converges and defines an $\mathcal{O}_v$ -analytic function on $\hat{A}(R_v)$. From $\theta'_{\Delta,0,v}(\mathfrak{a}) = [\Delta, \mathfrak{a}]$ for $\mathfrak{a} \in I^2$, by putting $\mathfrak{a} = (a + b) - (a) - (b) + (0)$ with $a, b \in \hat{A}(R_v)$ we have

$$\theta'_{\Delta,0,v}(a + b) \theta'_{\Delta,0,v}(a)^{-1} \theta'_{\Delta,0,v}(b)^{-1} = [\Delta, \mathfrak{a}] = \sigma_\Delta(a + b) \sigma_\Delta(a)^{-1} \sigma_\Delta(b)^{-1};$$

here $\sigma_\Delta: a \mapsto ((a) - (0), 1)$ is the K -rational section of $\pi: X \rightarrow A$ such that $\sigma_\Delta(0) = 1$. Hence $\sigma_\Delta \cdot \theta'_{\Delta,0,v}{}^{-1}: \hat{A}(R_v) \rightarrow \hat{X}(R_v)$ is a section of (2) as abstract groups. As $\theta'_{\Delta,0,v}$ is $\mathcal{O}_v$ -analytic and σ_Δ is K -rational, from [8, (4.2), Corollary 1], $\sigma_\Delta \cdot \theta'_{\Delta,0,v}{}^{-1}$ is in fact a formal group homomorphism defined over $\mathcal{O}_v$. We define $\psi'': \hat{X}(R_v) \rightarrow 1 + \mathfrak{m}_v$ by

$$\psi''(x) = x/(\sigma_\Delta \cdot \theta'_{\Delta,0,v}{}^{-1}(\pi(x))),$$

then ψ'' is a splitting of (2) as abstract groups. As $\psi' \cdot \psi''^{-1}$ is trivial on $1 + \mathfrak{m}_v$, it induces a formal group homomorphism $\psi' \cdot \psi''^{-1}: \hat{A} \rightarrow \hat{\mathbf{G}}_m$ defined over $\mathcal{O}_v$ (cf. *ibid.*). We need the following Lemma.

LEMMA. *Let $\hat{A}$ be the formal group of an abelian variety defined over K_v , which has good reduction mod v . Then any formal group homomorphism $\phi: \hat{A} \rightarrow \hat{\mathbf{G}}_m$ defined over $\mathcal{O}_v$ is trivial.*

PROOF OF THE LEMMA. If ϕ were nontrivial, there should exist a surjective homomorphism ψ such that $\phi = \psi^m$ for some m (cf. [8, (4.2), Corollary 1]). The reduction mod v of the homomorphism ψ is a surjective homomorphism of formal groups over k_v , $\psi/k_v: \hat{A}/k_v \rightarrow \hat{\mathbf{G}}_m/k_v$. Consider the characteristic polynomials of the Frobenius endomorphism of $\bar{k}_v$ over k_v acting on these p -divisible groups over k_v (cf. [2, §4.e]). The multiplicativity of the characteristic polynomials and the Riemann-Weil hypothesis lead to a contradiction.

By the Lemma, $\psi' = \psi''$. We apply the Mazur-Tate's construction of p -adic height pairing recalled as before. Let $m = [A(K_v): \hat{A}(\mathcal{O}_v)]$, which is the cardinality of the group of rational points of the special fiber of A at v . As before, let $\mathfrak{a} = (a) - (0)$. Then we have

$$\begin{aligned} \psi_{p,v}((\mathfrak{a}, 1)) &= \frac{1}{m} \psi_{p,v}((m\mathfrak{a}, 1)) \\ &= \frac{1}{m} \rho_v([\Delta, m\mathfrak{a} - m\delta\mathfrak{a}]) + \frac{1}{m} \rho_v \circ \psi'((m\delta\mathfrak{a}, 1)) \\ &= \frac{1}{m} \rho_v([\Delta, m\mathfrak{a} - m\delta\mathfrak{a}]) + \frac{1}{m} \rho_v \circ \psi''(\sigma_\Delta(m\mathfrak{a})) \\ &= \frac{1}{m} \rho_v([\Delta, m\mathfrak{a} - m\delta\mathfrak{a}]) + \frac{1}{m} \rho_v \circ \theta'_{\Delta,0,v}(m\mathfrak{a}) \\ &= \rho_v(\tilde{\theta}'_{\Delta,0,v}(\mathfrak{a})). \end{aligned}$$

So $\psi_{p,v}((\mathfrak{a}, 1)) = \rho_v(\tilde{\theta}'_{\Delta,0,v}(\mathfrak{a}))$ for all nonarchimedean primes v . The Theorem is proved.

3. In this section we prove the Proposition of §1. Let the notations be the same as in the Proposition. Let f be the K -rational function on A such that $\text{div}(f) = (p\delta)^{-1}\Delta - p\Delta$, $f(0) = 1$. We expand f as a power series in $t_1, \dots, t_d$ near 0, then from [4, p. 158], we see that $f \in \mathcal{O}[[t_1, \dots, t_d]]$ and $f - 1 \equiv 0 \pmod{\deg 2}$. Also from the arguments used in *ibid.* p. 158, to prove the convergence of $\theta'_{\Delta,0}$, it suffices to show that for $a \in \hat{A}(R)$, $\text{ord}_p((f(p^\nu a) - 1)/p^\nu)$ diverges to ∞ as $\nu \rightarrow \infty$, where ord_p is the additive valuation such $\text{ord}_p(p) = 1$. Let $\hat{A}$ be the formal group of A , and let I be the integer ring of the completion of the maximal unramified

extension of K . Then $\hat{A}$ is isomorphic to $\hat{G}_m^d$ over I . Hence $\hat{A}(R) \simeq (1 + \mathfrak{m})^d$ over I . As this isomorphism may be expressed by some invertible power series, we see that $\text{ord}_p(t_i(p^\nu a)) = \text{ord}_p((1+z)^{p^\nu} - 1)$ for some $z \in \mathfrak{m}$. We expand $(1+z)^{p^\nu}$ with binomial coefficients $\binom{p^\nu}{r}$. For $r < p^{\nu-1}$, using the fact that $\text{ord}_p(r!) = \sum_{k=0}^{\infty} [r/p^k]$, where $[x]$ is the largest integer such that $[x] \leq x$, after some computations we have $\text{ord}_p\left(\binom{p^\nu}{r}\right) = \nu - \text{ord}_p(r)$. Hence

$$\text{ord}_p((1+z)^{p^\nu} - 1) \geq \text{Min}(p^{\nu-1} \text{ord}_p z, \nu - \text{ord}_p r + r \text{ord}_p z),$$

where r ranges from 1 to $p^{\nu-1} - 1$.

As $\text{ord}_p(t_i(p^\nu a))$ satisfies similar inequality, we see that

$$\text{ord}_p((f(p^\nu a) - 1)/p^\nu) \geq \text{Min}(2p^{\nu-1} \text{ord}_p(z_i) - \nu, \nu - 2 \text{ord}_p r + 2r \text{ord}_p(z_i)),$$

where r ranges from 1 to $p^{\nu-1} - 1$, i ranges from 1 to d , and z_i are elements in $\mathfrak{m}$.

From $\text{ord}_p(z_i) > 0$, we see easily that the right-hand side of the above inequality diverges to ∞ when $\nu \rightarrow \infty$.

When $\text{ord}_p(z_i)$ ranges through a compact subset of $\{x \in \mathbf{R} | x > 0\}$, this convergence is uniform. Hence $\theta'_{\Delta,0}$ defines an $\mathcal{O}$ -analytic function on $\hat{A}(R)$.

Now let $\mathfrak{a} = (a+b) - (a) - (b) + (0) \in I^2$. By Néron's result we have $\theta'_{\Delta,0}(\mathfrak{a}) = [\Delta, \mathfrak{a}]$ for $a, b \in \hat{A}(R)$ sufficiently close to 0 (cf. [4, Théorème]). For a fixed b , both sides of the above equation are given by power series in coordinates of a . Hence we have $\theta'_{\Delta,0}(\mathfrak{a}) = [\Delta, \mathfrak{a}]$ for all $\mathfrak{a} \in I^2$. Uniqueness is clear from the Lemma in §2.

REFERENCES

1. S. Bloch, *A note on height pairings, Tamagawa numbers and the Birch and Swinnerton-Dyer conjecture*, Invent. Math. **58** (1980), 65–76.
2. B. Mazur, *Rational points of abelian varieties with values in towers of number fields*, Invent. Math. **18** (1972), 183–266.
3. B. Mazur and J. Tate, *Canonical height pairing via biextensions*, Arithmetic and Geometry (Volume dedicated to Shafarevich, vol. 1), Progress in Math., vol. 35, Birkhäuser, Boston, Basel, and Stuttgart, 1983, pp. 195–273.
4. A. Néron, *Fonctions θ - p -adiques et hauteurs p -adiques* (Séminaire de théorie de nombres, Paris 1980–1981), Progress in Math., vol. 22, Birkhäuser, Boston, Basel, and Stuttgart, 1982, pp. 149–174.
5. J. Oesterlé, *Construction de hauteurs archimédiennes et p -adiques suivant la méthode de Bloch* (Séminaire de théorie de nombres, Paris 1980–1981), Progress in Math., vol. 22, Birkhäuser, Boston, Basel, and Stuttgart, 1982, pp. 175–192.
6. B. Perrin-Riou, *Hauteurs p -adiques* (Séminaire de théorie de nombres, Paris 1982–1983), Progress in Math., vol. 51, Birkhäuser, Boston, Basel, and Stuttgart, 1984, pp. 233–257.
7. P. Schneider, *p -adic height pairings. I*, Invent. Math. **69** (1982), 401–409.
8. J. Tate, *p -divisible groups*, Proceedings of a Conference on Local Fields, NUFFIC Summer School held at Driebergen, 1966, Springer, Berlin, Heidelberg, and New York, 1967, pp. 158–183.

DEPARTMENT OF MATHEMATICS, COLLEGE OF GENERAL EDUCATION, TÔHOKU UNIVERSITY, SENDAI, 980, JAPAN