TWO-GENERATOR GROUPS WITH PERFECT FRATTINI SUBGROUPS

M. J. EVANS

ABSTRACT. In this paper, 2-generator groups with perfect Frattini subgroups are constructed. The construction also shows that every countable group can be embedded in the Frattini subgroup of a 2-generator group.

1. Introduction. It is well-known that the Frattini subgroup $\Phi(G)$ of a finite group G is nilpotent. However, for infinite groups the situation is considerably more complicated. For instance, in [1] P. Hall constructed infinite 2-generator soluble groups with non-nilpotent Frattini subgroups. More recently, M. F. Newman (unpublished) has constructed finitely generated, infinite p -groups with insoluble Frattini subgroups.

In this note we use a construction similar to Hall's (loc. cit.) to obtain 2-generator groups with perfect Frattini subgroups. More precisely, we prove

THEOREM 1.1. *Let K be a countable perfect group. Then there exists a 2-generator group G_K , such that $\Phi(G_K)$ is a direct product of $\aleph_0$ copies of K .*

Since each countable group can be embedded in a countable, nonabelian, simple (and therefore perfect) group (e.g. [2, Theorem 3.4]), Theorem 1.1 implies the following.

COROLLARY 1.2. *Every countable group can be embedded in the Frattini subgroup of a 2-generator group.*

2. The construction. Throughout the sequel M denotes a countable perfect group of infinite exponent.

Suppose $M = \{m_1, m_2, m_3, \dots\}$ and let S denote the sequence $m_1, m_2, m_1, m_2, m_3, m_1, m_2, m_3, m_4, m_1, \dots, m_i, m_1, \dots, m_i, m_{i+1}, m_1, \dots$. Write S_j for the j th term of S .

REMARK. Note that for any integer N and any $m \in M$ there exists an integer $n > N$ such that $S_n = m$. This fact will be crucial in the proof of Lemma 3.1.

Let $C = \text{Cr}_{i \in \mathbf{Z}} M_i$ and $H = \text{Dr}_{i \in \mathbf{Z}} M_i$, where $M_i \simeq M$ for all $i \in \mathbf{Z}$. Thus C is the cartesian product and H the direct product of $\aleph_0$ copies of M . Given any $g \in C$ we write $(g)_i$ for the i th coordinate of g and define the *support* of g to be the set $\sigma(g) = \{i | (g)_i \neq 1\}$. Clearly $g \in H$ if and only if $\sigma(g)$ is a finite set.

Let z generate an infinite cyclic group that is disjoint from C and define an action of $\langle z \rangle$ on C by

$$(g^z)_{i+1} = (g)_i, \quad \text{for all } g \in C \text{ and } i \in \mathbf{Z}.$$

Define C^* to be the split extension of C by $\langle z \rangle$.

Received by the editors March 6, 1986.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 20F26, 20E28.

©1987 American Mathematical Society
 0002-9939/87 \$1.00 + \$.25 per page

Let $w \in C$ be the element such that

$$(w)_{2^j} = S_j \quad \text{if } j \geq 1, \\ (w)_l = 1 \quad \text{if } l \neq 2^j \text{ for all } j \geq 1.$$

Finally, define $G_M = \langle H, w, z \rangle$, a subgroup of C^* . Observe that $M_i^z = M_{i+1}$ and $M_i^w = M_i$, for all $i \in \mathbf{Z}$, so $H \triangleleft G_M$.

The notation introduced above will be used for the remainder of this note. We shall show that $\Phi(G_M) = H$ and use this fact to deduce Theorem 1.1.

3. The lemmas. Our main lemma is the following.

LEMMA 3.1. *Let h_1 and h_2 be arbitrary elements of H . Then $G_M = \langle h_1 w, h_2 z \rangle$.*

PROOF. Since $\sigma(h_1)$ and $\sigma(h_2)$ are finite sets, there exists an integer $N > 0$ such that $(h_1 w)_n = (w)_n$ and $(h_2)_n = 1$ for all $n \geq N$. We consider such an N fixed once for all and define p to be the least integer such that $2^p \geq N$. We shall prove that $M_{2^p} \leq \langle h_1 w, h_2 z \rangle$. Since $(h_2 z)^{-k} M_{2^p} (h_2 z)^k = M_{2^p+k}$, for all $k \in \mathbf{Z}$, and $H = \text{Dr}_{i \in \mathbf{Z}} M_i$, the result then follows immediately.

As M is perfect, it suffices to show that, given arbitrary elements $m_k, m_j \in M$, there exists $y \in \langle h_1 w, h_2 z \rangle \cap H$ such that

$$(1) \quad (y)_{2^p} = [m_k, m_j]$$

and

$$(2) \quad (y)_l = 1 \quad \text{if } l \neq 2^p.$$

Suppose that m_k and m_j are given elements of M and let t denote the least integer in $\sigma(h_1 w)$. It follows from the remark in §2 that there exists an integer $r > p$ such that $S_r = m_k$. Furthermore, there exists an integer $u > r$ such that $S_u = m_j$ and $2^r - t < 2^{u-1}$. The letters r and u retain this meaning throughout.

Define $\lambda_1, \lambda_2 \in \langle h_1 w, h_2 z \rangle$ by $\lambda_i = (h_2 z)^{q_i} h_1 w (h_2 z)^{-q_i}$, where $q_1 = 2^r - 2^p$ and $q_2 = 2^u - 2^p$. We claim that $y = [\lambda_1, \lambda_2]$ is the desired element satisfying (1) and (2).

Bearing in mind that $q_i > 0$, for $i = 1, 2$, we have

$$(h_2 z)^{q_i} = h_2 h_2^{z^{-1}} \cdots h_2^{z^{-(q_i-1)}} z^{q_i} = f_i z^{q_i}, \quad \text{say,}$$

where $f_i \in H$. Thus $(\lambda_i)_{2^p} = (f_i)_{2^p} ((h_1 w)^{z^{-q_i}})_{2^p} (f_i^{-1})_{2^p}$. However, $f_i = \prod_{a=0}^{q_i-1} h_2^{z^{-a}}$ so

$$(f_i)_{2^p} = \prod_{a=0}^{q_i-1} (h_2^{z^{-a}})_{2^p} = \prod_{a=0}^{q_i-1} (h_2)_{2^p+a}$$

for $i = 1, 2$. Now $2^p + a \geq N$ for $a = 0, 1, \dots, q_i - 1$ and, by our choice of N , $(h_2)_n = 1$ for all $n \geq N$. Hence $(f_i)_{2^p} = 1$ for $i = 1, 2$. Therefore $(\lambda_i)_{2^p} = ((h_1 w)^{z^{-q_i}})_{2^p} = (h_1 w)_{2^p+q_i}$ for $i = 1, 2$. Recall that $(h_1 w)_n = (w)_n$ for all $n \geq N$. It follows that $(\lambda_1)_{2^p} = (h_1 w)_{2^r} = (w)_{2^r} = S_r = m_k$ by our choice of r , and similarly $(\lambda_2)_{2^p} = (h_1 w)_{2^u} = S_u = m_j$. Hence $([\lambda_1, \lambda_2])_{2^p} = [(\lambda_1)_{2^p}, (\lambda_2)_{2^p}] = [m_k, m_j]$ and $y = [\lambda_1, \lambda_2]$ satisfies (1).

We next show that $\sigma(\lambda_1) \cap \sigma(\lambda_2) \subseteq \{2^p\}$. Suppose that $b \in \sigma(\lambda_1) \cap \sigma(\lambda_2)$ and observe that $\sigma(\lambda_i) = \{v - q_i | v \in \sigma(h_1 w)\}$, for $i = 1, 2$. It follows that $b = v_1 - q_1 =$

$v_2 - q_2$ where $v_1, v_2 \in \sigma(h_1 w)$. Therefore $v_2 = v_1 - q_1 + q_2 = v_1 - 2^r + 2^u$ and as t is the least integer in $\sigma(h_1 w)$ this implies that $v_2 \geq t - 2^r + 2^u$. However $2^r - t < 2^{u-1}$, so $v_2 > 2^{u-1} \geq N$. It follows from our choice of N and the fact that $v_2 \in \sigma(h_1 w)$ that $v_2 = 2^{n_2}$ for some integer $n_2 \geq u$. Now $v_1 = 2^{n_2} + q_1 - q_2 = 2^{n_2} + 2^r - 2^u \geq 2^r \geq N$ so $v_1 = 2^{n_1}$ for some integer $n_1 \geq r$. Hence $b = 2^{n_1} + 2^p - 2^r = 2^{n_2} + 2^p - 2^u$ so $2^r(2^{n_1-r} - 1) = 2^r(2^{n_2-r} - 2^{u-r})$ which implies that $n_1 = r$ and $n_2 = u$. Thus $b = 2^p$ and $\sigma(\lambda_1) \cap \sigma(\lambda_2) \subseteq \{2^p\}$.

Note that $\sigma([\lambda_1, \lambda_2]) \subseteq \sigma(\lambda_1) \cap \sigma(\lambda_2)$ so $([\lambda_1, \lambda_2])_l = 1$ if $l \neq 2^p$ and $y = [\lambda_1, \lambda_2]$ satisfies (2). The proof is complete.

Lemma 3.1 has the following easy consequence.

LEMMA 3.2. *The subgroup H of G_M is omissible.*

PROOF. Suppose $G_M = \langle H, X \rangle$ where X is a subset of G_M . Define $N = \langle X \rangle$ so that $G = HN$. Now there exists $h_1, h_2 \in H$ and $n_1, n_2 \in N$ such that $w = h_1^{-1}n_1$ and $z = h_2^{-1}n_2$. Thus $h_1 w, h_2 z \in N$, so $N = G_M$, by Lemma 3.1. Therefore $N = G_M = \langle X \rangle$, as required.

Of course, Lemma 3.2 implies that $H \leq \Phi(G_M)$.

Our present goal, as mentioned at the end of §2, is to show that $\Phi(G_M) = H$. We have established that $\Phi(G_M) \geq H$, so it is enough to prove that $\Phi(G_M/H) = 1$. A routine argument, along the lines of [1, Theorem 7], and using the fact that M is of infinite exponent, shows that $G_M/H \simeq \mathbf{Z} \wr \mathbf{Z}$, the restricted wreath product of two infinite cyclic groups. Moreover, it is easily seen that the Frattini subgroup of $\mathbf{Z} \wr \mathbf{Z}$ is trivial. Summing up, we have

LEMMA 3.3. *Let M be a countable perfect group of infinite exponent and let G_M be the group constructed above. Then $\Phi(G_M) = H$, a direct product of $\aleph_0$ copies of M .*

4. Proof of Theorem 1.1. If K is of infinite exponent the result follows by Lemma 3.3, so suppose K is of finite exponent.

Let N be a countable perfect group of infinite exponent and define $M = K \times N$, a countable perfect group of infinite exponent. Consider the group G_M constructed above. By Lemma 3.3 we have $\Phi(G_M) = H = \text{Dr}_{i \in \mathbf{Z}} M_i$, where $M_i \simeq M$ and $M_i^{z^j} = M_{i+j}$, for all $i, j \in \mathbf{Z}$.

We may write M_0 as a direct product $M_0 = K_0 \times N_0$, where $K_0 \simeq K$ and $N_0 \simeq N$. Let $K_i = K_0^{z^i}$, $N_i = N_0^{z^i}$, and notice that $M_i = K_i \times N_i$, $N_i^z = N_{i+1}$, and $N_i^w = N_i$, for all $i \in \mathbf{Z}$. Lemma 3.1 shows that $G_M = \langle w, z \rangle$ and it follows that $D = \text{Dr}_{i \in \mathbf{Z}} N_i$ is a normal subgroup of G_M .

Clearly $H/D \simeq \text{Dr}_{i \in \mathbf{Z}} K_i$ and the result follows on setting $G_K = G_M/D$.

ACKNOWLEDGMENTS. The results presented here are contained in Chapter V of my Ph.D. thesis. I would like to record my gratitude to my supervisor, James Wiegold, who gave me much help and encouragement during my time as a research student. I am also indebted to the University of Wales for providing me with financial support.

REFERENCES

1. P. Hall *The Frattini subgroups of finitely generated groups*, Proc. London Math. Soc. (3) **11** (1961), 327–352.
2. R. C. Lyndon and P. E. Schupp, *Combinatorial group theory*, Springer-Verlag, Berlin-Heidelberg-New York, 1977.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ALABAMA, UNIVERSITY, ALABAMA
35486