RESTRICTED LEFT INVERTIBLE TOEPLITZ OPERATORS ON MULTIPLY CONNECTED DOMAINS

KEIJI IZUCHI AND SHŪICHI OHNO

Dedicated to Professor Junzo Wada on his sixtieth birthday

ABSTRACT. A characterization of restricted left invertible Toeplitz operators on multiply connected domains is given. To prove this, some extension theorems are given.

1. Introduction. Let Ω be a bounded connected open subset of the plane whose boundary Γ consists of finitely many disjoint analytic Jordan curves. We denote by $H^\infty(\Omega)$ the space of bounded holomorphic functions on Ω . In [1], Abrahamse studied Toeplitz operators on Ω and gave a left invertibility criterion for Toeplitz operators in a generalized sense. In [3], Clancey and Gosselin studied local properties of Toeplitz operators on the unit open disk and gave a characterization of restricted left invertible Toeplitz operators with the help of Younis' extension theorem [7].

In this paper, we will extend the Clancey-Gosselin-Younis theorem to multiply connected domains (§3). To prove this, we need some extension theorems as in [7, 8]. Since $H^\infty(\Omega)$ is not a strongly logmodular algebra, here we cannot use the Younis extension theorems. We will give more general extension theorems (§2).

2. Extension theorems. Let A be a function algebra on X . Throughout this section, we assume that X is a totally disconnected compact Hausdorff space. A closed subset E of X will be called a peak set for A if there is a function f in A , which is called a peaking function for E , such that $f = 1$ on E and $|f| < 1$ on $X \setminus E$. A subset S of X is called a weak peak set for A if it is an intersection of some peak sets. We consider the following separation condition.

(#) For any open and closed subset U of X , there is a function G in A such that $|G| > 1$ on U and $|G| < 1$ on $X \setminus U$.

The following theorem is a generalization of [8, Theorem 3.2 and Corollary 3.3].

THEOREM 1. *Suppose that A satisfies separation condition (#). Let S be a weak peak set for A . If a function f in A satisfies $|f| > 0$ on S , then there exists a function F in A such that $F = f$ on S and $|F| > 0$ on X .*

PROOF. We may assume that f vanishes somewhere on X . Take an open and closed subset U of X such that $S \subset U$, $|f| > 0$ on U , and

$$(1) \quad |f| \leq 1 \quad \text{on } X \setminus U.$$

Received by the editors December 9, 1985 and, in revised form, March 24, 1986.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 30H05, 46J10, 47B35.

Key words and phrases. Restricted left invertible, a family of Toeplitz operators, function algebras.

Put $c = \inf\{|f(x)|; x \in U\}$. Then $c > 0$. Since S is a weak peak set, there is a peak set E for A with $S \subset E \subset U$ [5, p. 56]. Let $h \in A$ be a peaking function for E . Replacing h by high powers of h , we may assume

$$(2) \quad |h| \leq 1/2 \quad \text{on } X \setminus U.$$

By our separation condition, there is a function G in A such that

$$(3) \quad |G| \leq c/3 \quad \text{on } U \quad \text{and} \quad |G| \geq 3 \quad \text{on } X \setminus U.$$

Put $F = f + G(1 - h)$. Clearly $F \in A$ and $F = f$ on S . For $x \in U$, by (3) we have

$$|F(x)| \geq |f(x)| - |G(x)| |1 - h(x)| \geq c/3 > 0.$$

For $x \in X \setminus U$, by (1), (2), and (3) we have

$$|F(x)| \geq |G(x)| |1 - h(x)| - |f(x)| \geq 1/2 > 0.$$

Thus $|F| > 0$ on X .

REMARK 1. Let A be a strongly logmodular algebra on a totally disconnected compact space X , that is, $\log |A^{-1}| = C_R(X)$, where A^{-1} denotes the set of invertible elements in A and $C_R(X)$ denotes the space of real continuous functions on X . Then A satisfies (#).

REMARK 2. For a general function algebra A , the assertion of Theorem 1 is not true. Put $X = \{(z, t); z \text{ is a complex number and } t \text{ is a real number with } |z| = 1 - |t| \text{ and } |t| \leq 1\}$. Let $A = \{g \in C(X); \text{ for each } t \text{ with } |t| < 1, g(z, t) \text{ has a continuous holomorphic extension to } \{|z| \leq 1 - |t|\}\}$. Then $S = \{(z, 0); |z| = 1\}$ is a peak set for A . Put $f(z, t) = (1 - |t|)z$. Then $f \in A$ and $|f| > 0$ on S , but it is easy to see that there is no $F \in A$ with $F = f$ on S and $|F| > 0$ on X .

For a weak peak set S , put $A_S = \{f \in C(X); f|_S \in A|_S\}$. Then A_S is a closed subalgebra of $C(X)$. For $\psi \in C(X)$ and $B \subset C(X)$, put $d(\psi, B) = \inf\{\|\psi - f\|; f \in B\}$. For $\psi \in C(X)$ and $F \subset X$, put $\|\psi\|_F = \sup\{|\psi(x)|; x \in F\}$. The following theorem is a generalization of [8, Theorem 3.1].

THEOREM 2. Suppose that A satisfies separation condition (#). Let S be a weak peak set for A , and let u be a function in $C(X)$ such that $\|u\| = 1$, $|u| = 1$ on S , and $d(u, A_S) < 1$. Then there is a function $\tilde{u}$ in $C(X)$ such that $|\tilde{u}| = |u|$ on X , $\tilde{u} = u$ on S , and $d(\tilde{u}, A) < 1$.

PROOF. Since $d(u, A_S) < 1$, there exists a function f in A such that $\|u - f\|_S < 1$. Since $|u| = 1$ on S , $|f| > 0$ on S . By Theorem 1, there is a function F in A such that $F = f$ on S , and

$$(1) \quad |F| > 0 \quad \text{on } X.$$

Since $\|u - F\|_S = \|u - f\|_S < 1$, there is an open and closed subset U of X such that $S \subset U$ and

$$(2) \quad \|u - F\|_U < 1.$$

Since S is a weak peak set, there is a peak set E for A such that $S \subset E \subset U$. Let $h \in A$ be a peaking function for E . Replacing h by high powers of $(1 + h)/2$, we may assume that

$$(3) \quad |h| > 0 \quad \text{on } X \quad \text{and} \quad 0 < |hF| < 1 \quad \text{on } X \setminus U.$$

Put

$$\tilde{u} = \begin{cases} uh/|h| & \text{on } U, \\ |u| hF/|hF| & \text{on } X \setminus U. \end{cases}$$

By (3), $\tilde{u} \in C(X)$. Clearly $|\tilde{u}| = |u|$ on X and $\tilde{u} = u$ on S . To prove $d(\tilde{u}, A) < 1$, it is sufficient to prove $\|\tilde{u} - hF\| < 1$, because $hF \in A$. We have

$$\begin{aligned} \|\tilde{u} - hF\|_{X \setminus U} &= \left\| \frac{|u| hF}{|hF|} - \frac{hF |hF|}{|hF|} \right\|_{X \setminus U} \\ &= \| |u| - |hF| \|_{X \setminus U} \\ &< 1 \quad \text{by (3) and } \|u\| \leq 1. \end{aligned}$$

Also we have

$$\|\tilde{u} - hF\|_U = \left\| \frac{uh}{|h|} - \frac{hF|h|}{|h|} \right\|_U = \|u - F|h|\|_U.$$

Let $x \in U$. Then

$$\begin{aligned} |u(x) - F(x)|h(x)| &= |(1 - |h(x)|)u(x) + |h(x)|(u(x) - F(x))| \\ &\leq 1 - |h(x)| + |h(x)| \|u - F\|_U \quad \text{by } \|u\| \leq 1, \|h\| \leq 1 \\ &= 1 - |h(x)|\{1 - \|u - F\|_U\} \\ &< 1 \quad \text{by (2)}. \end{aligned}$$

Thus we get $\|\tilde{u} - hF\|_U < 1$, hence $\|\tilde{u} - hF\| < 1$.

If A is a strongly logmodular algebra, then A_S , where S is a weak peak set for A , is generated by A and $\{f^{-1}; f \in A \cap A_S^{-1}\}$ [2]. Younis used this property to prove his theorem [8, Theorem 3.1]. So the following theorem is another generalization of [8, Theorem 3.1].

THEOREM 3. *Suppose that A satisfies separation condition (#). Let S be a weak peak set for A such that A_S is generated by A and $\{f^{-1}; f \in A \cap A_S^{-1}\}$. If u is a function in $C(X)$ with $\|u\| \leq 1$ and $d(u, A_S) < 1$, then there exists a function $\tilde{u}$ in $C(X)$ such that $|\tilde{u}| = |u|$ on X , $\tilde{u} = u$ on S , and $d(\tilde{u}, A) < 1$.*

NOTE. We do not assume $|u| = 1$ on S .

PROOF. By our assumption, there exist functions g in A and f in $A \cap A_S^{-1}$ such that

$$(1) \quad \|u - f^{-1}g\| < 1.$$

Since $f^{-1} \in A_S$, $|f^{-1}| > 0$ on X . By Theorem 1, there is a function F in A such that $F = f^{-1}$ on S and $|F| > 0$ on X . Put $\tilde{u} = ufF/|fF|$. Since $|fF| > 0$ on X , $\tilde{u} \in C(X)$. Clearly $|\tilde{u}| = |u|$ on X and $\tilde{u} = u$ on S . To prove $d(\tilde{u}, A) < 1$, suppose not. Since $\|\tilde{u}\| = \|u\| \leq 1$, we have $d(\tilde{u}, A) = 1$. We note that the space of bounded linear functionals of $C(X)/A$ may be identified with $A^\perp$, the set of regular Borel measures μ on X such that $\int_X \phi d\mu = 0$ for every $\phi \in A$. Hence there exists $\mu \in A^\perp$ with the unit total variation, $\|\mu\| = 1$, such that

$$1 = \int_X \tilde{u} d\mu = \int_X \frac{ufF}{|fF|} d\mu.$$

Since $\|u\| \leq 1$, we get $((ufF)/|fF|)\mu = |\mu|$, hence

$$(2) \quad ufF\mu = |fF| |\mu|.$$

Then

$$\begin{aligned}
 1 &> \left| \int_X (u - f^{-1}g) \frac{fF}{\|fF\mu\|} d\mu \right| && \text{by (1)} \\
 &= \frac{1}{\|fF\mu\|} \left| \int_X u fF d\mu \right| && \text{by } \mu \in A^\perp \\
 &= \frac{1}{\|fF\mu\|} \int_X |fF| d|\mu| && \text{by (2)} \\
 &= 1.
 \end{aligned}$$

This contradiction shows $d(\tilde{u}, A) < 1$.

3. Restricted left invertible Toeplitz operators. Let Ω be a bounded connected open subset of the plane whose boundary Γ consists of finitely many disjoint analytic Jordan curves. Identifying a function in $H^\infty(\Omega)$ with its boundary function, we may regard $H^\infty(\Omega)$ as an essentially supremum norm closed subalgebra of $L^\infty(m)$, where m is the arc length measure on Γ . A closed subspace $\mathcal{M}$ of $L^2(m)$ with $\mathcal{M} \neq \{0\}$ is called invariant if $H^\infty(\Omega)\mathcal{M} \subset \mathcal{M}$, and it is called reducing if $H^\infty(\Omega)\mathcal{M} \subset \mathcal{M}$ and $\bar{H}^\infty(\bar{\Omega})\mathcal{M} \subset \mathcal{M}$. An invariant subspace of $L^2(m)$ is called simple if it contains no reducing subspaces. For a closed subspace $\mathcal{M}$, $P_{\mathcal{M}}$ denotes the orthogonal projection of $L^2(m)$ onto $\mathcal{M}$. For a function ψ in $L^\infty(m)$, put $T_\psi^{\mathcal{M}}(f) = P_{\mathcal{M}}(\psi f)$ for every f in $\mathcal{M}$. $\tau_\psi = \{T_\psi^{\mathcal{M}}; \mathcal{M} \text{ is a simply invariant subspace}\}$ is called a family of Toeplitz operators with a symbol ψ . In the case that Ω is the unit open disk, by Beurling's theorem, $T_\psi^{\mathcal{M}}$ is unitarily equivalent to the usual Toeplitz operator T_ψ . We call τ_ψ left invertible if $T_\psi^{\mathcal{M}}$ is a left invertible operator on $\mathcal{M}$ for every simply invariant subspace $\mathcal{M}$. In [1, Theorem 4.1], Abrahamse proved the following theorem which is a generalization of the Devinatz-Rabinathan theorem (see [6, p. 119]).

THEOREM 4. *Let ψ be a function in $L^\infty(m)$ with $|\psi| = 1$ a.e. dm . Then the following conditions are equivalent:*

- (i) τ_ψ is left invertible.
- (ii) $d(\psi, H^\infty(\Omega)) < 1$.

We note that by his proof, (i) $\Rightarrow$ (ii) is true for every $\psi \in L^\infty(m)$ with $\|\psi\| \leq 1$.

Let X be the maximal ideal space of $L^\infty(m)$. Then X is a totally disconnected compact Hausdorff space [4, p. 190]. Identifying a function in $L^\infty(m)$ with its Gelfand transform, we have $L^\infty(m) = C(X)$. We may consider $H^\infty(\Omega)$ as a function algebra on X [4, p. 123]. Let S be a weak peak subset of X for $H^\infty(\Omega)$. A family of Toeplitz operators τ_ψ , $\psi \in C(X)$, is called S -restricted left invertible if there is a function Ψ in $C(X)$ such that $\Psi = \psi$ on S and τ_Ψ is left invertible. In the case that Ω is the open unit disk, Clancey-Gosselin-Younis [3, 7] gave a characterization of S -restricted left invertible Toeplitz operators as follows: If $|\psi| = 1$ a.e. dm , then the Toeplitz operator T_ψ is S -restricted left invertible if and only if $d(\psi, H_S^\infty) < 1$. We shall give a generalization of the above theorem to multiply connected domains as an application of §2.

THEOREM 5. *Let S be a weak peak subset of X for $H^\infty(\Omega)$. Let ψ be a function in $C(X)$ with $|\psi| = 1$ on S . Then the following conditions are equivalent:*

- (i) $\mathcal{T}_\psi$ is S -restricted left invertible.
- (ii) $d(\psi, H^\infty(\Omega)_S) < 1$.

PROOF. (ii) $\Rightarrow$ (i) Since X is totally disconnected, we may assume that $|\psi| = 1$ on X . We note that $A = H^\infty(\Omega)$ satisfies separation condition (#) in §2 [5, p. 119]. Then $H^\infty(\Omega)$, ψ , and S satisfy all assumption of Theorem 2. Hence there exists a function Ψ in $C(X)$ such that $|\Psi| = 1$ on X , $\Psi = \psi$ on S , and $d(\Psi, H^\infty(\Omega)) < 1$. By Theorem 4, $\mathcal{T}_\Psi$ is left invertible. So (i) holds.

(i) $\Rightarrow$ (ii) Suppose that $\mathcal{T}_\psi$ is S -restricted left invertible. By our definition, there is a function Ψ in $C(X)$ such that $\Psi = \psi$ on S and $\mathcal{T}_\Psi$ is left invertible. First we shall show that there is a function h in $H^\infty(\Omega)$ such that

$$(1) \quad h = 1 \text{ on } S, \quad \|\Psi h\| \leq 1, \quad \text{and} \quad |h| > 0 \text{ on } X.$$

To prove this, put

$$(2) \quad \phi = \max\{|\Psi|, 1\}.$$

Then $\phi \in C(X)$, $\phi \geq 1$ on X , and $\phi = 1$ on S . By [5, p. 58], there is a function g in $H^\infty(\Omega)$ such that $g = 1$ on S and

$$(3) \quad |g| \leq 1/\phi \text{ on } X.$$

Then $\|g\| = 1$. Take a positive integer N with

$$(4) \quad N \geq 4 \quad \text{and} \quad \|\Psi\|(5/6)^N \leq 1.$$

Put $h = ((g + 2)/3)^N$. Then $h \in H^\infty(\Omega)$, $h = 1$ on S , and $|h| > 0$ on X . If $x \in X$ with $|g(x)| \leq 1/2$, then

$$|\Psi(x)h(x)| \leq \|\Psi\|(5/6)^N \leq 1 \quad \text{by (4).}$$

If $x \in X$ with $1/2 \leq |g(x)| \leq 1$, then

$$\begin{aligned} |\Psi(x)h(x)| &\leq ((|g(x)| + 2)/3)^N |\phi(x)| \quad \text{by (2)} \\ &\leq ((|g(x)| + 2)/3)^N / |g(x)| \quad \text{by (3)} \\ &\leq 1. \end{aligned}$$

The last inequality follows from $((t + 2)/3)^N \leq t$ for $1/2 \leq t \leq 1$ and $N \geq 4$. Thus $\|\Psi h\| \leq 1$ and we get (1).

We put $\Phi = \Psi h \in L^\infty(m)$. By (1),

$$(5) \quad \|\Phi\| \leq 1 \quad \text{and} \quad \Phi = \psi \text{ on } S.$$

Also we have that $\mathcal{T}_\Phi$ is left invertible. To see this, let $\mathcal{M}$ be a simply invariant subspace of $L^2(m)$. Since $h \in H^\infty(\Omega)$ and $|h| > 0$ on X , $T_h^\mathcal{M}$ is left invertible. Since $\mathcal{T}_\Psi$ is left invertible, $T_\Phi^\mathcal{M} = T_\Psi^\mathcal{M} T_h^\mathcal{M}$ is left invertible, so $\mathcal{T}_\Phi$ is left invertible. As noted after Theorem 4, we get $d(\Phi, H^\infty(\Omega)) < \|\Phi\| \leq 1$. Hence

$$\begin{aligned} d(\psi, H^\infty(\Omega)_S) &= d(\Phi, H^\infty(\Omega)_S) \quad \text{by (5)} \\ &\leq d(\Phi, H^\infty(\Omega)) < 1. \end{aligned}$$

This completes the proof.

REFERENCES

1. M. B. Abrahamse, *Toeplitz operators in multiply connected regions*, Amer. J. Math. **96** (1974), 261–297.
2. S. Axler, *Subalgebras of L^∞* , Thesis, University of California, Berkeley, 1975.
3. K. Clancey and J. Gosselin, *On the local theory of Toeplitz operators*, Illinois J. Math. **22** (1978), 449–458.
4. S. Fisher, *Function theory on plane domains*, Wiley, New York, 1983.
5. T. W. Gamelin, *Uniform algebras*, Prentice-Hall, Englewood Cliffs, N.J., 1969.
6. D. Sarason, *Function theory on the unit circle*, Lecture Notes, Virginia Polytech Inst. and State Univ., Blacksburg, Virginia, 1978.
7. R. Younis, *Extension results in the Hardy space associated with a logmodular algebra*, J. Funct. Anal. **39** (1980), 16–22.
8. ———, *Interpolation in strongly logmodular algebras*, Pacific J. Math. **102** (1982), 247–251.

DEPARTMENT OF MATHEMATICS, KANAGAWA UNIVERSITY, YOKOHAMA 221, JAPAN