ORIENTATION PRESERVING ACTIONS OF FINITE ABELIAN GROUPS ON SPHERES

RONALD M. DOTZEL

ABSTRACT. If G is a finite Abelian group acting as a $\mathbf{Z}_{(\mathcal{P})}$ -homology n -sphere X (where $\mathcal{P}$ is the set of primes dividing $|G|$), then there is an integer valued function $n(\cdot, G)$ defined on the prime power subgroups H of G such that X^H has the $\mathbf{Z}_{(p)}$ -homology of a sphere $S^{n(H,G)}$. We prove here that there exists a real representation R of G such that for any prime power subgroup H of G , $\dim(S(R^H)) = n(H, G)$ where $S(R^H)$ is the unit sphere of R^H , provided that $n - n(H, G)$ is even whenever H is a 2-subgroup of G .

0. Introduction. Suppose that G is a finite Abelian group and let $\mathcal{P}$ be the set of primes dividing $|G|$. If G acts on a finite CW-complex X which has the $\mathbf{Z}_{(\mathcal{P})}$ -homology of S^n , then for any $p \in \mathcal{P}$ and p -subgroup H of G the fixed point set of H on X , X^H , has the $\mathbf{Z}_{(p)}$ -homology of $S^{n(H,G)}$ for some integer $n(H, G) \geq -1$ (-1 signifies empty). This is a well-known consequence of Smith theory ([2, III or 1, IV], e.g.). Thus we obtain in this way an integer valued function, $n(\cdot, G)$, defined on the set of p -subgroups of G by $H \mapsto n(H, G)$ (note that $n(e, G) = n$). This function is called the “dimension function” and has a considerable literature (see [5, 3, 6, 7, 9]; [10] gives a related extensive bibliography).

The function $n(\cdot, G)$ satisfies the following well-known conditions (see [1, XIII, 2.3; IV, 4.4, 4.7]):

1. (Borel Formula) If $H \leq K$ are both p -subgroups of G and $K/H = \mathbf{Z}_p + \mathbf{Z}_p$, then $n(H, G) - n(K, G) = \sum (n(K', G) - n(K, G))$ with the sum over all $H \leq K' \leq K$ such that $K'/H = \mathbf{Z}_p$.

2. If $H \leq K$ are p -subgroups of G , then $n(K, G) \leq n(H, G)$.

3. If $H \leq K$ are p -subgroups of G with $K/H = \mathbf{Z}_p$ and p odd, then $n(H, G) - n(K, G)$ is even.

4. If $H \leq K' \leq K$ are 2-subgroups of G such that $K/H = \mathbf{Z}_4$, $K'/H = \mathbf{Z}_2$, then $n(H, G) - n(K', G)$ is even.

For each $p \in \mathcal{P}$, let $G(p)$ denote the p -Sylow subgroup of G and set $N(\cdot, G) = n(\cdot, G) + 1$. The function $N(\cdot, G)$ restricted to the subgroups of $G(p)$ will naturally be denoted by $N(\cdot, G(p))$. In [8] it was shown that $N(\cdot, G(p))$ is realized by a real representation $V(p)$ of $G(p)$ which means that for each $H \leq G(p)$, $\dim V(p)^H = N(H, G(p)) = N(H, G)$. If $S(V(p))$ denotes the unit sphere of $V(p)$, then $\dim S(V(p)^H) = n(H, G)$.

Here we are interested in the existence of a real representation R of the Abelian group G such that for any p -subgroup H of G (for any $p \in \mathcal{P}$), $\dim R^H = N(H, G)$. Thus R would be a simultaneous realization of the functions $N(\cdot, G(p))$, $p \in \mathcal{P}$.

Received by the editors March 22, 1985. These results were presented to the AMS at the 829th meeting in Charlotte, North Carolina, in October 1986.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 57S17, 57S25.

It should be noted that in the special case where X^H is a homology sphere for all $H \leq G$, then $N(\cdot, G)$ is defined on all subgroups of G and tom Dieck has shown in [3] that $N(\cdot, G)$ is realized by a difference of representations. In general, this is best possible.

Clearly $N(\cdot, G)$ satisfies conditions 1-4 if $n(\cdot, G)$ does. We will denote $N(e, G)$ by N . As an example to show that some condition beyond 1-4 is needed, suppose $G = \mathbf{Z}_6$, $N = 2$, $N(\mathbf{Z}_3, G) = 0$, and $N(\mathbf{Z}_2, G) = 1$. Then $N(\cdot, G)$ satisfies conditions 1-4 but there is no real representation of G which realizes these numbers simultaneously as dimensions. From now on we will assume the following "orientation preserving" condition holds, in addition to conditions 1-4:

5. If H is any 2-subgroup of G , then $N - N(H, G)$ is even.

We obtain the following theorem and corollary:

THEOREM. *Let G be a finite Abelian group and suppose $N(\cdot, G)$ is a nonnegative integer valued function defined on the p -subgroups of G for all $p \mid |G|$, satisfying conditions 1-5. Then there exists a real representation R of G such that for any p -subgroup H of G , $p \mid |G|$, $\dim R^H = N(H, G)$. Furthermore, if $\bar{R}$ is another such representation of G then for all subgroups H of G ,*

$$\dim R^H \equiv \dim \bar{R}^H \pmod{2}.$$

COROLLARY. *Let G be a finite Abelian group and suppose the 2-Sylow subgroup of G is cyclic. If $N(\cdot, G)$ is a nonnegative integer valued function defined on the p -subgroups of G , $p \mid |G|$ satisfying only conditions 1-4, then there exists a real representation R such that for any p -subgroup H of G , $p \mid |G|$, $\dim R^H = N(H, G) + 1$.*

In §§1 and 2 we prove the theorem and corollary respectively. We thank the referee for several suggestions leading to an improved exposition.

1. Proof of the theorem. Let G be a finite Abelian group and suppose $N(\cdot, G)$ is a nonnegative integer valued function on the p -subgroups of G , for all $p \in \mathcal{P}$, satisfying conditions 1-5. By [8], for each $p \in \mathcal{P}$ there is a representation $V(p)$ of the p -Sylow subgroup $G(p)$ of G such that $\dim V(p)^H = N(H, G(p))$ for all $H \leq G(p)$. Let $\mathcal{V} = \bigotimes_{p \mid |G|} V(p)$. Then $\mathcal{V}$ is a representation of G and we will prove by induction on $|G|$ that $\mathcal{V}$ contains a subrepresentation R of G of dimension $N = N(e, G)$ such that $\dim R^H = N(H, G)$ for all p -subgroups H of G , all $p \in \mathcal{P}$. So if $|K| < |G|$, $N(\cdot, K)$ is a nonnegative integer valued function on the prime power subgroups of K and $W(p)$ is a representation of $K(p)$ realizing $N(\cdot, K(p))$, we can assume $\mathcal{W} = \bigotimes_{p \mid |K|} W(p)$ contains a subrepresentation realizing $N(\cdot, K)$.

Suppose that $N(\cdot, G)$ is a nonnegative integer valued function defined on the prime power order subgroups of an Abelian group G satisfying conditions 1-5 and suppose $N(e, G) = N(H, G)$ for some $H \leq G(p)$, $|H| = p$. Then for any prime power order subgroup K of G , define $N(K/K \cap H, G/H) = N(K, G)$. It is clear that $N(\cdot, G/H)$ satisfies conditions 1-5. Moreover, for any $K \leq G(p)$,

$$N(K, G) \stackrel{\text{def}}{=} N(K, G(p)) = N(KH, G(p)) \stackrel{\text{def}}{=} N(KH, G) \stackrel{\text{def}}{=} N(KH/H, G/H).$$

For by induction we can assume $N(K', G(p)) = N(K'H, G(p))$ for any $K' \not\leq K$ and clearly we can assume $H \not\leq K$. Select $K' < K$ such that $|K/K'| = p$ and use

condition 1 (Borel Formula) on $K' \leq KH$ to obtain $N(K, G(p)) = N(KH, G(p))$. It follows that in this case, a representation of G/H realizing $N(\cdot, G/H)$ can be regarded as an unfaithful representation of G (with kernel at least H) realizing $N(\cdot, G)$.

Now for each $p \in \mathcal{P}$ and each $\mathbf{Z}_p \leq G$, we must have $N - N(\mathbf{Z}_p, G) > 0$, otherwise by the observation above we could assume we are given a dimension function on $G/\mathbf{Z}_p$. Of all the differences $N - N(\mathbf{Z}_p, G)$, $p \in \mathcal{P}$, let p_0 and $H_0 = \mathbf{Z}_{p_0}$ be such that $N - N(H_0, G)$ is a minimum. Then the representation $V(p_0)$ of $G(p_0)$ (the p_0 -Sylow subgroup of G) contains an irreducible subrepresentation $W(p_0)$ of $G(p_0)$ on which H_0 acts without (nonzero) fixed points. For $q \neq p_0$ select $H = \mathbf{Z}_q \leq G(q)$ such that $N - N(H, G)$ is least for q (in general $N - N(H_0, G) \leq N - N(H, G)$) and let $W(q)$ be an irreducible subrepresentation of $V(q)$ on which H acts without fixed points. Then $\mathcal{W} = \bigotimes_{q \mid |G|} W(q)$ is a G -subrepresentation of $\mathcal{V} = \bigotimes_{q \mid |G|} V(q)$.

Let R_1 be an irreducible G -subrepresentation of $\mathcal{W}$. If $|G|$ is larger than 2, R_1 has dimension 2, since R_1 induces a free, irreducible, real representation of a cyclic group of order larger than 2 (the cyclic group is $G/\text{kernel of } R_1 = \text{kernel of } \mathcal{W}$).

Now R_1 , being a representation of G , has associated to it a dimension function $N_1(\cdot, G)$ defined on all subgroups H of G by $N_1(H, G) = \dim R_1^H$. Set $\bar{N}_1(\cdot, G) = N(\cdot, G) - N_1(\cdot, G)$. It is easy to verify that $\bar{N}_1(\cdot, G)$ is a dimension function defined on the prime power subgroups of G satisfying conditions 1-5.

Since $\bar{N}_1(e, G) < N(e, G)$ and $\bar{N}_1(H_0, G) = N(H_0, G)$ we are presented with two situations: (a) $\bar{N}_1(e, G) = \bar{N}_1(H_0, G)$ or (b) $\bar{N}_1(e, G) > \bar{N}_1(H_0, G)$.

In case (a) the function $\bar{N}_1(\cdot, G)$ may be replaced (as noted above) by a dimension function defined on the prime power subgroups of G/H_0 . Since for any q -subgroup K of G , $\dim R_1^K = \dim W(q)^K$, the subrepresentation $W(q)^\perp$ of $V(q)$ realizes the dimension function $\bar{N}_1(\cdot, G(q))$. Since $|G/H_0| < |G|$ by induction the tensor product of all the $W(q)^\perp$ contains a subrepresentation R^* of G/H_0 (which may be thought of as an unfaithful representation of G). R^* is a G -subrepresentation of $\mathcal{V}$, the tensor product of all the $V(q)$. $R^* \oplus R_1$ is the required representation in this case.

In (b), where we have $\bar{N}_1(e, G) > \bar{N}_1(H_0, G)$, note that $\bar{N}_1(e, G) - \bar{N}_1(H_0, G)$ is still a minimum of all differences $\bar{N}_1(e, G) - \bar{N}_1(H, G)$. Since the function $\bar{N}_1(\cdot, G(Q))$ is realized by the subrepresentation $W(q)^\perp$ of $V(q)$, we can repeat the procedure again obtaining another irreducible subrepresentation R_2 of G with an associated dimension function $N_2(\cdot, G)$ defined on the prime power subgroups of G (it is the restriction of a dimension function defined on all subgroups of G). Letting $\bar{N}_2(\cdot, G) = \bar{N}_1(\cdot, G) - N_2(\cdot, G)$ we again have a dimension function satisfying conditions 1-5 and we proceed as above. Eventually we obtain a dimension function $\bar{N}_k(\cdot, G)$ such that $\bar{N}_k(e, G) = \bar{N}_k(H_0, G)$ ($k = N - N(H_0, G)$). By case (a) and induction there is a G -subrepresentation of $\mathcal{V}$, R^* realizing $\bar{N}_k(\cdot, G)$. The representation $R = R^* \oplus R_1 \oplus R_2 \oplus \cdots \oplus R_k$ is the required G -subrepresentation of $\mathcal{V}$.

Now suppose $\bar{R}$ is another G -subrepresentation such that for any prime power order subgroup H of G , $\dim \bar{R}^H = N(H, G)$. Let H be an arbitrary subgroup of G and by induction assume $\dim \bar{R}^K - \dim R^K$ is even for all subgroups K of G with $|K| < |H|$. Select $K \leq H$ so that $|H/K|$ is an odd prime p (if this is not possible

then H is a 2-group and $\dim \bar{R}^H - \dim R^H$ is zero). The group $H/K = \mathbf{Z}_p$ acts on both $\bar{R}^K$ and R^K . It follows that both $\dim \bar{R}^K - \dim \bar{R}^H$ and $\dim R^K - \dim R^H$ are even and therefore $\dim \bar{R}^H - \dim R^H$ is even. This completes the proof of the theorem. $\square$

2. Proof of the corollary. Suppose G is a finite Abelian group with cyclic 2-Sylow subgroup, $G(2)$, and suppose $N(, G)$ is a nonnegative integer valued function defined on the p -subgroups of G , $p \mid |G|$, satisfying conditions 1–4. By condition 4, for any proper subgroup H of $G(2)$, $N - N(H, G)$ is even. Suppose that $N - N(G(2), G)$ is odd. For each $p \mid |G|$, let $\bar{V}(p) = V(p) \oplus 1$, where 1 denotes the trivial one-dimensional representation of $G(p)$. It is easy to see that $\bar{V}(p)$ realizes $N(, G(p)) + 1$. The function $N^*(, G) = N(, G) + 1$ corresponds to the G -action on the unreduced suspension on X .

Now since $N - N(G(2), G)$ is odd, $\bar{V}(2)$ has an irreducible summand of dimension 1 on which H , the maximal proper subgroup of $G(2)$, acts trivially and on which $G(2)$ acts nontrivially. Denote this summand by $W(2)$ for any $p \neq 2$ let $W(p)$ be a one-dimensional trivial subrepresentation of $\bar{V}(p)$. Then $R_1 \otimes_{p \mid |G|} W(p)$ is a 1-dimensional G -representation with a very large kernel and is a subrepresentation of $\bar{V} = \bigotimes_{p \mid |G|} \bar{V}(p)$. Let $N_1(, G)$ be the dimension function associated with R_1 ($N_1(, G)$ is actually defined on all subgroups of G). Setting $\bar{N}(, G) = N^*(, G) - N_1(, G)$ we see that $N - \bar{N}(H, G)$ is now even for all prime power subgroups of G so $\bar{N}(, G)$ satisfies conditions 1–5. By the argument §1, $\bar{N}(, G)$ is realized by a subrepresentation R of $\bigotimes_{p \mid |G|} W(p)^\perp$, since $\bar{N}(, G(p))$ is realized by $W(p)^\perp$ for all $p \mid |G|$. Then $R \oplus R_1$ is a subrepresentation of the G -representation $\bar{V}$ which realizes $N^*(, G)$. This establishes the corollary.

EXAMPLES. Let $G = \mathbf{Z}_6$, $N = 2$, $N(\mathbf{Z}_2, G) = 1$, $N(\mathbf{Z}_3, G) = 0$. If we “suspend” $N(, G)$ we have $N^* = 3$, $N^*(\mathbf{Z}_2, G) = 2$, $N^*(\mathbf{Z}_3, G) = 1$. Then the construction of §§2 and 1 yields the 3-dimensional representation of G , given on a generator by

$$\begin{pmatrix} -1 & 0 & 0 \\ 0 & R(2\pi/3) & \\ 0 & & \end{pmatrix}$$

where $R(2\pi/3)$ is a 2×2 rotation matrix.

REFERENCES

1. A. Borel et al., *Seminar on transformation groups*, Ann. of Math. Studies, no. 46, Princeton Univ. Press, Princeton, N. J.,
2. G. Bredon, *Introduction to compact transformation groups*, Academic Press, New York,
3. T. tom Dieck, *Homotopiedarstellungen endlicher Gruppen: Dimensionsfunktionen*, Invent. Math. **67** (1982), 231–252.
4. —, *Transformation groups and representation theory*, Lecture Notes in Math., vol. 766, Springer-Verlag, 1979.
5. —, *Semi-linear group actions on spheres: Dimension functions*, Lecture Notes in Math., vol. 763, Springer-Verlag, 1978, pp. 448–457.
6. T. tom Dieck and T. Petrie, *Homotopy representation of finite groups*, Inst. Hautes Étude Sci. Publ. Math. **56** (1983), 129–169.
7. —, *Geometric modules over the Burnside ring*, Invent. Math. **47** (1978), 273–287.

8. R. Dotzel and G. Hamrick, *p-group actions on homology spheres*, Invent. Math. **62** (1981), 437–442.
9. K. H. Dovermann and T. Petrie, *Artin relations for smooth representations*, Proc. Nat. Acad. Sci. U.S.A. **77** (1980), 5620–5621.
10. R. Schultz, *Nonlinear analogs of linear group actions*, Bull. Amer. Math. Soc. (N.S.) **11** (1984), 263–285.

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF MISSOURI-ST. LOUIS,
ST. LOUIS, MISSOURI 63121