SOME HOMOTOPY PROPERTIES OF THE HOMEOMORPHISM GROUPS OF $R^\infty(Q^\infty)$ -MANIFOLDS

VO THANH LIEM

ABSTRACT. In this note we will prove that, given an $R^\infty(Q^\infty)$ -manifold M , there is a deformation of $\text{Homo}(M)$ into $\text{Homeo}(M)$ whose final stage is a weak homotopy equivalence, and that if M has the homotopy type of a finite simplicial complex, then $\text{Homeo}(M)$ is an $\text{ANE}(\mathcal{CW}(M))$ and an $\text{ANE}(\mathcal{CW}(C), G_\delta)$.

0. Introduction. All spaces are Hausdorff and maps are continuous. The composition of two maps $f: A \rightarrow B$ and $g: B \rightarrow C$ will be denoted by gf . Let $\mathcal{CG}$ denote the category whose objects are compactly generated spaces (k -spaces) and whose morphisms are continuous maps [Gr, 8.1]. Given two spaces X and Y , following [Gr], let $X \times Y$ denote the product of X and Y in $\mathcal{CG}$, let $X \times_c Y$ be the Cartesian product, let $C(X, Y)$ be the space of maps from X to Y with the compact-open topology, and let $Y^X = kC(X, Y)$ [Gr, 8.14]. Similarly, let $\text{Homeo}(X)$ ($\text{Homo}(X)$, resp.) denote the space of homeomorphisms (homotopy equivalences, resp.) of X with compact-open topology as a subspace of $C(X, X)$. Recall that $k\text{Homeo}(X)$ and $k\text{Homo}(X)$ are subspaces of X^X in the sense of [Hy, p. 201]. Following [L₃], let $\mathcal{CW}(C)$ and $\mathcal{CW}(M)$ denote the classes of pseudo CW-complexes generated by the class C of Hausdorff compact spaces and by the class M of metric spaces, respectively. Let $E^\infty = \text{dirlim } E^n$, where E^n is either the n -Euclidean space or the n -fold product of the Hilbert cube Q . By an E^∞ -manifold, we mean a separable paracompact space which is locally homeomorphic to E^∞ .

In this note we will prove that given an E^∞ -manifold M , there is a deformation of $\text{Homo}(M)$ into $\text{Homeo}(M)$ whose final stage is a weak homotopy equivalence (this partially answers a question raised by P. J. Kahn some time ago [L₃, §0]), and that if M has the homotopy type of a finite complex, then $\text{Homeo}(M)$ is an absolute neighborhood extensor [Hu] (abbreviated ANE) for the class $\mathcal{CW}(M)$ and is an $\text{ANE}(\mathcal{CW}(C), G_\delta)$. By a space $S \in \text{ANE}(\mathcal{X}, G_\delta)$, we mean that if $X \in \mathcal{X}$ and A is a closed G_δ -subset of X , then every map $f: A \rightarrow S$ has a continuous extension over a neighborhood of A in X .

Throughout this note $I = [0, 1]$. By an E^∞ -manifold fiber bundle, we mean a locally trivial bundle in $\mathcal{CG}$, $p: X \rightarrow B$, whose fiber is an E^∞ -manifold. If (X, p, B) is a bundle, W a subset of X , and A a subset of B , we write $W_A = W \cap p^{-1}(A)$; specially, $X_A = p^{-1}(A)$. Let (X, p, B) and (Y, p', B) be bundles and W a subset of X . A map $f: W \rightarrow Y$ is said to be *fiber preserving* (f.p.) if $p'f = p|_W$. If $f: W \rightarrow B \times Z$ is an f.p. map, for each $b \in B$ let $f_b: W_{\{b\}} \rightarrow Z$ denote the map $p_Z f|_{W_{\{b\}}}$ where $p_Z: B \times Z \rightarrow Z$ is the projection. Given an $R^\infty(Q^\infty)$ -manifold M ,

Received by the editors January 30, 1986.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 54C20, 54C55.

Key words and phrases. Homotopy, k -space, evaluation map, fiber-preserving map.

following $[\mathbf{L}_1, \mathbf{L}_2]$, we write $M = \text{dirlim } M_n$, where M_n is a compact submanifold (compact Q -submanifold) of M_{n+1} for each $n = 1, 2, \dots$.

Given a map $f: B \rightarrow Y^X$ or $f: B \rightarrow C(X, Y)$, by the *associate* of f we mean the f.p. function $\hat{f}: B \times Y \rightarrow B \times Y$ defined by $\hat{f}(b, x) = (b, f(b)(x))$, and vice versa; we also call f the associate of $\hat{f}$ [$\mathbf{D}$, p. 261]. By [$\mathbf{Gr}$, 8.17], the continuity of f and that of its associate are equivalent if $B, X, Y \in \mathcal{CG}$. Given a subset A of X , let i_A denote the inclusion $A \rightarrow X$ and id_A the identity map on A .

1. A canonical deformation. The main result of this section is Theorem 1.3 which proves the existence of a deformation of the space of homotopy equivalences into the space of homeomorphisms. The proof of the following lemma is straightforward.

LEMMA 1.1. *Let X, Y , and Z be topological spaces. Let*

$$F: C(X, Y) \rightarrow C(X \times_c Z, Y \times_c Z)$$

be the map defined by $F(g) = g \times \text{id}_Z$. Then F is an embedding; i.e., $C(X, Y)$ is homeomorphic to the subspace $F(C(X, Y))$ of $C(X \times_c Z, Y \times_c Z)$. $\square$

We need another lemma for the proof of Theorem 1.3. Recall that $X \times_c I \cong X \times I$ for $X \in \mathcal{CG}$ [$\mathbf{Gr}$, 8.11].

LEMMA 1.2. *Let M, N be topological spaces. If $g: M \times_c I \rightarrow M$ and $h: N \times_c I \rightarrow N$ are given homotopies, then g and h induce the homotopies G and H on $C(M, N)$ defined by $G(f, t) = fg_t$ and $H(f, t) = h_t f$ for all $f \in C(M, N)$ and $t \in I$.*

PROOF. For the continuity of G , consider the following composition:

$$\bar{g}: C(M, N) \xrightarrow{\tilde{p}_M} C(M \times_c I, N) \xrightarrow{\tilde{g}'} C(M \times_c I, N) \xrightarrow{\sigma} C(I, C(M, N)),$$

where $\tilde{p}_M, \tilde{g}'$ are maps naturally induced [$\mathbf{D}$, p. 259] from the projections $p_M: M \times_c I \rightarrow M$ and $g': M \times_c I \rightarrow M \times_c I$ defined by $g'(x, t) = (g(x, t), t)$, and where σ is the embedding defined by $\sigma(f)(t)(x) = f(x, t)$ for all $t \in I$ and $x \in M$ [$\mathbf{J}$, Theorem 1.1]. Define

$$G: C(M, N) \times_c I \xrightarrow{\bar{g} \times \text{id}_I} C(I, C(M, N)) \times_c I \xrightarrow{e} C(M, N),$$

where e is the evaluation map. Then G is a homotopy. Let $f \in C(M, N)$ and $t \in I$. We have

$$\begin{aligned} G(f, t) &= e(\bar{g} \times \text{id}_I)(f, t) = e((\sigma \tilde{g}' \tilde{p}_M) \times \text{id}_I)(f, t) \\ &= e(\sigma \tilde{g}' \tilde{p}_M(f), t) = e(\sigma(fp_M g'), t) = \sigma(fp_M g')(t) \\ &= fp_M g'(\cdot, t) = fg(\cdot, t) = fg_t. \end{aligned}$$

For the continuity of H , consider the composition

$$\bar{h}: C(M, N) \xrightarrow{F} C(M \times_c I, N \times_c I) \xrightarrow{\tilde{h}} C(M \times_c I, N) \xrightarrow{\sigma} C(I, C(M, N)),$$

where the embedding F is given by Lemma 1.1, $\tilde{h}$ is induced from h and σ is from [$\mathbf{J}$, Theorem 1.1] as above. Define

$$H: C(M, N) \times_c I \xrightarrow{\bar{h} \times \text{id}_I} C(I, C(M, N)) \times_c I \xrightarrow{e} C(M, N).$$

Then H is continuous. Finally, let $f \in C(M, N)$ and $t \in I$. We have

$$\begin{aligned} H(f, t) &= e(\bar{h} \times \text{id}_I)(f, t) = e(\bar{h}(f), t) = \bar{h}(f)(t) \\ &= (\sigma \tilde{h} F(f))(t) = \tilde{h}(F(f))(\cdot, t) \\ &= (\tilde{h}(f \times \text{id}_I))(\cdot, t) = h((f \times \text{id}_I)(\cdot, t)) \\ &= h(f(\cdot), t) = h_t f. \quad \square \end{aligned}$$

REMARK 1. In Lemma 1.2, if M and N are k -spaces, then G and H are also homotopies on $N^M = kC(M, N)$ (by use of [Gr, 8.8(vi)]).

THEOREM 1.3. *Let M be an E^∞ -manifold. Then, there is a deformation of $\text{Homo}(M)$ in itself into $\text{Homeo}(M)$ whose final map $r: \text{Homo}(M) \rightarrow \text{Homeo}(M)$ is a weak homotopy equivalence.*

PROOF. To avoid confusion, we let N be another copy of M and consider $\text{Homo}(M, N)$ and $\text{Homeo}(M, N)$ instead of $\text{Homo}(M)$ and $\text{Homeo}(M)$, respectively. By [He, Theorem C], we can identify M and N with $K \times E^\infty$ and $L \times E^\infty$, respectively, where K and L are locally finite simplicial complexes. We also identify K and L with $K \times 0$ and $L \times 0$, respectively. Let $g: M \times I \rightarrow M$ and $h: N \times I \rightarrow N$ be natural strong deformation retractions of M and N onto K and L , respectively, with $g_1 = p_K$ and $h_1 = p_L$ (the projections). Let G and H be the homotopies on $C(M, N)$ induced by g and h given by Lemma 1.2. Define $D_t: C(M, N) \rightarrow C(M, N)$ by

$$D_t = \begin{cases} H_{4t} & \text{if } t \in [0, 1/4], \\ G_{4t-1} & \text{if } t \in [1/4, 1/2], \\ G_{4t-2}^* & \text{if } t \in [1/2, 3/4], \\ H_{4t-3}^* & \text{if } t \in [3/4, 1], \end{cases}$$

where $G_t^* = G_{1-t}$ and $H_t^* = H_{1-t}$. Then

$$D: f \simeq p_L f \simeq p_L f p_K \simeq p_L((p_L f|_K) \times \text{id}_{E^\infty}) \simeq (p_L f|_K) \times \text{id}_{E^\infty}, w_3$$

and

$$D_1(C(M, N)) = \{q \times \text{id}_{E^\infty} \mid q \in C(K, L)\} \cong C(K, L)$$

(by Lemma 1.1) which is a metrizable space [Ku, Theorem 1, p. 93].

Observe that $D(\text{Homo}(M, N) \times_c I) \subset \text{Homo}(M, N)$. So, D deforms $\text{Homo}(M, N)$ in itself into the metrizable subspace $B \equiv \text{Homo}(M, N) \cap \{q \times \text{id}_{E^\infty} \mid q \in C(K, L)\}$. Recall that since B is metrizable, it also is a subspace of $k\text{Homo}(M, N)$. Now, we would like to deform B into $\text{Homeo}(M, N)$. Consider the associate g of the inclusion $i_B: B \rightarrow \text{Homo}(M, N)$, i.e.,

$$g: B \times K \times E^\infty \rightarrow B \times L \times E^\infty$$

defined by $g(b, x, y) = (b, q(x), y)$, where $b = q \times \text{id}_{E^\infty}$. Then, g is continuous [Gr, 8.17], and g is an f.p. homotopy equivalence [Ja, Proposition 7.58]. Therefore, by [L3, Lemma 3.3], g is f.p. homotopic to an f.p. homeomorphism, say $\bar{g}$. This homotopy induces a homotopy $R: B \times I \rightarrow \text{Homo}(M, N)$ with $R_0 = i_B$ and $R_1(B) \subset \text{Homeo}(M, N)$. Combining D and R , we will obtain a desired deformation with a final map $r = R_1 D_1$.

Finally, let $i: \text{Homeo}(M) \rightarrow \text{Homo}(M)$ be the natural inclusion. Then we have $ir = r \simeq \text{id}_{\text{Homo}(M)}$. By [L3, Lemma 3.3], i is a weak homotopy equivalence (i.e., i

induces isomorphisms on homotopy groups). It follows easily that r is also a weak homotopy equivalence. $\square$

To conclude this section, we prove the following proposition which will be used in the next section. In fact, the proposition indicates that $\text{Homeo}(M)$ is not a closed subset of $\text{Homo}(M)$. Therefore, it is interesting to know whether the deformation in the above theorem can be chosen such that $\text{Homeo}(M)$ is deformed in itself.

PROPOSITION 1.4. *Let A be a closed subset (G_δ -subset, resp.) of $B \in \mathcal{CW}(M)$ ($B \in \mathcal{CW}(C)$, resp.), and (X, p, B) an E^∞ -manifold fiber bundle. If $f: X \rightarrow X$ is an f.p. homotopy equivalence with $f|_{X_A}$ a continuous injection, then there is an f.p. map $h: X \rightarrow X$ such that $h|_{X_{B-A}}: X_{B-A} \rightarrow X_{B-A}$ is an f.p. homeomorphism and that f is f.p. homotopic to h (rel X_A).*

PROOF. Similar to the proof of [L₃, Theorem 3.4], we will assume that $X = B \times M$, where M is an E^∞ -manifold. Since A is a closed G_δ -set in B , we can write $B - A = \bigcup \{C_n | n = 1, 2, \dots\}$, where each C_n is a closed G_δ -set in B and $C_n \subset$ (interior of C_{n+1}). Write $M = \text{dirlim } M_n$ and assume $C_0 = \emptyset$ and $M_0 = \emptyset$. We now define by induction a sequence of f.p. homotopies $H^n: B \times M \times I \rightarrow B \times M$ and $h_n = H_1^n$ such that

- (i)_n $H^n: h_{n-1} \simeq h_n$ (f.p.) rel $(A \times M) \cup (C_{n-1} \times M) \cup (B \times M_{n-1})$,
- (ii)_n $h_n|_{C_n \times M}: C_n \times M \rightarrow C_n \times M$ is an f.p. homeomorphism, and
- (iii)_n $h_n|_{B \times M_n}: B \times M_n \rightarrow B \times M$ is an f.p. embedding.

First, since $f|_{A \times M_1}$ is an f.p. embedding, there is from [L₃, Lemma 3.3] an f.p. homotopy $F: B \times M \times I \rightarrow B \times M$ (rel $A \times M_1$) such that $F_0 = f$ and F_1 is an f.p. homeomorphism. Let $s: B \rightarrow I$ be a map with $s^{-1}(0) = A$ and $s^{-1}(1) = C_1$. Define $H^1: B \times M \times I \rightarrow B \times M$ by $H^1(b, x, t) = F(b, x, s(b)t)$. Then we have

- (i)₁ $H^1: f (\equiv h_0) \simeq h_1$ (f.p.) rel $A \times M$,
- (ii)₁ $h_1|_{C_1 \times M} = F_1|_{C_1 \times M}$ is an f.p. homeomorphism, and
- (iii)₁ $h_1|_{B \times M_1}: B \times M_1 \rightarrow B \times M$ is an f.p. embedding, by use of [L₃, Lemma 1.2].

Second, assume that H^{n-1} has been defined such that (i)_{n-1}, (ii)_{n-1}, and (iii)_{n-1} are satisfied. By [L₃, Lemma 3.3], there is an f.p. homotopy $G: B \times M \times I \rightarrow B \times M$ (rel $A \times M_n \cup C_{n-1} \times M \cup B \times M_{n-1}$) such that $G_0 = h_{n-1}$ and G_1 is an f.p. homeomorphism. Similar to above, by use of an Urysohn function and [L₃, Lemma 1.2], we can obtain a homotopy H^n such that (i)_n, (ii)_n, and (iii)_n are satisfied.

Finally, define $h = \lim h_n$; then, h is well defined and continuous by use of (iii)_n; and, $h|(B - A) \times M$ is an f.p. homeomorphism by use of (ii)_n and the condition $C_n \subset$ (interior of C_{n+1}). Moreover, a desired homotopy $H: B \times M \times I \rightarrow B \times M$ from f to h can be defined by

$$H(b, x, t) = \begin{cases} H^n(b, x, 2^n(t-1) + 2) & \text{if } t \in J_n, \\ h(x) & \text{if } t = 1, \end{cases}$$

where $J_n = [1 - (1/2^{n-1}), 1 - (1/2^n)]$. Then, H is well defined and continuous by (i)_n. $\square$

2. Absolute neighborhood extensor properties of $\text{Homeo}(M)$. We first prove some lemmas that will be used in the proof of the main result of this section, Theorem 2.3.

LEMMA 2.1. *Let K be a compact metric space and M an E^∞ -manifold. Then the space $M^K \in \mathcal{CW}(M)$; moreover, it is an $\text{ANE}(M)$ and an $\text{ANE}(\mathcal{C})$. Consequently, M^K is an $\text{ANE}(\mathcal{CW}(M))$ and an $\text{ANE}(\mathcal{CW}(\mathcal{C}))$, and so is $C(K, M)$.*

PROOF. The proof is implicit in that of [Hy, Theorem 8.2]. Because of its simplicity in this case, we will give its outline.

Write $M = \text{dirlim } M_n$, where each M_n is a compact ANR metric space with $M_n \subset M_{n+1}$. Observe that

(1) by [Hy, Proposition 2.7], $M^K = \bigcup \{(M_n)^K | n = 1, 2, \dots\}$,

(2) by [Hy, Lemma 8.1.b], $(M_n)^K$ is a closed subspace of $(M_{n+1})^K$ for each n , and

(3) each $(M_n)^K$ is an ANR complete metric space (so, it is an $\text{ANE}(\mathcal{C})$ by [M, Theorem 3.1(b)]).

Therefore, by [Hy, Theorem 11.3], $\text{dirlim}(M_n)^K \in \mathcal{CW}(M)$ (or an M -space of [Hy]) and is an $\text{ANE}(M)$. On the other hand, let P be a compact subset of M^K and $e: M^K \times K \rightarrow M$ the evaluation map. Then $e(P \times K)$ is a compact subset of M . So, there is an n such that $e(P \times K) \subset M_n$; in other words, $P \subset (M_n)^K$ for some n . Therefore, $M^K = \text{dirlim}(M_n)^K$ by [Hy, Lemma 5.5]. Also, observe that M^K is an $\text{ANE}(\mathcal{C})$ by use of (3).

Now, it follows from [Hy, Theorem 10.2] and its proof that $M^K = \text{dirlim}(M_n)^K$ is an $\text{ANE}(\mathcal{CW}(M))$ and an $\text{ANE}(\mathcal{CW}(\mathcal{C}))$. Finally, it follows easily that $C(K, M)$ is an $\text{ANE}(\mathcal{CW}(M))$ and an $\text{ANE}(\mathcal{CW}(\mathcal{C}))$ by use of [Gr, 8.8] and $\mathcal{CW}(M) \cup \mathcal{CW}(\mathcal{C}) \subset \mathcal{CG}$ [L₃, §0]. $\square$

LEMMA 2.2. *Let K be a finite complex and $N = K \times E^\infty$. Let A be a closed subset (G_δ -subset, resp.) of $B \in \mathcal{CW}(M)$ ($B \in \mathcal{CW}(\mathcal{C})$, resp.). Then each map $f: A \rightarrow \text{Homeo}(N) \subset \text{Homo}(N)$ has a continuous extension $g: U \rightarrow \text{Homo}(N)$ over a neighborhood U of A in B ; i.e., the associate $g: U \times N \rightarrow U \times N$ of g is an f.p. homotopy equivalence.*

PROOF. Identify K with $K \times 0 \subset N$ and let $\hat{f}: A \times N \rightarrow A \times N$ be the associate of f . We will first show that $\hat{f}$ has an f.p. extension $q: V \times N \rightarrow V \times N$, where V is a closed neighborhood of A in B . Then, we will choose a suitable neighborhood U of A in V such that $\hat{g} = q|U \times N$ is an f.p. homotopy equivalence.

Define

$$f_K: A \xrightarrow{f} \text{Homeo}(N) \subset C(N, N) \xrightarrow{\tilde{i}_K} C(K, N),$$

where $\tilde{i}_K$ is induced from i_K . We have f_K being continuous. It follows from Lemma 2.1 that there is a closed neighborhood V of A in B and an extension $h: V \rightarrow C(K, N)$ of f_K . Let $\hat{h}: V \times K \rightarrow V \times N$ be its associate [Gr, 8.17] and $R: N \times I \rightarrow N$ a strong deformation retraction of N onto K . Define $d: V \times N \rightarrow V \times N$ by $d = \hat{h}(\text{id}_V \times R_1)$. Then,

$$\begin{aligned} d|A \times N &= \hat{h}(\text{id}_A \times R_1) = \hat{f}_K(\text{id}_A \times R_1) = \hat{f}(\text{id}_A \times R_1) \\ &\simeq \hat{f}(\text{id}_A \times \text{id}_N) \quad (\text{f.p.}) \text{ since } R_1 \simeq \text{id}_N \\ &= \hat{f}. \end{aligned}$$

Therefore, by [L₃, Lemma 1.1], d is f.p. homotopic to a map $q: V \times N \rightarrow V \times N$ such that $q|A \times N = \hat{f}$.

Fix an $a \in A$. Consider $\hat{G} = q(\text{id}_V \times \hat{f}_a^{-1})$ with its continuous associate $G: V \rightarrow C(N, N)$. We have $G(a) = \hat{f}_a \hat{f}_a^{-1} = \text{id}_N$. From the proof of [L₄, Theorem 1], there is an open neighborhood W of id_N in $C(N, N)$ such that for every map $s: X \rightarrow W$, its associate $\hat{s}: X \times N \rightarrow X \times N$ is f.p. homotopic to $\text{id}_{X \times N}$. Let $U_a = G^{-1}(W)$. Then, the restriction $\hat{G}|_{U_a \times N}$ is an f.p. homotopy equivalence; hence, so is $q|_{U_a \times N} = \hat{G}(\text{id}_{U_a} \times \hat{f}_a)$. Define $U = \bigcup \{U_a | a \in A\}$. Then, $\hat{g} = q|_{U \times N}$ is a local f.p. homotopy equivalence; hence, it is an f.p. homotopy equivalence by [Ja, Theorem 5.57], and its associate $g: U \rightarrow \text{Homo}(N)$ is an extension of f as desired. $\square$

THEOREM 2.3. *If M is an E^∞ -manifold having homotopy type of a finite complex, then $\text{Homeo}(M)$ is an $\text{ANE}(\mathcal{CW}(M))$ and an $\text{ANE}(\mathcal{CW}(C), G_\delta)$; hence, so is $k\text{Homeo}(M)$.*

PROOF. By [He, Theorem C], we can write $M = K \times E^\infty$, where K is a finite complex. Let A be a closed subset (G_δ -subset, resp.) of $B \in \mathcal{CW}(M)$ ($B \in \mathcal{CW}(C)$, resp.), and $f: A \rightarrow \text{Homeo}(M)$ a map. Recall that since $A, B \in \mathcal{CG}$, we can use either topology on $\text{Homeo}(M)$ and $\text{Homo}(M)$. From Lemma 2.2, f has an extension $g: U \rightarrow \text{Homo}(M)$, where U is a closed neighborhood of A in B , such that its associate $\hat{g}: U \times M \rightarrow U \times M$ is an f.p. homotopy equivalence. Then, it follows from Proposition 1.4 that there is an f.p. map $\hat{h}: U \times M \rightarrow U \times M$ whose associate $h: U \rightarrow C(M, M)$ satisfies the following properties:

- (a) $h(U) \subset \text{Homeo}(M)$, and
- (b) $h|_A = f$.

In other words, h is an extension of f over U into $\text{Homeo}(M)$. $\square$

REFERENCES

- [D] J. Dugundji, *Topology*, Allyn and Bacon, Boston, Mass., 1978.
- [Gr] B. Gray, *Homotopy theory: An introduction to algebraic topology*, Academic Press, New York, 1975.
- [He] R. E. Heisey, *Manifolds modelled on the direct limit of lines*, Pacific J. Math. **102** (1982), 47–54.
- [Hu] S. T. Hu, *Theory of retracts*, Wayne State Univ. Press, 1965.
- [Hy] D. M. Hyman, *A category slightly larger than the metric and CW categories*, Michigan J. Math. **15** (1968), 193–214.
- [Ja] I. M. James, *General topology and homotopy theory*, Springer-Verlag, New York, 1984.
- [J] J. R. Jackson, *Spaces of mappings on topological products with applications to homotopy theory*, Proc. Amer. Math. Soc. **3** (1952), 327–333.
- [Ku] K. Kuratowski, *Topology*, Vol. II, Academic Press, New York, 1968.
- [L₁] V. T. Liem, *An unknotting theorem in Q^∞ -manifolds*, Proc. Amer. Math. Soc. **82** (1981), 125–132.
- [L₂] —, *An α -approximation theorem for R^∞ -manifolds*, Rocky Mountain J. Math. (to appear).
- [L₃] —, *On $R^\infty(Q^\infty)$ -manifold bundles over CW-complexes*, Trans. Amer. Math. Soc. **297** (1986), 563–585.
- [L₄] —, *A local strong UV^∞ -property of the homeomorphism groups of $R^\infty(Q^\infty)$ -manifolds*, Proc. Amer. Math. Soc. **98** (1986), 145–149.
- [M] E. Michael, *Some extension theorems for continuous functions*, Pacific J. Math. **3** (1953), 789–806.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ALABAMA, TUSCALOOSA, ALABAMA 35487