ISOMORPHISMS OF GRAPH GROUPS

CARL DROMS

ABSTRACT. Given a graph X , define the presentation PX to have generators the vertices of X , and a relation $xy = yx$ for each pair x, y of adjacent vertices. Let GX be the group with presentation PX , and given a field K , let KX denote the K -algebra with presentation PX . Given graphs X and Y and a field K , it is known that the algebras KX and KY are isomorphic if and only if the graphs X and Y are isomorphic. In this paper, we use this fact to prove that if the groups GX and GY are isomorphic, then so are the graphs X and Y .

Given a graph X with vertex set $V(X)$, we define the presentation PX to be that having as generators the elements of $V(X)$, with a defining relation $vw = wv$ for each pair v and w of adjacent vertices of X . PX can be regarded as a presentation of a group GX , or of a K -algebra KX over a field K . In [1], Kim, Makar-Limanov, Neggers, and Roush proved that if the algebras KX and KY are isomorphic, then so are the graphs X and Y . (In their formulation, two generators commute provided they are *not* adjacent in the graph. This is sufficient for our purposes, since if two graphs are isomorphic, so are their complements.)

Let K be a field. In this note we will show that if the groups GX and GY are isomorphic, then so are the algebras KX and KY , thus demonstrating:

THEOREM. *If the groups GX and GY are isomorphic, then so are the graphs X and Y .*

Let $f: GX \rightarrow GY$ be an isomorphism. Denote by G_2X the quotient group $GX/[GX, (GX)']$. Then f induces an isomorphism $f_2: G_2X \rightarrow G_2Y$. Let $V(X)$ and $V(Y)$ be totally ordered, and denote both orderings by $<$. We will not distinguish by notation between a vertex of X , the corresponding element of GX , and the image of this element in G_2X . For each vertex x of X , $f_2(x)$ can be written uniquely in the form $y_1^{a_1}y_2^{a_2}\cdots y_n^{a_n}C_x$, where C_x is an element of the commutator subgroup $(G_2Y)'$, $y_1 < y_2 < \cdots < y_n$, and the integers a_r are all nonzero. Define $f_*(x) = a_1y_1 + a_2y_2 + \cdots + a_ny_n$. We will show that the function $f_*: X \rightarrow KY$ extends to a homomorphism $f_*: KX \rightarrow KY$ by showing that if $xx' = x'x$ is a relation of PX , then $f_*(x)f_*(x') = f_*(x)f_*(x')$ in KY .

LEMMA. *The commutators $\{[x_i, x_j] \mid x_i < x_j \text{ and } x_i \text{ and } x_j \text{ are not adjacent in } X\}$ of G_2X are linearly independent.*

PROOF. Consider the exact sequence

$$1 \rightarrow N \rightarrow FX \rightarrow GX \rightarrow 1$$

Received by the editors January 21, 1985 and, in revised form, May 12, 1986.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 20F05; Secondary 20F12.

©1987 American Mathematical Society
 0002-9939/87 \$1.00 + \$.25 per page

associated with the presentation PX . Since N is a subgroup of FX' , there is an exact sequence

$$1 \rightarrow \hat{N} \rightarrow (F_2X)' \rightarrow (G_2X)' \rightarrow 1$$

where $\hat{N}$ denotes the image of N in F_2X . By the Basis Theorem, $(F_2X)'$ is a free abelian group with a basis consisting of the commutators $\{[x_i, x_j] \mid x_i < x_j\}$. Since $\hat{N}$ is the subgroup of $(F_2X)'$ generated by the collection $\{[x_i, x_j] \mid x_i < x_j, x_i \text{ and } x_j \text{ adjacent}\}$, the assertion follows.

Suppose that x and x' are adjacent vertices of X . Let $f_2(x) = y_1^{a_1} y_2^{a_2} \cdots y_n^{a_n} C_x$ and $f_2(x') = y_1^{b_1} y_2^{b_2} \cdots y_n^{b_n} C_{x'}$, with $y_1 < y_2 < \cdots < y_n$, where we allow some of the exponents a_r and b_s to be 0 so that we may use the same elements of $V(Y)$ in both expressions. Writing $(G_2Y)'$ additively, and recalling that in G_2Y , commutators are central, we have

$$\begin{aligned} 0 &= [f_2(x), f_2(x')] = \sum_{r,s} [y_r^{a_r}, y_s^{b_s}] \\ &= \sum_{r,s} a_r b_s [y_r, y_s] = \sum_{r < s} (a_r b_s - a_s b_r) [y_r, y_s]. \end{aligned}$$

Thus, by the Lemma, if y_r and y_s are nonadjacent vertices of Y , then $a_r b_s - a_s b_r = 0$. Clearly, $f_*(x) = \sum_r a_r y_r$ and $f_*(x') = \sum_s b_s y_s$, so that

$$f_*(x)f_*(x') - f_*(x')f_*(x) = \sum_{r,s} (a_r b_s - b_r a_s) y_r y_s.$$

If y_r and y_s are adjacent vertices of Y , then $y_r y_s = y_s y_r$ in KY , so the net coefficient of $y_r y_s$ in this sum is $(a_r b_s - b_r a_s) + (a_s b_r - b_s a_r) = 0$. If y_r and y_s are nonadjacent vertices of Y , then the net coefficient of $y_r y_s$ is $a_r b_s - b_r a_s$, and we saw above that this must be 0 when y_r and y_s are not adjacent. Finally, for each r , the coefficient of y_r^2 is $a_r b_r - b_r a_r = 0$. Thus, $f_*(x)f_*(x') - f_*(x')f_*(x) = 0$, so f_* extends to a homomorphism from KX to KY .

It is now an easy matter to check that for each vertex x of X , $(f^{-1})_* f_*(x) = x$, and for each vertex y of Y , $f_*(f^{-1})_*(y) = y$, so that f_* is in fact an isomorphism.

The following shorter proof of our theorem has been pointed out by an anonymous reviewer: Define $G_0 = GX$ and $G_{n+1} = [G_n, G]$ for $n \geq 0$, and let LX denote the Lie algebra $K \otimes \sum G_n / G_{n+1}$. Then the algebra KX is the universal enveloping algebra of LX , so that if GX and GY are isomorphic groups, then the Lie algebras LX and LY , and hence the algebras KX and KY , are isomorphic.

REFERENCES

1. K. H. Kim, L. Makar-Limanov, J. Neggers and F. Roush, *Graph algebras*, J. Algebra **64** (1980), 46–51.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, JAMES MADISON UNIVERSITY, HARRISONBURG, VIRGINIA 22807