ARENS PRODUCT AND THE ALGEBRA OF DOUBLE MULTIPLIERS. II

PAK-KEN WONG

ABSTRACT. We show that the algebra of double multipliers of a certain Banach algebra A can be embedded in the second conjugate space A^{**} of A . This generalizes the previous work by the author for B^* -algebras.

1. Introduction. Let A be a semisimple Banach algebra and $M(A)$ the algebra of double multipliers on A . The author recently showed that $M(A)$ is isomorphic to $(A^{**}, \circ)$ if and only if A has the following properties: (1) A is Arens regular, (2) A has an approximate identity, and (3) $\pi(A)$ is an ideal of $(A^{**}, \circ)$ [10, Theorem, p. 442]. In general, not many Banach algebras possess all these properties. The purpose of this paper is to establish an embedding of $M(A)$ in the second conjugate space A^{**} of A for a semisimple Banach algebra A with an approximate identity. In fact, we show that $M(A)$ is isometrically isomorphic to M^{**}/N^{**} , where $M^{**} = \{F \in A^{**}: F \circ \pi(x) \text{ and } \pi(x) \circ F \in \pi(A) \text{ for all } x \in A\}$ and $N^{**} = \{F \in A^{**}: F(f \circ \pi) = F(x \circ' f) = 0 \text{ for all } x \in A \text{ and } f \in a^*\}$. The main idea of the proof of this result is essentially contained in the proofs of [8, Lemma 2.1, p. 80 and 10, Theorem, p. 422]. This embedding was studied in [1 and 8] for B^* -algebras.

2. Notation and preliminaries. Definitions not explicitly given are taken from Rickart's book [6].

Let A be a Banach algebra. Then A^* and A^{**} will denote the first and second conjugate spaces of A , and π the canonical map of A into A^{**} . The two Arens products on A^{**} are defined in stages according to the following rules (see [2]). Let $x, y \in A$, $f \in A^*$, and $F, G \in A^{**}$.

Define $f \circ x$ by $(f \circ x)(y) = f(xy)$. Then $f \circ x \in A^*$.

Define $G \circ f$ by $(G \circ f)(x) = G(f \circ x)$. Then $G \circ f \in A^*$.

Define $F \circ G$ by $(F \circ G)(f) = F(G \circ f)$. Then $F \circ G \in A^{**}$.

Define $x \circ' f$ by $(x \circ' f)(y) = f(yx)$. Then $x \circ' f \in A^*$.

Define $f \circ' F$ by $(f \circ' F)(x) = F(x \circ' f)$. Then $f \circ' F \in A^*$.

Define $F \circ' G$ by $(F \circ' G)(f) = G(f \circ' F)$. Then $F \circ' G \in A^{**}$.

Received by the editors February 25, 1986.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 46H10; Secondary 46H99.

Key words and phrases. Arens products, Arens regularity, approximate identity, double multiplier, B^* -algebra, conjugate space.

This research was done while the author was visiting the University of Hong Kong. The author gratefully acknowledges the hospitality of the Department of Mathematics of the University.

©1987 American Mathematical Society
0002-9939/87 \$1.00 + \$.25 per page

A^{**} is a Banach algebra under the products $F \circ G$ and $F \circ' G$ and π is an algebra isomorphism of A into $(A^{**}, \circ)$ and $(A^{**}, \circ')$. In general, $\circ$ and $\circ'$ are distinct on A^{**} . If they agree on A^{**} , then A is called Arens regular.

The following properties of the Arens products will be used often in the rest of this paper.

LEMMA 2.1. *Let A be a Banach algebra. Then for all $x \in A$, $f \in A^*$, and $F, G \in A^{**}$, we have*

- (1) $\pi(x) \circ F = \pi(x) \circ' F$ and $F \circ \pi(x) = F \circ' \pi(x)$.
- (2) If $\{F_\beta\} \subset A^{**}$ and $F_\beta \rightarrow F$ weakly in A^{**} , then $F_\beta \circ G \rightarrow F \circ G$ and $G \circ' F_\beta \rightarrow G \circ' F$ weakly.
- (3) $F(x \circ' f) = (F \circ \pi(x))(f)$.

PROOF. (1) is proved in [2, p. 843] and (2) in [2, p. 842].

(3) It is easy to see that $\pi(x) \circ f = x \circ' f$. Therefore

$$(F \circ \pi(x))(f) = F(\pi(x) \circ f) = F(x \circ' f).$$

This completes the proof of the lemma.

We say that a Banach algebra A has an approximate identity if there exists a net $\{e_\alpha\}$ in A such that $\|e_\alpha\| \leq 1$ for all α and $x = \lim e_\alpha x = \lim x e_\alpha$ for all $x \in A$.

Let A be a semisimple Banach algebra. A pair (T_1, T_2) of operators from A to A is called a double multiplier (centralizer) on A provided that $x(T_1 y) = (T_2 x)y$ for all $x, y \in A$. It is known that T_1 and T_2 are continuous linear operators on A such that $T_1(xy) = (T_1 x)y$ and $T_2(xy) = x(T_2 y)$. The set $M(A)$ of all double multipliers on A is a Banach algebra with identity and A can be identified as a two-sided ideal of $M(A)$ (see [4 and 5]).

In this paper, all algebras and spaces under consideration are over the complex field.

3. Extensions of double multipliers. Let A be a semisimple Banach algebra. We show that each double multiplier $T = (T_1, T_2)$ on A can be extended to a double multiplier on $(A^{**}, \circ)$. In fact, for all $x \in A$ and $f \in A^*$, we define

$$(f \circ T_1)(x) = f(T_1 x) \quad \text{and} \quad (f \circ T_2)(x) = f(T_2 x).$$

Then $f \circ T_1$ and $f \circ T_2 \in A^*$. For all $F \in A^{**}$, we define

$$T_1^\#(F)(f) = F(f \circ T_1) \quad \text{and} \quad T_2^\#(F)(f) = F(f \circ T_2).$$

Then $T_1^\#(f)$ and $T_2^\#(F) \in A^{**}$.

THEOREM 3.1. *Let A be a semisimple Banach algebra. Then for each $T = (T_1, T_2) \in M(A)$, $T^\# = (T_1^\#, T_2^\#)$ is a continuous double multiplier on $(A^{**}, \circ)$ and $T^\# \upharpoonright \pi(A) = T$.*

PROOF. Let F and $G \in A^{**}$. Then by Goldstine's Theorem, there exist nets $\{x_m\}$ and $\{y_n\}$ in A such that $\pi(x_m) \rightarrow F$ and $\pi(y_n) \rightarrow G$ weakly in A^{**} . Then for all $f \in A^*$, we have

$$\begin{aligned} (F \circ T_1^\#(G))(f) &= \lim_m (T_1^\#(G) \circ f)(x_m) = \lim_m G((f \circ x_m) \circ T_1) \\ &= \lim_m \lim_n ((f \circ x_m) \circ T_1)(y_n) \\ &= \lim_m \lim_n f((T_2 x_m) y_n) = \lim_m G(f \circ T_2 x_m) \\ &= \lim_m ((G \circ f) \circ T_2)(x_m) = (T_2^\#(F) \circ G)(f). \end{aligned}$$

Therefore $F \circ T_1^\#(G) = T_2^\#(F) \circ G$. Similarly, we can show that $T_1^\#(F \circ G) = T_1^\#(F) \circ G$ and $T_2^\#(F \circ G) = F \circ T_2^\#(G)$. It is easy to see that $\|T_1^\#\| \leq \|T_1\|$ and $\|T_2^\#\| \leq \|T_2\|$. Therefore $\|T^\#\| \leq \|T\|$. Since $T_1^\#(\pi(x)) = \pi(T_1 x)$ and $T_2^\#(\pi(x)) = \pi(T_2 x)$, we have $T^\# \mid \pi(A) = T$ and the theorem is proved.

4. The subalgebras M^{} and N^{**} of $(A^{**}, \circ)$.** For any Banach algebra A , set

$$M^{**} = \{F \in A^{**}: F \circ \pi(x) \text{ and } \pi(x) \circ F \in \pi(A) \text{ for all } x \in A\}$$

and

$$N^{**} = \{F \in A^{**}: F(f \circ x) = F(x \circ' f) = 0 \text{ for all } x \in A \text{ and } f \in A^*\}.$$

Then $\pi(A)$ is a two-sided ideal of $(A^{**}, \circ)$ if and only if $M^{**} = A^{**}$.

LEMMA 4.1. *Let A be a Banach algebra. Then M^{**} and N^{**} are closed subalgebras of $(A^{**}, \circ)$ and $(A^{**}, \circ')$. Also N^{**} is a two-sided ideal in M^{**} and $N^{**} = \{F \in A^{**}: A^{**} \circ F = F \circ' A^{**} = (0)\}$.*

PROOF. For any $F \in N^{**}$, easy calculations show that $\pi(A) \circ F = (0)$ and $F \circ \pi(A) = (0)$. Hence $A^{**} \circ F = F \circ' A^{**} = (0)$ and so $N^{**} = \{F \in A^{**}: A^{**} \circ F = F \circ' A^{**} = (0)\}$. Thus if $F, G \in N^{**}$, then $\pi(A) \circ (F \circ G) = (F \circ G) \circ \pi(A) = (0)$ and so $F \circ G \in N^{**}$. Similarly, $F \circ' G \in N^{**}$. It is easy to check that M^{**} and N^{**} are closed subalgebras in $(A^{**}, \circ)$ and $(A^{**}, \circ')$ and N^{**} is a two-sided ideal of M^{**} . Therefore the lemma is proved.

In the rest of this section, A will be a semisimple Banach algebra with an approximate identity $\{e_\alpha\}$. It follows from [3, Proposition 7, p. 146] that $(A^{**}, \circ)$ has a right identity I . Also I is a left identity for $(A^{**}, \circ')$.

THEOREM 4.2. *Let A be a semisimple Banach algebra with an approximate identity $\{e_\alpha\}$. Then the Arens products $\circ$ and $\circ'$ agree on M^{**}/N^{**} and M^{**}/N^{**} is a Banach algebra with an identity.*

PROOF. Let $F, G \in M^{**}$, $f \in A^*$, and $x \in A$. Then by Cohen's Factorization Theorem [3, Theorem 10, p. 61], $x = yz$ with $y, z \in A$. Hence

$$\begin{aligned} (F \circ G - F \circ' G)(f \circ x) &= (F \circ G - F \circ' G)((f \circ y) \circ z) \\ &= (\pi(z) \circ F \circ G - \pi(z) \circ F \circ G)(f \circ y) = 0. \end{aligned}$$

Similarly, $(F \circ G - F \circ' G)(x \circ' f) = 0$. Therefore $F \circ G - F \circ' G \in N^{**}$ and so $\circ$ and $\circ'$ agree on M^{**}/N^{**} . Since I is a right identity for $(A^{**}, \circ)$ and a left identity for $(A^{**}, \circ')$ and $I \in M^{**}$, it follows that M^{**}/N^{**} is a Banach algebra with an identity.

REMARK. For a certain Banach algebra A , N^{**} is the radical of $(A^{**}, \circ)$ (for example, see [9, Theorem 4.1, p. 440]).

The following result is essentially contained in the proof of [8, Lemma 2.1, p. 80]. It is useful in the next section.

LEMMA 4.3. *Let A be a semisimple Banach algebra with an approximate identity $\{e_\alpha\}$, $T = (T_1, T_2) \in M(A)$, and $x \in A$. Then*

- (1) $T_1^\#(I) \circ \pi(x) = T_2^\#(I) \circ \pi(x) = \pi(T_1 x)$.
- (2) $\pi(x) \circ T_1^\#(I) = \pi(x) \circ T_2^\#(I) = \pi(T_2 x)$.
- (3) $T_1^\#(I) = T_2^\#(I) + N$ for some $N \in N^{**}$.
- (4) If $T_1^\#(I)$ and $T_2^\#(I)$ are both in N^{**} , then $T = 0$.

PROOF. (1) For all $f \in A^*$, we have

$$\begin{aligned} (T_1^\#(I) \circ \pi(x))(f) &= I((\pi(x) \circ f) \circ T_1) = \lim_{\alpha} ((\pi(x) \circ f) \circ T_1)(e_\alpha) \\ &= \lim_{\alpha} f(T_1(e_\alpha x)) = \lim_{\alpha} (f \circ T_1)(e_\alpha x) \\ &= (f \circ T_1)(x) = f(T_1 x). \end{aligned}$$

Hence $T_1^\#(I) \circ \pi(x) = \pi(T_1 x)$. Similarly, $T_2^\#(I) \circ \pi(x) = \pi(T_2 x)$. Therefore (1) is proved. Similarly we can prove (2).

(3) Let $N = T_1^\#(I) - T_2^\#(I)$. Then

$$\begin{aligned} N(f \circ x) &= \pi(x)(N \circ f) = (\pi(x) \circ N)(f) \\ &= (\pi(x) \circ T_1^\#(I) - \pi(x) \circ T_2^\#(I))(f) \\ &= (\pi(T_1 x) - \pi(T_2 x))(f) = 0. \end{aligned}$$

Similarly, $N(x \circ' f) = 0$. Therefore $N \in N^{**}$ and so $T_1^\#(I) = T_2^\#(I) + N$.

(4) Since $T_1^\#(I)$ and $T_2^\#(I) \in N^{**}$, we have

$$\pi(T_1 x)(f) = (T_2^\#(I) \circ \pi(x))(f) = T_2^\#(I)(x \circ' f) = 0.$$

Hence $T_1 x = 0$ and so $T_1 = 0$. Also

$$\pi(T_2 x)(f) = (\pi(x) \circ T_1^\#(I))(f) = T_1^\#(I)(f \circ x) = 0.$$

Therefore $T_2 = 0$ and so $T = 0$. This completes the proof of the lemma.

Notation. In the rest of this paper, for each $F \in M^{**}$, we write $\tilde{F} = F + N^{**}$.

5. The algebras $M(A)$ and M^{}/N^{**} .** We have the main result of this paper.

THEOREM 5.1. *Let A be a semisimple Banach algebra with an approximate identity $\{e_\alpha\}$. Then $M(A)$ is isometrically isomorphic to M^{**}/N^{**} .*

PROOF. Let $T = (T_1, T_2) \in M(A)$. Then by Lemma 4.3, $T_1^\#(I)$ and $T_2^\#(I) \in M^{**}$ and $\widetilde{T_1^\#(I)} = \widetilde{T_2^\#(I)}$. We define a mapping Φ from $M(A)$ to M^{**}/N^{**} by

$$\Phi(T) = \widetilde{T_1^\#(I)} = \widetilde{T_2^\#(I)}.$$

We show that Φ is an isometric isomorphism from $M(A)$ onto M^{**}/N^{**} . It is clear that Φ is linear. Let $x \in A$ and $f \in A^*$. If $\Phi(T) = 0$, then $T_1^\#(I)$ and $T_2^\#(I)$ are both in N^{**} . Hence by Lemma 4.3, $T = 0$. Therefore Φ is one-one. Let $S = (S_1, S_2) \in M(A)$. Since $ST = (S_1T_1, T_2S_2)$, by Lemma 4.3, we have

$$\begin{aligned} (S_1^\#(I) \circ T_1^\#(I))(f \circ x) &= (\pi(x) \circ S_1^\#(I) \circ T_1^\#(I))(f) \\ &= (\pi(S_2x) \circ T_1^\#(I))(f) = \pi(T_2S_2x)(f) \\ &= (\pi(x) \circ (S_1T_1)^\#(I))(f) = (S_1T_1)^\#(I)(f \circ x). \end{aligned}$$

Similarly, we have

$$(S_1^\#(I) \circ T_1^\#(I))(x \circ' f) = (S_1T_1)^\#(I)(x \circ' f).$$

Therefore $\Phi(ST) = \Phi(S)\Phi(T)$ and so Φ is an isomorphism. Let $F \in M^{**}$. Define

$$P_1(x) = F \circ \pi(x) \quad \text{and} \quad P_2(x) = \pi(x) \circ F \quad (x \in A).$$

Since $F \in M^{**}$, $P = (P_1, P_2) \in M(A)$. Also

$$\begin{aligned} P_1^\#(I)(f \circ x) &= I((f \circ x) \circ P_1) = \lim_{\alpha} ((f \circ x) \circ P_1)(e_{\alpha}) \\ &= \lim_{\alpha} f(x(F \circ \pi(e_{\alpha}))) = \lim_{\alpha} f((\pi(x) \circ F)e_{\alpha}) \\ &= \lim_{\alpha} f(e_{\alpha}(\pi(x) \circ F)) = \lim_{\alpha} f(\pi(e_{\alpha}x) \circ F) \\ &= \lim_{\alpha} (\pi(e_{\alpha}x) \circ F)(f) = \lim_{\alpha} \pi(e_{\alpha}x)(F \circ f) \\ &= (F \circ f)(x) = F(f \circ x). \end{aligned}$$

Similarly, we have $P_1^\#(I)(x \circ' f) = F(x \circ' f)$. Therefore $\Phi(P) = \widetilde{P_1^\#} = \tilde{F}$. Hence Φ is an onto mapping. It is easy to see that $\|\Phi(T)\| \leq \|T\|$. Since $\|T_1\| = \sup\{|f(T_1x)| : \|x\| \leq 1, \|f\| \leq 1 \text{ for all } x \in A \text{ and } f \in A^*\}$, for given $\varepsilon > 0$, there exists $y \in A$ and $g \in A^*$ with $\|y\| \leq 1$ and $\|g\| \leq 1$ such that $|g(T_1y)| \geq \|T_1\| - \varepsilon$. Then

$$\begin{aligned} I((y \circ' g) \circ T_1) &= \lim_{\alpha} (y \circ' g)(T_1e_{\alpha}) = \lim_{\alpha} g((T_1e_{\alpha})y) = \lim_{\alpha} g(T_1(e_{\alpha}y)) \\ &= \lim_{\alpha} (g \circ T_1)(e_{\alpha}y) = (g \circ T_1)(y) = g(T_1y). \end{aligned}$$

Hence for all $N \in N^{**}$, we have

$$\begin{aligned} \|T_1^\#(I) + N\| &\geq |(T_1^\#(I) + N)(y \circ' g)| = |T_1^\#(I)(y \circ' g)| \\ &= |I((y \circ' g) \circ T_1)| = |g(T_1y)| \geq \|T_1\| - \varepsilon. \end{aligned}$$

Since ε is arbitrary, we have

$$\|\Phi(T)\| = \|\widetilde{T_1^\#(I)}\| \geq \|T_1\|.$$

Similarly, we can show that $\|\Phi(T)\| \geq \|T_2\|$ and so $\|\Phi(T)\| \geq \|T\|$. Therefore $\|\Phi(T)\| = \|T\|$. This completes the proof of the theorem.

COROLLARY 5.2. *Let A be as in Theorem 5.1. If $(A^{**}, \circ)$ has an identity I , then $N^{**} = (0)$ and so $M(A)$ is isometrically isomorphic to M^{**} .*

PROOF. Let $N \in N^{**}$. Then for all $f \in A^*$, we have

$$N(f) = (I \circ N)(f) = \lim_{\alpha} (\pi(e_{\alpha}) \circ N)(f) = \lim_{\alpha} N(f \circ e_{\alpha}) = 0.$$

Therefore $N^{**} = (0)$ and the corollary follows from Theorem 5.1.

COROLLARY 5.3. *Let A be as in Theorem 5.1. If A is Arens regular, then $M(A)$ is isometrically isomorphic to M^{**} .*

PROOF. Since A is Arens regular, $(A^{**}, \circ)$ has an identity I . Therefore the result follows from Corollary 5.2.

We now have a slight improvement of [10, Theorem, p. 442].

THEOREM 5.4. *Let A be a semisimple Banach algebra. Then $M(A)$ is isometrically isomorphic to $(A^{**}, \circ)$ if and only if A has the following properties:*

- (1) A is Arens regular.
- (2) A has an approximate identity.
- (3) $\pi(A)$ is a two-sided ideal of $(A^{**}, \circ)$.

PROOF. Suppose that $M(A)$ is isometrically isomorphic to $(A^{**}, \circ)$. Then by Theorem 5.1, we have $N^{**} = (0)$ and $M^{**} = A^{**}$. Hence $\pi(A)$ is a two-sided ideal of $(A^{**}, \circ)$. By Theorem 4.2, A is Arens regular and $(A^{**}, \circ)$ has an identity. Hence it follows from [3, Proposition 7, p. 147] that A has an approximate identity. Therefore A has properties (1), (2), and (3).

Conversely, suppose that A has properties (1), (2), and (3). Then $M^{**} = A^{**}$ and so by Corollary 5.3, $M(A)$ is isometrically isomorphic to $(A^{**}, \circ)$. This completes the proof of the theorem.

6. Banach $*$ -algebras. Let A be a Banach $*$ -algebra with a continuous involution. For all $x \in A$, $f \in A^*$, and $F \in A^{**}$, we define

$$f^*(x) = \overline{f(x^*)} \quad \text{and} \quad F^*(f) = \overline{F(f^*)},$$

where the bar denotes the complex conjugation. Then $f^* \in A$ and $F^* \in A^{**}$. If A is a B^* -algebra, then $(A^{**}, \circ)$ is a B^* -algebra under the involution $F \rightarrow F^*$ (see [7, p. 192]).

LEMMA 6.1. *Let A be a Banach $*$ -algebra with a continuous involution. Then*

- (1) *For all F and $G \in A^{**}$, we have*

$$(F \circ G)^* = G^* \circ' F^* \quad \text{and} \quad (F \circ' G)^* = G^* \circ F^*.$$

- (2) *A is Arens regular if and only if $(A^{**}, \circ)$ is a $*$ -algebra.*

PROOF. (1) For all $x \in A$ and $F \in A^*$, it is easy to show that $f^* \circ x = (x^* \circ' f)^*$ and so $G \circ f^* = (f \circ' G^*)^*$. Then

$$\begin{aligned} (F \circ G)^*(f) &= \overline{F(G \circ f^*)} = \overline{F((f \circ' G^*)^*)} \\ &= F^*(f \circ' G^*) = (G^* \circ' F^*)(f). \end{aligned}$$

Hence $(F \circ G)^* = G^* \circ' F^*$. Similarly, we have $(F \circ' G)^* = G^* \circ F^*$.

(2) If A is Arens regular, then it follows from (1) that $(A^{**}, \circ)$ is a $*$ -algebra. Conversely, suppose that $(A^{**}, \circ)$ is a $*$ -algebra. Then

$$F \circ G = (F \circ G)^{**} = (G^* \circ F^*)^* = F^{**} \circ' G^{**} = F \circ' G.$$

Hence A is Arens regular and the lemma is proved.

If A is a semisimple Banach $*$ -algebra and $T = (T_1, T_2) \in M(A)$, then $T = (T_1, T_2) \rightarrow T^* = (T_1^*, T_2^*)$ is a continuous involution on $M(A)$, where $T_i^*(x) = (T_i(x^*))^*$ ($i = 1, 2$).

THEOREM 6.2. *Let A be a semisimple Banach $*$ -algebra. If A is Arens regular and A has an approximate identity $\{e_\alpha\}$ with $e_\alpha^* = e_\alpha$, then $M(A)$ is isometrically $*$ -isomorphic to M^{**} .*

PROOF. By Corollary 5.3, $M(A)$ is isometrically isomorphic to M^{**} . By Lemma 6.1, $(A^{**}, \circ)$ is a $*$ -algebra and so M^{**} is a closed $*$ -subalgebra of $(A^{**}, \circ)$. Let $T = (T_1, T_2) \in M(A)$. Then, for all $f \in A^*$, we have

$$\begin{aligned} (T_1^*)^\#(I)(f) &= \lim_\alpha (f \circ T_1^*)(e_\alpha) = \lim_\alpha f(T_1^*e_\alpha) \\ &= \lim_\alpha f((T_1(e_\alpha))^*) = \lim_\alpha \overline{(f^* \circ T_1)(e_\alpha)} = \overline{I(f^* \circ T_1)} \\ &= T_1^\#(I)(f^*) = (T_1^\#(I))^*(f). \end{aligned}$$

Hence $(T_1^*)^\#(I) = (T_1^\#(I))^*$. Therefore $\Phi(T^*) = (\Phi(T))^*$. This completes the proof of the theorem.

If A is a B^* -algebra, then A is Arens regular and A has an approximate identity. Therefore we have the following result:

COROLLARY 6.3. *If A is a B^* -algebra, then $M(A)$ is isometrically $*$ -isomorphic to M^{**} .*

REMARK. Corollary 6.3 is [8, Lemma 2.1, p. 80].

REFERENCES

1. C. A. Akemann, G. K. Pedersen, and J. Tomiyama, *Multipliers of C^* -algebras*, J. Funct. Anal. **13** (1973), 277–301.
2. R. F. Arens, *The adjoint of a bilinear operation*, Proc. Amer. Math. Soc. **2** (1951), 839–848.
3. F. F. Bonsall and J. Duncan, *Complete normed algebras*, Springer, Berlin, 1973.
4. R. C. Busby, *Double centralizers and extensions of C^* -algebras*, Trans. Amer. Math. Soc. **132** (1968), 79–99.
5. B. E. Johnson, *An introduction to the theory of centralizers*, Proc. London Math. Soc. **14** (1964), 299–320.
6. C. E. Rickart, *General theory of Banach algebras*, University Series in Higher Math., Van Nostrand, Princeton, N. J., 1960.
7. M. Tomita, *The second dual of a C^* -algebra*, Mem. Fac. Sci. Kyushu Univ. Ser. A **21** (1967), 185–193.
8. P. K. Wong, *On the Arens product and annihilator algebras*, Proc. Amer. Math. Soc. **30** (1971), 79–83.
9. ———, *On the Arens product and certain Banach algebras*, Trans. Amer. Math. Soc. **180** (1973), 437–448.
10. ———, *Arens product and the algebra of double multipliers*, Proc. Amer. Math. Soc. **94** (1985), 441–444.