*-PURE SUBGROUPS OF COMPLETELY DECOMPOSABLE ABELIAN GROUPS

LOYISO G. NONGXA

ABSTRACT. In this note we prove that

- (i) homogeneous *-pure subgroups of completely decomposable groups are completely decomposable,
- (ii) *-pure subgroups of finite rank completely decomposable groups are almost completely decomposable.

We also characterize those reduced completely decomposable groups G , with $T(G)$ satisfying the maximum condition, any *-pure subgroup of which is also completely decomposable.

In [2], a subgroup H of a torsion-free group G is said to be **-balanced* if H is balanced in G and

$$(i) \quad \langle H^*(\tau) \rangle_* = H \cap \langle G^*(\tau) \rangle_*$$

for every type τ . We shall say that H is **-pure* in G if H is pure in G and equation (i) above is satisfied for every type τ .

Throughout this note, unless otherwise specified; G will denote a completely decomposable abelian group and its extractable typeset will be denoted by $\mathcal{E}(G)$. We let $G = \bigoplus_{\tau \in \mathcal{E}(G)} G_\tau$ be a homogeneous decomposition of G and, for every $\tau \in \mathcal{E}(G)$, $\pi_\tau: G \rightarrow G_\tau$ is the projection such that $\ker \pi_\tau = \bigoplus_{\tau' \neq \tau} G_{\tau'}$.

In [4] we proved that homogeneous pure subgroups of G are also completely decomposable provided that $\mathcal{E}(G)$ is countable. The following theorem implies that the countability condition of $\mathcal{E}(G)$ can be relaxed if we consider homogeneous *-pure subgroups of G .

THEOREM 1. *Homogeneous *-pure subgroups of completely decomposable groups are completely decomposable.*

PROOF. Let H be a homogeneous *-pure subgroup of a completely decomposable group G . Let $0 \neq h \in H$ and suppose that $\text{type}_G(h) = \tau_0$. Then $h = \sum_{\tau \geq \tau_0} \pi_\tau(h)$. If $\tau_0 \notin \mathcal{E}(G)$, then

$$h = \sum_{\tau > \tau_0} \pi_\tau(h) \in H \cap G^*(\tau_0) = \langle H^*(\tau_0) \rangle_* = \{0\},$$

a contradiction. Thus $\tau_0 \in \mathcal{E}(G)$ and $\pi_{\tau_0}(h) \neq 0$. Hence $H \cong \pi_{\tau_0}(H) \subseteq G_{\tau_0}$. Since $\pi_{\tau_0}(H)$ is homogeneous of type τ_0 and is a subgroup of a homogeneous completely decomposable group G_{τ_0} , also of type τ_0 , it is completely decomposable by Lemma 86.6 in [3].

The following example illustrates that the homogeneity condition on H in the previous theorem is necessary.

Received by the editors February 5, 1986.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 20K15, 20K20, 20K27.

©1987 American Mathematical Society
0002-9939/87 \$1.00 + \$.25 per page

EXAMPLE. Let $G = G_1 \oplus G_2 \oplus G_3$ be a torsion-free group, where G_i is of rank one and type τ_i and $G_i \not\cong \mathbb{Q}$, the group of rationals, $1 \leq i \leq 3$. Assume that τ_1 and τ_2 are incomparable and $\tau_3 \geq \sup\{\tau_1, \tau_2\} > \tau_i$, $i = 1, 2$. By Lemma 1 in [1], G contains an indecomposable pure subgroup H with typeset $T(H) = \{\tau_0, \tau_1, \tau_2\}$, where $\tau_0 = \inf\{\tau_1, \tau_2\}$. Since τ_i is maximal in $T(H)$,

$$\langle H^*(\tau_i) \rangle_* = \{0\} = H \cap G_3 = H \cap G^*(\tau_i), \quad i = 1, 2,$$

and obviously

$$H = \langle H^*(\tau_0) \rangle_* = H \cap G = H \cap G^*(\tau_0).$$

Thus H is $*$ -pure in G .

Proposition 12 in [2] asserts that, if A is a torsion-free almost completely decomposable group, any balanced exact sequence

$$0 \rightarrow B \rightarrow C \rightarrow A \rightarrow 0$$

with C completely decomposable is $*$ -balanced. A special case of the following theorem implies that B is also almost completely decomposable.

THEOREM 2. *$*$ -pure subgroups of finite rank completely decomposable groups are almost completely decomposable.*

PROOF. Let H be $*$ -pure in a completely decomposable group G and define

$$\mathcal{E}(H) = \{\tau \in T(H) : \langle H^*(\tau) \rangle_* \subsetneq H(\tau)\},$$

where $T(H)$ is the typeset of H . Then, for every $\tau \in \mathcal{E}(H)$, $H(\tau) = H_\tau \oplus \langle H^*(\tau) \rangle_*$, where H_τ is a nonzero homogeneous completely decomposable group of type τ . If $\tau \in \mathcal{E}(H) \setminus \mathcal{E}(G)$, then $G^*(\tau) = G(\tau)$ which implies that $\langle H^*(\tau) \rangle_* = H \cap G^*(\tau) = H \cap G(\tau) = H(\tau)$, a contradiction. Thus $\mathcal{E}(H) \subseteq \mathcal{E}(G)$ and for every $0 \neq h \in H_\tau$, $\tau \in \mathcal{E}(H)$, $\pi_\tau(h) \neq 0$.

Let $\tau \in \mathcal{E}(H)$ and let $0 \neq h \in H_\tau$. We shall now prove that there exists a homomorphism $\psi: G \rightarrow H_\tau$ such that $\psi(h) = nh$ for some nonzero integer n . Let $\{p_1, p_2, \dots, p_k\}$ be the set of all primes such that

$$h_{p_i}^G(h) = h_{p_i}^H(h) \leq h_{p_i}^G(\pi_\tau(h)), \quad i = 1, 2, \dots, k.$$

Let $n = \prod_{i=1}^k p_i^{n(i)}$, where $n(i) = h_{p_i}^G(\pi_\tau(h)) - h_{p_i}^G(h)$, $i = 1, 2, \dots, k$.

It is easy to see that n divides $\pi_\tau(h)$ in $\langle \pi_\tau(h) \rangle_*$ and this group is a direct summand of $G_\tau = \pi_\tau(G)$. We let $\pi_\tau(h) = ng'$ and let $\theta: G \rightarrow \langle \pi_\tau(h) \rangle_*$ be a projection onto $\langle \pi_\tau(h) \rangle_*$. Then, for every prime $p \notin \{p_1, p_2, \dots, p_k\}$,

$$h_p^H(h) = h_p^G(\pi_\tau(h)) = h_p^G(g')$$

and, for every $p \in \{p_1, p_2, \dots, p_k\}$,

$$h_p^G(\pi_\tau(h)) = h_p^G(g') + h_p^G(\pi_\tau(h)) - h_p^G(h)$$

which implies that $h_p^G(g') = h_p^G(h)$. Thus, the height-sequence of g' in G is identical to the height-sequence of h in G which implies that there is an isomorphism $\psi': \langle g' \rangle_* \rightarrow \langle h \rangle_*$ with $\psi'(g') = h$. We then have

$$\psi' \theta \pi_\tau(h) = \psi' \theta (ng') = n \psi'(g') = nh \quad \text{and} \quad \psi = \psi' \theta \pi_\tau \in \text{Hom}(G, H_\tau).$$

We now prove that the regulating subgroup $\sum_{\tau \in \mathcal{E}(H)} H_\tau$ of H is completely decomposable. Let $0 = \sum_{i=1}^k h_{\tau_i}$, where $h_{\tau_i} \in H_{\tau_i}$, $\tau_i \in \mathcal{E}(H)$, $1 \leq i \leq k$. There exist homomorphisms $\psi_i: G \rightarrow H_{\tau_i}$ such that $\psi_i(h_{\tau_i}) = n_i h_{\tau_i}$ for some nonzero integers n_i , $1 \leq i \leq k$. If τ_1 , say, is minimal in $\{\tau_1, \tau_2, \dots, \tau_k\}$ we have

$$0 = \sum_{i=1}^k \psi_1(h_{\tau_i}) = \psi_1(h_{\tau_1}) = n_1 h_{\tau_1}$$

which implies that $h_{\tau_1} = 0$. It is now easy to see that $h_{\tau_i} = 0$, $1 \leq i \leq k$, which implies that $\sum_{\tau \in \mathcal{E}(H)} H_\tau = \bigoplus_{\tau \in \mathcal{E}(H)} H_\tau$. Thus H is almost completely decomposable.

Recall that a subgroup B of a torsion-free group A is said to be *regular* in A if, for every $b \in B$, $\text{type}_B(b) = \text{type}_A(b)$. We shall call a subgroup B of a torsion-free group A *strongly regular* if, for every $b \in B$, there exists a nonzero integer n and a homomorphism $\psi: A \rightarrow B$ such that $\psi(b) = nb$. This definition is an obvious generalization of the concept of strong purity introduced by K. M. Rangaswamy and S. Janakiraman. Strongly regular subgroups are regular but not necessarily pure. A subgroup that is both strongly regular and pure will be called *strongly regular pure*. Obviously, strongly regular pure subgroups are *-pure but the converse is not true. For finite rank completely decomposable groups we have the following theorem.

THEOREM 3. *A subgroup H of a finite rank completely decomposable group G is *-pure if and only if it is strongly regular pure.*

PROOF. We need only prove that, if H is *-pure in G , then H is strongly regular in G . Let $\mathcal{E}(H)$ be defined as in Theorem 2. Then, for $\tau \in \mathcal{E}(H)$, $H(\tau) = H_\tau \oplus \langle H^*(\tau) \rangle_*$ and, from the proof of Theorem 2, H_τ is strongly regular in G . We shall prove that $\bigoplus_{\tau \in \mathcal{E}(G)} H_\tau$ is strongly regular in G .

Let $0 \neq h = \sum_{i=1}^n h_{\tau_i}$, where $0 \neq h_{\tau_i} \in H_{\tau_i}$, $\tau_i \in \mathcal{E}(H)$, $1 \leq i \leq n$. There exist homomorphisms $\psi_i: G \rightarrow \langle h_{\tau_i} \rangle_*$ and nonzero integers n_i such that $\psi_i(h_{\tau_i}) = n_i h_{\tau_i}$, $1 \leq i \leq n$. Then, for every $i \in \{1, 2, \dots, n\}$

$$\psi_i(h) = n_i h_{\tau_i} + \sum_{j \neq i} \psi_i(h_{\tau_j})$$

and there exist integers m_i and m'_i such that $m_i h_{\tau_i} = m'_i (\sum_{j \neq i} \psi_i(h_{\tau_j}))$, since $\sum_{j \neq i} \psi_i(h_{\tau_j}) \in \langle h_{\tau_i} \rangle_*$. Thus

$$m'_i \psi_i(h) = m'_i n_i h_{\tau_i} + m'_i \left(\sum_{j \neq i} \psi_i(h_{\tau_j}) \right) = r_i h_{\tau_i}, \quad \text{where } r_i = m'_i n_i + m_i.$$

Thus

$$\begin{aligned} \left(\prod_{i=1}^n r_i \right) h &= \sum_{i=1}^n \left(\prod_{i=1}^n r_i \right) h_{\tau_i} = \sum_{i=1}^n \left(\prod_{j \neq i} r_j \right) r_i h_{\tau_i} \\ &= \sum_{i=1}^n r'_i m'_i \psi_i(h), \quad \text{where } r'_i = \prod_{j \neq i} r_j \\ &= \left(\sum_{i=1}^n r'_i m'_i \psi_i \right) (h). \end{aligned}$$

Hence

$$\psi = \sum_{i=1}^n r'_i m'_i \psi_i \in \text{Hom} \left(G, \bigoplus_{\tau \in \mathcal{E}(H)} H_\tau \right),$$

and $\psi(h) = rh$, where $r = \prod_{i=1}^n r_i$. This implies that $\bigoplus_{\tau \in \mathcal{E}(H)} H_\tau$ is strongly regular in G . Since $H / \bigoplus_{\tau \in \mathcal{E}(H)} H_\tau$ is finite, for every $h' \in H$ there exists an integer n' such that $n'h' \in \bigoplus_{\tau \in \mathcal{E}(H)} H_\tau$ and from the previous paragraphs there exists $\psi: G \rightarrow \bigoplus_{\tau \in \mathcal{E}(H)} H_\tau \subseteq H$ such that $\psi(n'h') = rn'h' = n'\psi(h')$ for some nonzero integer r . By torsion-freeness, $\psi(h') = rh'$ which implies that H is strongly regular in G and the proof is complete.

From the proof of the previous theorem we conclude that

- (i) finite rank $*$ -pure subgroups of separable torsion-free groups are strongly regular,
- (ii) completely decomposable $*$ -pure subgroups of completely decomposable groups are strongly regular.

It seems likely that the concepts of $*$ -purity and strongly regular purity are equivalent for separable torsion-free groups.

Finally, we wish to characterize those completely decomposable groups whose typesets satisfy the maximum condition any $*$ -pure subgroup of which is completely decomposable.

A set of types $\mathcal{T}$ is called a *tree* if, for any two incomparable types τ_1 and τ_2 in $\mathcal{T}$, there is no τ in $\mathcal{T}$ satisfying $\tau \geq \sup\{\tau_1, \tau_2\}$.

THEOREM 4. *Let G be a reduced completely decomposable group whose typeset satisfies the maximum condition. Every $*$ -pure subgroup of G is completely decomposable if and only if $\mathcal{E}(G)$ is a tree.*

PROOF. Suppose every $*$ -pure subgroup of G is completely decomposable. Let $\tau_1, \tau_2 \in \mathcal{E}(G)$ be incomparable. If there exists $\tau \in \mathcal{E}(G)$ such that $\tau \succeq \tau_i$, $i = 1, 2$, then by the example after Theorem 1, G contains a $*$ -pure subgroup which is not completely decomposable, a contradiction. This implies that $\mathcal{E}(G)$ is a tree.

Conversely, assume that $\mathcal{E}(G)$ is a tree and let H be a $*$ -pure subgroup of G . Define

$$\mathcal{E}(H) = \{\tau \in T(H) : \langle H^*(\tau) \rangle_* \subsetneq H(\tau)\}.$$

Then $\mathcal{E}(H)$ is nonempty since every maximal element of $T(H)$ belongs to $\mathcal{E}(H)$. For every $\tau \in \mathcal{E}(H)$

$$\begin{aligned} H(\tau)/\langle H^*(\tau) \rangle_* &= (H \cap G(\tau))/(H \cap G^*(\tau)) \\ &\cong \{(H \cap G(\tau)) + G^*(\tau)\}/G^*(\tau) \\ &\subseteq G(\tau)/G^*(\tau) \cong G_\tau. \end{aligned}$$

Thus $H(\tau)/\langle H^*(\tau) \rangle_*$ is a homogeneous group of type τ isomorphic to a subgroup of G_τ and therefore completely decomposable. This also implies that $\langle H^*(\tau) \rangle_*$ is balanced in $H(\tau)$ and therefore $H(\tau) = H_\tau \oplus \langle H^*(\tau) \rangle_*$ where H_τ is a homogeneous completely decomposable group of type τ . From the proof of Theorem 2, H_τ is strongly regular in G for every $\tau \in \mathcal{E}(H)$ and $\sum_{\tau \in \mathcal{E}(H)} H_\tau = \bigoplus_{\tau \in \mathcal{E}(H)} H_\tau$, a completely decomposable group. We shall prove that $H = \bigoplus_{\tau \in \mathcal{E}(H)} H_\tau$.

Let $h \in H$ be of type τ_0 and, since $T(H)$ satisfies the maximum condition, we may assume that every element of H of type greater than τ_0 belongs to $\bigoplus_{\tau \in \mathcal{E}(H)} H_\tau$, i.e. $H^*(\tau_0) \subseteq \bigoplus_{\tau \in \mathcal{E}(H)} H_\tau$. We first show that this assumption implies that $\langle H^*(\tau_0) \rangle_* \subseteq \bigoplus_{\tau \in \mathcal{E}(H)} H_\tau$. Let $h' \in \langle H^*(\tau_0) \rangle_*$ and let

$$\mathcal{E}(h') = \{\tau \in \mathcal{E}(G) : \pi_\tau(h) \neq 0\}.$$

Since $\langle H^*(\tau_0) \rangle_* = H \cap G^*(\tau_0)$, $\tau \geq \tau_0$ for every $\tau \in \mathcal{E}(h')$. Let $\mathcal{M}$ be the set of all minimal types in $\mathcal{E}(h')$ and, for every $\tau \in \mathcal{M}$, let

$$\mathcal{M}(\tau) = \{\tau' \in \mathcal{E}(h') : \tau' \geq \tau\}.$$

Since $\mathcal{E}(G)$ is a tree, $\{\mathcal{M}(\tau) : \tau \in \mathcal{M}\}$ is a partition of $\mathcal{E}(h')$ and also $\sum_{\tau \in \mathcal{M}} G(\tau) = \bigoplus_{\tau \in \mathcal{M}} G(\tau)$, a summand of G . From the proof of Theorem 3 $\bigoplus_{\tau \in \mathcal{E}(H)} H_\tau$ is strongly regular in G and, by assumption, there exists a nonzero integer n such that $nh' \in \bigoplus_{\tau \in \mathcal{E}(H)} H_\tau$. Thus, there is a homomorphism $\psi : G \rightarrow \bigoplus_{\tau \in \mathcal{E}(H)} H_\tau$ such that $\psi(h') = kh'$ for some nonzero integer k . Then

$$kh' = \sum_{\tau \in \mathcal{E}(h')} \psi \pi_\tau(h'),$$

where $\tau_0 \leq \tau \leq \text{type}_H(\psi \pi_\tau(h'))$ for every $\tau \in \mathcal{E}(h')$. By assumption, this implies that $\psi \pi_\tau(h') \in \bigoplus_{\tau \in \mathcal{E}(H)} H_\tau$. We then have

$$\begin{aligned} kh' &= \sum_{\tau \in \mathcal{E}(h')} \psi \pi_\tau(h') = \sum_{\tau \in \mathcal{M}} \sum_{\tau' \in \mathcal{M}(\tau)} \psi \pi_{\tau'}(h') \\ &\in \sum_{\tau \in \mathcal{M}} \langle \psi \pi_{\tau'}(h') : \tau' \in \mathcal{M}(\tau) \rangle_*, \end{aligned}$$

where $\langle \psi \pi_{\tau'}(h') : \tau' \in \mathcal{M}(\tau) \rangle_*$ is a pure subgroup of G for every $\tau \in \mathcal{M}$. Thus

$$\sum_{\tau \in \mathcal{M}} \langle \psi \pi_{\tau'}(h') : \tau' \in \mathcal{M}(\tau) \rangle_* = \bigoplus_{\tau \in \mathcal{M}} \langle \psi \pi_{\tau'}(h') : \tau' \in \mathcal{M}(\tau) \rangle_*$$

is pure in $\bigoplus_{\tau \in \mathcal{M}} G(\tau)$ and therefore pure in G . This implies that

$$h' \in \bigoplus_{\tau \in \mathcal{M}} \langle \psi \pi_{\tau'}(h') : \tau' \in \mathcal{M}(\tau) \rangle_* \subseteq \bigoplus_{\tau \in \mathcal{E}(H)} H_\tau$$

which proves that $\langle H^*(\tau_0) \rangle_* \subseteq \bigoplus_{\tau \in \mathcal{E}(H)} H_\tau$. But then $h \in H(\tau_0) = H_\tau \oplus \langle H^*(\tau_0) \rangle_*$ which proves that $H = \bigoplus_{\tau \in \mathcal{E}(H)} H_\tau$ is completely decomposable.

In the general case, one observes that if $\mathcal{E}(G)$ (and therefore $T(G)$) is linearly ordered, then every pure subgroup of G is $*$ -pure. One can therefore characterize completely decomposable groups any $*$ -pure subgroup of which is completely decomposable by combining Theorem 2 in [1] and Theorem 4 above.

REFERENCES

1. L. Bican, *Completely decomposable abelian groups any pure subgroup of which is completely decomposable*, Czechoslovak Math. J. **24** (1974), 176–191.
2. H. Bowman and K. M. Rangaswamy, *On special balanced subgroups of torsion-free abelian groups*, preprint.
3. L. Fuchs, *Infinite abelian groups*, Vol. 2, Academic Press, New York, 1973.
4. Loyiso G. Nongxa, *Homogeneous subgroups of completely decomposable groups*, Arch. Math. **42** (1984), 208–213.

DEPARTMENT OF MATHEMATICS AND APPLIED MATHEMATICS, UNIVERSITY OF NATAL, KING GEORGE V AVENUE, DURBAN 4001, SOUTH AFRICA