ON THE PURELY INSEPARABLE CLOSURE OF RINGS

SHIZUKA SATO

ABSTRACT. Let $K \subseteq R$ be commutative rings with identity 1. Let $D = \{D_i\}$ be a higher derivation of R . We shall prove in this paper that if K is invariant with respect to D , the purely inseparable closure $\overline{K_R}$ of K in R is invariant with respect to D and the formal power series ring $\overline{K_R}[[t]]$ is purely inseparably closed in $R[[t]]$.

1. Introduction. Let $K \subseteq R$ be commutative rings with identity 1. Let $D = \{D_i\}$ be a higher derivation of R . We shall write $D(K) \subseteq K$ if $D_i(K) \subseteq K$ for all i . We shall call R purely inseparable over K if for every element x of R there is an integer n such that $(x \otimes 1 - 1 \otimes x)^n = 0$ in $R \otimes_K R$ (cf. [1]). It is immediately proved that there exists the maximum purely inseparable subalgebra over K contained in R . We shall call this subalgebra the purely inseparable closure of K in R and we shall write $\overline{K_R}$. We shall prove that $D(K) \subseteq K$ implies $D(\overline{K_R}) \subseteq \overline{K_R}$ and the formal power series ring $\overline{K_R}[[t]]$ is purely inseparably closed in $R[[t]]$.

2. The purely inseparable closure. Let $K \subseteq R$ be commutative rings with identity 1. Let $\{K_\lambda\}_{\lambda \in \Lambda}$ be the set of purely inseparable subalgebras over K contained in R and let $\overline{K_R}$ be the subalgebra of R generated by $\{K_\lambda\}_{\lambda \in \Lambda}$ over K . Then it is evident that $\overline{K_R}$ is the maximum purely inseparable subalgebra over K in R . We shall call $\overline{K_R}$ the purely inseparable closure of K in R . If $\overline{K_R} = K$, we shall call K purely inseparably closed in R .

EXAMPLE. Let K be a field of characteristic $p > 0$ and let X, Y , and Z be three indeterminates over K . Let $R = K[X, Y, Z]/(Y^p - X) = K[x, y, z]$. Then R is a domain and $K[x, y]$ is purely inseparable over $K[x]$. Assume S is a purely inseparable subalgebra over $K[x]$ containing $K[x, y]$. An element $f \in S$ can be written in the form

$$f = \sum_{i=0}^n g_i(x, y) z^i, \quad g_i(x, y) \in K[x, y].$$

Then we have $f^{p^m} \otimes 1 - 1 \otimes f^{p^m} = 0$ for some integer m in $S \otimes_{K[x]} S$. Since $g_i(x, y)^{p^m} \in K[x]$, we have

$$\sum_{i=0}^n g_i(x, y)^{p^m} (z^{ip^m} \otimes 1 - 1 \otimes z^{ip^m}) = 0$$

and hence

$$g_i(x, y)^{p^m} = 0 \quad \text{for all } i \geq 1.$$

Received by the editors June 4, 1986.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 13B02; Secondary 13B10.

Key words and phrases. Purely inseparable algebras, higher derivations.

©1987 American Mathematical Society
 0002-9939/87 \$1.00 + \$.25 per page

Since R is a domain D , we have $g_i(x, y) = 0$ for all i and $f = g_0(x, y)$. Therefore we have $S \subseteq K[x, y]$ and it follows that the purely inseparable closure $\overline{K[x]_R}$ of $K[x]$ in R is $K[x, y]$.

A higher derivation D of R is an infinite sequence $D = \{D_0, D_1, D_2, \dots\}$ of mappings D_i of R into R such that

$$\begin{aligned} D_i(x + y) &= D_i(x) + D_i(y), \\ D_i(xy) &= \sum_{j=0}^i D_j(x) D_{i-j}(y) \quad \text{for } x, y \in R, \\ D_0(1) &= 1. \end{aligned}$$

Let $D = \{D_i\}$ be a higher derivation of R and let $e^{tD} = D_0 + tD_1 + t^2D_2 + \dots$ (t is a variable). Then e^{tD} is a homomorphism of R into $R[[t]]$.

LEMMA 1. Let $K \subseteq S \subseteq R$ be commutative rings with identity 1. Let $D = \{D_i\}$ be a higher derivation of R . Assume S is a subalgebra of R purely inseparable over K . If $D(K) \subseteq K$, then $e^{tD}(S)$ is a purely inseparable subalgebra of $R[[t]]$ over $e^{tD}(K)$.

PROOF. Let L be a reduced $e^{tD}(K)$ -algebra and let ψ, ϕ be $e^{tD}(K)$ -algebra homomorphisms of $e^{tD}(S)$ into L such that $\psi i = \phi i$ where i is the canonical injection of $e^{tD}(K)$ into $e^{tD}(S)$. We shall consider the commutative diagram:

$$\begin{array}{ccccc} R & \longrightarrow & R[[t]] & & \\ \uparrow & & \uparrow & & \\ S & \xrightarrow{e^{tD}} & e^{tD}(S) & \xrightleftharpoons[\phi]{\psi} & L \\ \uparrow i & & \uparrow i & & \\ K & \xrightarrow{e^{tD}} & e^{tD}(K) & \longrightarrow & 0 \end{array}$$

Then, since L is regarded as a K -algebra by e^{tD} , we have $\psi i e^{tD} = \phi i e^{tD}$. Hence we have $\psi i e^{tD} = \psi e^{tD} i$ and $\phi i e^{tD} = \phi e^{tD} i$. Since S is purely inseparable over K , it holds that $\psi e^{tD} = \phi e^{tD}$ and hence $\psi = \phi$. Therefore $e^{tD}(S)$ is purely inseparable over $e^{tD}(K)$.

LEMMA 2. Using the same terminology as in Lemma 1, let T be an intermediate ring as $K \subseteq T \subseteq R$. Then if S is a subalgebra of R purely inseparable over K then ST is purely inseparable over T .

PROOF. Let U be a reduced T -algebra and let ψ, ϕ be T -algebra homomorphisms of ST into U satisfying $\psi i = \phi i$. We shall consider the commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & S & \xrightarrow{j} & ST & \xrightleftharpoons[\phi]{\psi} & U \\ & & \uparrow i & & \uparrow i & & \\ 0 & \longrightarrow & K & \xrightarrow{j} & T & & \end{array}$$

(i, j are the canonical injections). From $\psi i = \phi i$ we have $\psi i j = \phi i j$ and hence $\psi j i = \phi j i$. Since S is purely inseparable over K , it holds that $\psi = \phi$ and therefore ST is purely inseparable over T .

LEMMA 3. Let $K \subseteq R$ be commutative rings with identity 1 and let f be a mapping of the product space $R[[t]] \times R[[t]]$ into $(R \otimes_K R)[[t]]$ (t is a variable) such that

$$f\left(\sum a_i t^i, \sum b_i t^i\right) = \sum_{k=0}^{\infty} \sum_{i+j=k} (a_i \otimes b_j) t^k, \quad a_i, b_j \in R.$$

Then f is bilinear and hence we have a linear mapping $\bar{f}$ of $R[[t]] \otimes_{K[[t]]} R[[t]]$ into $(R \otimes_K R)[[t]]$. By a simple calculation, this mapping is a ring homomorphism.

THEOREM 1. Let $K \subseteq R$ be commutative rings with identity 1 and let $D = \{D_i\}$ be a higher derivation of R . If $D(K) \subseteq K$, we have $D(\overline{K_R}) \subseteq \overline{K_R}$.

PROOF. Let S be the subalgebra of R generated by $\{D_i(a)\}_{a \in \overline{K_R}, i}$ over $\overline{K_R}$. By Lemmas 1 and 2, $e^{tD}(\overline{K_R})K[[t]]$ is purely inseparable over $e^{tD}(K)K[[t]] = K[[t]]$. Therefore, for any element $a \in \overline{K_R}$, there is an integer n such that

$$(e^{tD}(a) \otimes 1 - 1 \otimes e^{tD}(a))^n = 0$$

in

$$e^{tD}(\overline{K_R})K[[t]] \otimes K[[t]] e^{tD}(\overline{K_R})K[[t]].$$

Since $e^{tD}(\overline{K_R})K[[t]] \subseteq S[[t]]$, we have $e^{tD}(a) \otimes 1 - 1 \otimes e^{tD}(a)^n = 0$ in $S[[t]] \otimes_{K[[t]]} S[[t]]$ and hence

$$\bar{f}(e^{tD}(a) \otimes 1 - 1 \otimes e^{tD}(a))^n = 0$$

in $(S \otimes_K S)[[t]]$ by Lemma 3. By a simple calculation we have

$$\left[e^{tD}(a) \otimes 1 - 1 \otimes e^{tD}(a) - \sum_{i=0}^{m-1} (D_i(a) \otimes 1 - 1 \otimes D_i(a)) t^i \right]^{n2^m} = 0$$

in $(S \otimes_K S)[[t]]$ and hence $(D_m(a) \otimes 1 - 1 \otimes D_m(a))^{n2^m} = 0$ in $S \otimes_K S$. Since S is generated by $\{D_i(a)\}_{0 \leq i < \infty, a \in \overline{K_R}}$ over $\overline{K_R}$, S is purely inseparable over K . Therefore we have $\overline{K_R} = S$ and $D(\overline{K_R}) \subseteq \overline{K_R}$.

THEOREM 2. Let $K \subseteq R$ be commutative rings with identity 1. Then $\overline{K_R}[[t]]$ is purely inseparably closed in $R[[t]]$.

PROOF. Let $D = \{D_k\}$ be a sequence of mappings of $R[[t]]$ into $R[[t]]$ such that $D_k(\sum a_i t^i) = \sum a_i t^k$ for $\sum a_i t^i \in R[[t]]$, $k = 0, 1, 2, 3, \dots$. Then D is a higher derivation of $R[[t]]$ such that $D(\overline{K_R}[[t]]) \subseteq \overline{K_R}[[t]]$.

Let T be the purely inseparable closure of $\overline{K_R}[[t]]$ in $R[[t]]$ and let $\alpha = \sum a_i t^i \in T$. By Theorem 1 we have $D_k(\alpha) \in T$ and hence $(a_k \otimes 1 - 1 \otimes a_k)^n (t^{nk} \otimes 1) = 0$ in $T \otimes_{\overline{K_R}[[t]]} T$ for some integer n . Let S be a subring of R generated by coefficients of elements of T . Then we have $(a_k \otimes 1 - 1 \otimes a_k)^n (t^n \otimes 1) = 0$ in $S[[t]] \otimes_{\overline{K_R}[[t]]} S[[t]]$ and hence by Lemma 3 $(a_k \otimes 1 - 1 \otimes a_k)^n = 0$ in $S \otimes_{\overline{K_R}} S$. By the definition of S , S is purely inseparable over $\overline{K_R}$. Since $\overline{K_R}$ is purely inseparably closed in R , we have $\overline{K_R} = S$. Therefore it follows $T = \overline{K_R}[[t]]$ and hence $\overline{K_R}[[t]]$ is purely inseparably closed in $R[[t]]$.

REFERENCES

1. S. Sato, *On purely inseparable algebras and P.H.D. rings*, Trans. Amer. Math. Soc. **266** (1981), 483–498.

FACULTY OF ENGINEERING, OITA UNIVERSITY, OITA 870-11, JAPAN