A TRANSFORMATION FOR AN n -BALANCED ${}_3\Phi_2$

H. M. SRIVASTAVA¹

ABSTRACT. An interesting generalization of the familiar q -extension of the Pfaff-Saalschütz theorem is proved and is applied, for example, to derive a reduction formula for a certain double q -series. The main theorem (asserting the symmetry in n and N of a function $f(n, N)$ defined in terms of an n -balanced basic (or q -) hypergeometric ${}_3\Phi_2$ series by equation (8)) is essentially a q -extension of Sheppard's transformation.

1. Introduction and the main result. For real or complex q , $|q| < 1$, let

$$(1) \quad (\lambda; q)_\mu = \prod_{j=0}^{\infty} \left(\frac{1 - \lambda q^j}{1 - \lambda q^{\mu+j}} \right)$$

for arbitrary λ and μ , so that

$$(2) \quad (\lambda; q)_l = \begin{cases} 1, & \text{if } l = 0, \\ (1 - \lambda)(1 - \lambda q) \cdots (1 - \lambda q^{l-1}), & \forall l \in \{1, 2, 3, \dots\}, \end{cases}$$

and

$$(3) \quad (\lambda; q)_\infty = \prod_{j=0}^{\infty} (1 - \lambda q^j).$$

Denote by ${}_r\Phi_s$ a basic (or q -) hypergeometric series with r numerator and s denominator parameters (see Slater [6, Chapter 3] for details). Following Askey and Wilson [1, p. 6], we say that the q -hypergeometric series

$$(4) \quad {}_{p+1}\Phi_p \left[\begin{matrix} \alpha_0, \dots, \alpha_p; \\ \beta_1, \dots, \beta_p; \end{matrix} \middle| q, z \right] = \sum_{l=0}^{\infty} \frac{(\alpha_0; q)_l \cdots (\alpha_p; q)_l}{(\beta_1; q)_l \cdots (\beta_p; q)_l} \frac{z^l}{(q; q)_l}$$

is *balanced* if it terminates (that is, if at least one of the numerator parameters $\alpha_0, \dots, \alpha_p$ is of the form q^{-N} ($N = 0, 1, 2, \dots$)), if $z = q$, and if

$$(5) \quad \beta_1 \cdots \beta_p = q \alpha_0 \cdots \alpha_p,$$

it being understood, as usual, that no zeros appear in the denominator of (4). More generally, the q -hypergeometric series (4) is said to be *n-balanced* if this last condition (5) is replaced by

$$(6) \quad \beta_1 \cdots \beta_p = q^{n+1} \alpha_0 \cdots \alpha_p \quad (n = 0, 1, 2, \dots),$$

Received by the editors March 17, 1986 and, in revised form, May 29, 1986.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 33A30; Secondary 05A30, 11B65.

Key words and phrases. Pfaff-Saalschütz theorem, basic (or q -) hypergeometric series, n -balanced series, Sheppard's transformation, Gaussian (or q -binomial) coefficient, Jackson's sum, q -series identity, Sears's transformation, combinatorial analysis.

¹ This work was supported, in part, by the Natural Sciences and Engineering Research Council of Canada under Grant A-7353.

so that a zero-balanced q -hypergeometric series is simply called balanced.

We shall also need the Gaussian (or q -binomial) coefficient defined for all non-negative integers n and k by

$$(7) \quad \begin{bmatrix} n \\ k \end{bmatrix} = \begin{cases} 1, & \text{if } k = 0, \\ \prod_{j=1}^k \left(\frac{1-q^{n-j+1}}{1-q^j} \right), & \text{if } 1 \leq k \leq n, \\ 0, & \text{if } k > n. \end{cases}$$

The main result of the present paper is a transformation formula for an n -balanced ${}_3\Phi_2$ series, which is contained in the following

THEOREM. *Let n and N be arbitrary nonnegative integers. Then $f(n, N)$ defined in terms of an n -balanced ${}_3\Phi_2$ series by*

$$(8) \quad f(n, N) = \frac{(c; q)_N (c/ab; q)_N}{(c/a; q)_N (c/b; q)_N} {}_3\Phi_2 \left[\begin{matrix} a, b, q^{-N}; \\ cq^n, abq^{1-N}/c; \end{matrix} q, q \right]$$

is a symmetric function of n and N .

REMARK 1. The theorem can be stated in its *equivalent* form:

$$(9) \quad {}_3\Phi_2 \left[\begin{matrix} a, b, q^{-N}; \\ cq^n, abq^{1-N}/c; \end{matrix} q, q \right] = \frac{(cq^n/a; q)_{N-n} (cq^n/b; q)_{N-n}}{(cq^n; q)_{N-n} (c/ab; q)_N} \cdot \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix} \frac{(a; q)_k (b; q)_k (c/ab; q)_{n-k}}{(cq^N; q)_k} \left(\frac{c}{ab} \right)^k,$$

which, for $n = 0$, reduces immediately to Jackson's sum (cf. [3, p. 111, equation (B)]; see also [6, p. 97, equation (3.3.2.2)])

$$(10) \quad {}_3\Phi_2 \left[\begin{matrix} a, b, q^{-N}; \\ c, abq^{1-N}/c; \end{matrix} q, q \right] = \frac{(c/a; q)_N (c/b; q)_N}{(c; q)_N (c/ab; q)_N}$$

for a balanced ${}_3\Phi_2$ series.

REMARK 2. Upon replacing a, b and c by q^a, q^b and q^c , respectively, and letting $q \rightarrow 1$, formula (10) evidently yields the well-known Pfaff-Saalschütz theorem [6, p. 49, equation (2.3.1.3)]. The general result (9), on the other hand, similarly yields a ${}_3F_2$ transformation which is at least as old as Sheppard [5, p. 476, equation (18)].

2. Proof of the theorem. Our proof of the summation formula (9) is based upon (10) and the q -series identity (cf. [7, p. 229, equation (6.1)]):

$$(11) \quad \begin{aligned} \sum_{l,m=0}^{\infty} \Omega_{l+m}(\lambda; q)_l (\mu; q)_m \frac{(\mu z)^l}{(q; q)_l} \frac{z^m}{(q; q)_m} \\ = \sum_{n=0}^{\infty} \Omega_n(\lambda \mu; q)_n \frac{z^n}{(q; q)_n}, \end{aligned}$$

where $\{\Omega_n\}_{n=0}^{\infty}$ is a bounded sequence of complex numbers and the parameters λ and μ are essentially arbitrary.

Denote, for convenience, the left-hand side of the summation formula (9) by S . Applying the q -series identity (11) with, of course,

$$(12) \quad \begin{aligned} z &= q, \quad \lambda = q^{-n}, \quad \mu = q^{-N+n}, \\ \Omega_l &= \frac{(a; q)_l (b; q)_l}{(cq^n; q)_l (abq^{1-N}/c; q)_l}, \quad l \geq 0, \end{aligned}$$

we thus find that

$$\begin{aligned} S &= \sum_{l, m \geq 0} \frac{(a; q)_{l+m} (b; q)_{l+m}}{(cq^n; q)_{l+m} (abq^{1-N}/c; q)_{l+m}} \\ &\quad \cdot (q^{-n}; q)_l (q^{-N+n}; q)_m \frac{q^{(1-N+n)l}}{(q; q)_l} \frac{q^m}{(q; q)_m} \\ &= \sum_{l=0}^n \frac{(a; q)_l (b; q)_l (q^{-n}; q)_l}{(cq^n; q)_l (abq^{1-N}/c; q)_l} \frac{q^{(1-N+n)l}}{(q; q)_l} \\ &\quad \cdot {}_3\Phi_2 \left[\begin{matrix} aq^l, bq^l, q^{-N+n}; \\ cq^{n+l}, abq^{1-N+l}/c; \end{matrix} \quad q, q \right]. \end{aligned}$$

Summing this balanced ${}_3\Phi_2$ series by means of (10) with a, b, c and N replaced by aq^l, bq^l, cq^l and $N - n$, respectively, we have

$$\begin{aligned} (13) \quad S &= \frac{(cq^n/a; q)_{N-n} (cq^n/b; q)_{N-n}}{(cq^n; q)_{N-n}} \\ &\quad \cdot \sum_{l=0}^n \frac{(a; q)_l (b; q)_l (q^{-n}; q)_l}{(cq^N; q)_l (abq^{1-N}/c; q)_l (cq^{n-l}/ab; q)_{N-n}} \frac{q^{(1-N+n)l}}{(q; q)_l} \\ &= \frac{(cq^n/a; q)_{N-n} (cq^n/b; q)_{N-n}}{(cq^n; q)_{N-n} (c/ab; q)_N} \\ &\quad \cdot \sum_{l=0}^n (-1)^l q^{\frac{1}{2}l(1+2n-l)} \frac{(a; q)_l (b; q)_l (q^{-n}; q)_l}{(cq^N; q)_l} \frac{(c/ab)^l}{(q; q)_l}. \end{aligned}$$

From the definition (7) it is easily verified that

$$(14) \quad \left[\begin{matrix} n \\ k \end{matrix} \right] = \frac{(q; q)_n}{(q; q)_k (q; q)_{n-k}} = (-1)^k q^{\frac{1}{2}k(1+2n-k)} \frac{(q^{-n}; q)_k}{(q; q)_k}, \quad 0 \leq k \leq n,$$

which, when used in the last expression in (13), yields

$$S = \frac{(cq^n/a; q)_{N-n} (cq^n/b; q)_{N-n}}{(cq^n; q)_{N-n} (c/ab; q)_N} \sum_{l=0}^n \left[\begin{matrix} n \\ l \end{matrix} \right] \frac{(a; q)_l (b; q)_l (c/ab; q)_{n-l}}{(cq^N; q)_l} \left(\frac{c}{ab} \right)^l,$$

thus completing the proof of the theorem.

Alternatively, as suggested by the referee, the theorem can be proven by using Sears's transformation² [4, p. 167, equation (8.3)]

$$(15) \quad {}_4\Phi_3 \left[\begin{matrix} \alpha, \beta, \gamma, q^{-m}; \\ \lambda, \mu, \nu; \end{matrix} \quad q, q \right] = \frac{(\mu/\alpha; q)_m (\lambda\mu/\beta\gamma; q)_m}{(\mu; q)_m (\lambda\mu/\alpha\beta\gamma; q)_m} \cdot {}_4\Phi_3 \left[\begin{matrix} \alpha, \lambda/\beta, \lambda/\gamma, q^{-m}; \\ \lambda, \alpha q^{1-m}/\mu, \alpha q^{1-m}/\nu; \end{matrix} \quad q, q \right],$$

which holds true when each ${}_4\Phi_3$ series is balanced, that is, when m is a nonnegative integer and (cf. equation (5)) $\lambda\mu\nu = \alpha\beta\gamma q^{1-m}$. Letting $\gamma, \nu \rightarrow 0$, (15) readily yields the transformation

$$(16) \quad {}_3\Phi_2 \left[\begin{matrix} \alpha, \beta, q^{-m}; \\ \lambda, \mu; \end{matrix} \quad q, q \right] = \frac{(\mu/\alpha; q)_m}{(\mu; q)_m} \alpha^m {}_3\Phi_2 \left[\begin{matrix} \alpha, \lambda/\beta, q^{-m}; \\ \lambda, \alpha q^{1-m}/\mu; \end{matrix} \quad q, \frac{\beta q}{\mu} \right]$$

between two terminating ${}_3\Phi_2$ series.

Making use of (16), we observe that

$$(17) \quad \frac{(c; q)_N (c/ab; q)_N}{(c/a; q)_N (c/b; q)_N} {}_3\Phi_2 \left[\begin{matrix} a, b, q^{-N}; \\ cq^n, abq^{1-N}/c; \end{matrix} \quad q, q \right] \\ = \frac{(c; q)_\infty (c/ab; q)_\infty}{(c/a; q)_\infty (c/b; q)_\infty} {}_3\Phi_2 \left[\begin{matrix} a, b, cq^{n+N}; \\ cq^n, cq^N; \end{matrix} \quad q, \frac{c}{ab} \right]$$

when $b = q^{-m}$ ($m = 0, 1, 2, \dots$). Obviously, the right-hand side of (17) is symmetric in n and N . Thus the left-hand side of (17) is symmetric in n and N when $b = q^{-m}$. Since both ${}_3\Phi_2$ series in (17) are polynomials, (17) holds true for infinitely many values of b , and hence for all b , as long as no problems arise from division by zero.

A comparison between (17) and the definition (8) evidently completes the proof of the theorem.

3. Applications. Our assertion that $f(n, N)$ defined by (8) is a symmetric function of n and N might lead to a number of useful consequences.

For example, just as the Pfaff-Saalschütz theorem and its q -analogue (10), the general summation formula (9) has great potential for applications in various areas of combinatorial analysis (see, for details, Takács [9] and Goulden [2], and the references cited in these works). We choose to record here the following consequence of our theorem, which can indeed be proved by comparing coefficients of like powers of z on both sides, using the summation formula (9):

²See also Askey and Wilson [1, pp. 4-7] for a systematic account of Sears's transformation (15) and of some of its numerous interesting consequences.

COROLLARY. For every bounded sequence $\{\Omega_n\}_{n=0}^\infty$ of complex numbers, and for complex parameters α, β, γ and q , $|q| < 1$,

$$\begin{aligned}
 & \sum_{l,m=0}^{\infty} \Omega_{l+m} \frac{(\alpha; q)_l (\beta; q)_l (\gamma/\alpha\beta; q)_m}{(\gamma q^n; q)_l} \frac{(\gamma z/\alpha\beta)^l}{(q; q)_l} \frac{z^m}{(q; q)_m} \\
 (18) \quad &= \frac{(\gamma; q)_n}{(\gamma/\alpha; q)_n (\gamma/\beta; q)_n} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix} (\alpha; q)_k (\beta; q)_k (\gamma/\alpha\beta; q)_{n-k} \left(\frac{\gamma}{\alpha\beta} \right)^k \\
 & \cdot \sum_{N=0}^{\infty} \Omega_N \frac{(\gamma/\alpha; q)_N (\gamma/\beta; q)_N}{(\gamma; q)_{N+k}} \frac{z^N}{(q; q)_N},
 \end{aligned}$$

provided that each series involved converges absolutely.

REMARK 3. For $n = 0$, (18) yields a q -analogue of a known series identity [8, p. 313, equation (143) with $y = 0$].

REFERENCES

1. R. Askey and J. Wilson, *Some basic hypergeometric orthogonal polynomials that generalize Jacobi polynomials*, Mem. Amer. Math. Soc., Vol. 54, No. 319 (1985).
2. I. P. Goulden, *A bijective proof of the q -Saalschütz theorem*, Discrete Math. **57** (1985), 39–44.
3. F. H. Jackson, *Summation of q -hypergeometric series*, Messenger Math. **50** (1921), 101–112.
4. D. B. Sears, *On the transformation theory of basic hypergeometric functions*, Proc. London Math. Soc. (2) **53** (1951), 158–180.
5. W. F. Sheppard, *Summation of the coefficients of some terminating hypergeometric series*, Proc. London Math. Soc. (2) **10** (1912), 469–478.
6. L. J. Slater, *Generalized hypergeometric functions*, Cambridge Univ. Press, Cambridge, London and New York, 1966.
7. H. M. Srivastava and V. K. Jain, *q -Series identities and reducibility of basic double hypergeometric functions*, Canad. J. Math. **38** (1986), 215–231.
8. H. M. Srivastava and P. W. Karlsson, *Multiple Gaussian hypergeometric series*, Halsted Press (Ellis Horwood Limited, Chichester); Wiley, New York, Chichester, Brisbane, and Toronto, 1985.
9. L. Takács, *On an identity of Shih-Chieh Chu*, Acta Sci. Math. (Szeged) **34** (1973), 383–391.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF VICTORIA, VICTORIA, BRITISH COLUMBIA V8W 2Y2, CANADA