A SIMPLE PROOF OF JACOBI'S FOUR-SQUARE THEOREM

M. D. HIRSCHHORN

(Communicated by Larry J. Goldstein)

ABSTRACT. Jacobi's four-square theorem, which gives the number of representations of a positive integer as a sum of four squares, is shown to follow simply from the triple-product identity.

1. Recently [2], I showed how one can obtain Jacobi's two-square theorem from the triple-product identity. In this note I show that the triple-product identity also gives Jacobi's four-square theorem:

THEOREM 1. *The number $r_4(n)$ of representations of the positive integer n as a sum of four squares is given by*

$$r_4(n) = 8 \sum_{d|n, 4 \nmid d} d.$$

2. The triple product identity is

$$\prod_{n \geq 1} (1 + ax^{2n-1})(1 + a^{-1}x^{2n-1})(1 - x^{2n}) = \sum_{-\infty}^{\infty} a^n x^{n^2}.$$

It follows easily (loc. cit.) that

$$(1) \quad (a - a^{-1}) \prod_{n \geq 1} (1 - a^2 x^n)(1 - a^{-2} x^n)(1 - x^n) = \sum_{-\infty}^{\infty} (-1)^n a^{2n+1} x^{(n^2+n)/2}.$$

Differentiate (1) with respect to a , put $a = 1$, divide by 2 and we obtain the identity

$$(2) \quad \prod_{n \geq 1} (1 - x^n)^3 = \frac{1}{2} \sum_{-\infty}^{\infty} (-1)^n (2n+1) x^{(n^2+n)/2}.$$

This is a celebrated identity of Jacobi [1, Theorem 357], and can be considered our starting point.

Squaring (2) gives

$$\prod_{n \geq 1} (1 - x^n)^6 = \frac{1}{4} \sum_{m, n = -\infty}^{\infty} (-1)^{m+n} (2m+1)(2n+1) x^{(m^2+n^2+m+n)/2}.$$

Received by the editors April 15, 1986 and, in revised form, July 15, 1986.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 11E25, 11P05.

©1987 American Mathematical Society
0002-9939/87 \$1.00 + \$.25 per page

Now split the sum on the right into two, according as $m + n$ is even or odd, to obtain

$$\prod_{n \geq 1} (1 - x^n)^6 = \frac{1}{4} \left\{ \sum_{m \equiv n \pmod{2}} (2m+1)(2n+1)x^{(m^2+n^2+m+n)/2} - \sum_{m \not\equiv n \pmod{2}} (2m+1)(2n+1)x^{(m^2+n^2+m+n)/2} \right\}.$$

In the first sum, set $r = \frac{1}{2}(m+n)$, $s = \frac{1}{2}(m-n)$, and in the second set $r = \frac{1}{2}(m-n-1)$, $s = \frac{1}{2}(m+n+1)$, and, remarkably, the two sums coalesce to give

$$\prod_{n \geq 1} (1 - x^n)^6 = \frac{1}{2} \sum_{r, s = -\infty}^{\infty} ((2r+1)^2 - (2s)^2) x^{r^2+s^2+r}.$$

Once again splitting the sum, we obtain

$$\begin{aligned} \prod_{n \geq 1} (1 - x^n)^6 &= \frac{1}{2} \left\{ \sum_{s=-\infty}^{\infty} x^{s^2} \sum_{r=-\infty}^{\infty} (2r+1)^2 x^{r^2+r} - \sum_{r=-\infty}^{\infty} x^{r^2+r} \sum_{s=-\infty}^{\infty} (2s)^2 x^{s^2} \right\} \\ &= \frac{1}{2} \left\{ \sum_{s=-\infty}^{\infty} x^{s^2} \times \left(1 + 4x \frac{d}{dx}\right) \sum_{r=-\infty}^{\infty} x^{r^2+r} - \sum_{r=-\infty}^{\infty} x^{r^2+r} \times 4x \frac{d}{dx} \sum_{s=-\infty}^{\infty} x^{s^2} \right\}. \end{aligned}$$

Making use of the triple-product identity, we obtain

$$\begin{aligned} \prod_{n \geq 1} (1 - x^n)^6 &= \frac{1}{2} \left\{ \prod_{n \geq 1} (1 + x^{2n-1})^2 (1 - x^{2n}) \right. \\ &\quad \times \left(1 + 4x \frac{d}{dx}\right) 2 \prod_{n \geq 1} (1 + x^{2n})^2 (1 - x^{2n}) \\ &\quad - 2 \prod_{n \geq 1} (1 + x^{2n})^2 (1 - x^{2n}) \\ &\quad \left. \times 4x \frac{d}{dx} \prod_{n \geq 1} (1 + x^{2n-1})^2 (1 - x^{2n}) \right\}. \end{aligned}$$

Employing the product rule to evaluate the derivatives, we find

$$\begin{aligned} \prod_{n \geq 1} (1 - x^n)^6 &= \prod_{n \geq 1} (1 + x^{2n-1})^2 (1 - x^{2n}) (1 + x^{2n})^2 (1 - x^{2n}) \\ &\quad \times \left\{ 1 + 8 \sum_{n \geq 1} \frac{2nx^{2n}}{1 + x^{2n}} - 4 \sum_{n \geq 1} \frac{2nx^{2n}}{1 - x^{2n}} \right\} \\ &\quad - \prod_{n \geq 1} (1 + x^{2n})^2 (1 - x^{2n}) (1 + x^{2n-1})^2 (1 - x^{2n}) \\ &\quad \times \left\{ 8 \sum_{n \geq 1} \frac{(2n-1)x^{2n-1}}{1 + x^{2n-1}} - 4 \sum_{n \geq 1} \frac{2nx^{2n}}{1 - x^{2n}} \right\}, \end{aligned}$$

or,

$$\begin{aligned} \prod_{n \geq 1} (1 - x^n)^6 &= \prod_{n \geq 1} (1 + x^{2n-1})^2 (1 + x^{2n})^2 (1 - x^{2n})^2 \\ &\quad \times \left\{ 1 - 8 \sum_{n \geq 1} \left(\frac{(2n-1)x^{2n-1}}{1 + x^{2n-1}} - \frac{2nx^{2n}}{1 + x^{2n}} \right) \right\}. \end{aligned}$$

Dividing both sides by

$$\begin{aligned}\prod_{n \geq 1} (1 + x^n)^4 (1 - x^n)^2 &= \prod_{n \geq 1} (1 + x^n)^2 (1 - x^{2n})^2 \\ &= \prod_{n \geq 1} (1 + x^{2n-1})^2 (1 + x^{2n})^2 (1 - x^{2n})^2,\end{aligned}$$

we obtain

$$\prod_{n \geq 1} \left(\frac{1 - x^n}{1 + x^n} \right)^4 = 1 - 8 \sum_{n \geq 1} \left(\frac{(2n-1)x^{2n-1}}{1 + x^{2n-1}} - \frac{2nx^{2n}}{1 + x^{2n}} \right).$$

Now, it is a simple consequence of the triple-product identity that

$$\prod_{n \geq 1} \left(\frac{1 - x^n}{1 + x^n} \right) = \sum_{n=-\infty}^{\infty} (-1)^n x^{n^2},$$

so we have

$$\left(\sum_{n=-\infty}^{\infty} (-1)^n x^{n^2} \right)^4 = 1 - 8 \sum_{n \geq 1} \left(\frac{(2n-1)x^{2n-1}}{1 + x^{2n-1}} - \frac{2nx^{2n}}{1 + x^{2n}} \right).$$

Putting $-x$ for x , we obtain

$$\begin{aligned}\left(\sum_{n=-\infty}^{\infty} x^{n^2} \right)^4 &= 1 + 8 \sum_{n \geq 1} \left(\frac{(2n-1)x^{2n-1}}{1 - x^{2n-1}} + \frac{2nx^{2n}}{1 + x^{2n}} \right) \\ &= 1 + 8 \sum_{n \geq 1} \left(\frac{(2n-1)x^{2n-1}}{1 - x^{2n-1}} + \frac{2nx^{2n}}{1 - x^{2n}} \right) - 8 \sum_{n \geq 1} \left(\frac{2nx^{2n}}{1 - x^{2n}} - \frac{2nx^{2n}}{1 + x^{2n}} \right) \\ &= 1 + 8 \sum_{n \geq 1} \frac{nx^n}{1 - x^n} - 8 \sum_{n \geq 1} \frac{4nx^{4n}}{1 - x^{4n}},\end{aligned}$$

or,

$$\left(\sum_{n=-\infty}^{\infty} x^{n^2} \right)^4 = 1 + 8 \sum_{4 \nmid n} \frac{nx^n}{1 - x^n}$$

from which Theorem 1 follows directly.

REFERENCES

1. G. H. Hardy and E. M. Wright, *An introduction to the theory of numbers*, 4th ed., Clarendon Press, Oxford.
2. Michael D. Hirschhorn, *A simple proof of Jacobi's two-square theorem*, Amer. Math. Monthly **92** (1985), 579–580.

SCHOOL OF MATHEMATICS, UNIVERSITY OF NEW SOUTH WALES, P. O. BOX 1, KENSINGTON, NEW SOUTH WALES, AUSTRALIA 2033