RELATION BETWEEN RIGHT AND LEFT INVOLUTIONS OF A HILBERT ALGEBRA

P. P. SAWOROTNOW

(Communicated by John B. Conway)

ABSTRACT. Existence of a densely defined right involution in a Hilbert algebra implies existence of a left involution.

H^* -algebras were introduced by W. Ambrose [1] to characterize Hilbert-Schmidt operators. This notion was generalized by M. F. Smiley [7], who showed that the structure theorems are valid also for a right H^* -algebra (a Hilbert algebra whose involution $x \rightarrow x^r$ satisfies the condition " $(yx, z) = (y, zx^r)$ " but not the condition " $(xy, z) = (y, x^r z)$ "). (Hilbert algebra here is a Banach algebra with a Hilbert space norm.) The author showed in [5, Theorem 2] that a proper right H^* -algebra is also a left H^* -algebra, i.e. it also has another involution $x \rightarrow x^l$ (a *left* involution) which satisfied the condition " $(xy, z) = (y, x^l z)$."

In this paper we shall show that the same is true also for the case when the involution $x \rightarrow x^r$ (the *right* involution) is defined on a dense subset only.

DEFINITION. Let A be a Hilbert algebra (A is a Banach algebra with a scalar product $(\ , \)$ such that $(x, x) = \|x\|^2$). We shall say that A is a *weak right H^* -algebra* if there is a dense subset D_r of A with the property that for each $x \in D_r$ there is some x^r such that $(yx, z) = (y, zx^r)$ for all $y, z \in A$. We define *weak left H^* -algebra* in a similar fashion.

The algebra A is said to be *proper* if each x^r is unique, i.e. A has a right involution $x \rightarrow x^r$, defined on a dense subset. Note that A is proper if $r(A) = \{u \in A: Au = (0)\}$ consists of zero alone. Algebra A in Example 2, p. 54, of [5] is an example of weak right (as well as left) H^* -algebra. Also it is easy to show that each weak right H^* -algebra is a right complemented algebra.

THEOREM. *Each proper weak right H^* -algebra A is a proper weak left H^* -algebra.*

PROOF. Note that the involution $x \rightarrow x^r$ of A , defined on a dense subset D_r , is closable, i.e. the closure of its graph is a graph of some mapping: it is easy to show that if $x_n \in D_r$, $x_n \rightarrow 0$ and $x_n^r \rightarrow y \in A$, then $y = 0$ (see end of section 8 in §5, Chapter I of [3]).

Now replace the scalar product $(\ , \)$ of A and the multiplication λx of members x of A and complex number λ by $[\ , \]$ and $\lambda \circ x$ respectively, where these new

Received by the editors November 3, 1986.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 46K15; Secondary 46H20, 47C10.

Key words and phrases. Hilbert algebra, H^* -algebra, right H^* -algebra, left H^* -algebra, weak right H^* -algebra, weak left H^* -algebra, right involution, left involution.

products are defined as follows:

$$\begin{aligned} [x, y] &= (y, x), & x, y &\in A, \\ \lambda \circ x &= \bar{\lambda}x, & \lambda &\text{ is a complex number.} \end{aligned}$$

Let A' denote the algebra consisting of members of A but with these new operations. Note that A' is also a weak right H^* -algebra.

Let $T': A \rightarrow A$ be a mapping defined by $T'x = x^r$ and let T be the closed extension of T' (we remarked above that the map $x \rightarrow x^r$ is closable). It follows from II in section 9 of §5 in Chapter I of [3], that T has an adjoint T^* defined on a dense subset D_l of A' . We define $x^l = T^*x$ for each $x \in D_l$. It follows that $(x^l, y) = (T^*x, y) = [x, Ty] = (y^r, x)$ for all $y \in D_r$, from which we see that $(x^ly, z) = (x^l, zy^r) = ((zy^r)^r, x) = (yz^r, x) = (y, xz)$ for all $y, z \in D_r$ (and each $x \in D_l$).

We leave it to the reader to deduce that $(xy, z) = (y, x^lz)$ for all $y, z \in A$.

To conclude the paper it is appropriate to comment that, in the presence of semisimplicity of the algebra A , the above Theorem follows from Theorem 2 of [6]. However, semisimplicity of a proper weak right H^* -algebra would be much more difficult to establish (if it is true at all) than in the case of a right H^* -algebra. In the latter case it follows from the fact that each ideal (whether closed or not) contains a (right) selfadjoint element (it is easy to show that a selfadjoint element is not a generalized nilpotent).

REFERENCES

1. W. Ambrose, *Structure theorems for a special class of Banach algebras*, Trans. Amer. Math. Soc. **57** (1945), 365-386.
2. L. H. Loomis, *An introduction to abstract harmonic analysis*, Van Nostrand, 1953.
3. M. A. Naimark, *Normed rings*, "Nauka", Moscow, 1968; English transl., *Normed algebras*, Wolters-Noordhoff, Groningen, 1972.
4. Frigyes Riesz and Bela Sz.-Nagy, *Functional analysis*, Ungar, New York, 1955.
5. P. P. Saworotnow, *On a generalization of the notion of H^* -algebra*, Proc. Amer. Math. Soc. **8** (1957), 49-55.
6. —, *On the imbedding of a right complemented algebra into Ambrose's H^* -algebra*, Proc. Amer. Math. Soc. **8** (1957), 56-62.
7. M. F. Smiley, *Right H^* -algebra*, Proc. Amer. Math. Soc. **4** (1953), 1-4.

DEPARTMENT OF MATHEMATICS, THE CATHOLIC UNIVERSITY OF AMERICA, WASHINGTON, D.C. 20064