UNIQUE SOLVABILITY OF AN EXTENDED STIELTJES MOMENT PROBLEM

OLAV NJÅSTAD

(Communicated by Paul S. Muhly)

ABSTRACT. Let $a_1, \dots, a_p$ be given real numbers ordered by size, and let $[\alpha, \beta]$ be a real interval disjoint from the set $\{a_1, \dots, a_p\}$. Let $\{c_j^{(i)}: j = 1, 2, \dots\}$, $i = 1, \dots, p$, be sequences of real numbers and c_0 be a real number. The extended Stieltjes moment problem is to find a distribution function ψ with all its points of increase in $[\alpha, \beta]$ such that

$$\int_{\alpha}^{\beta} d\psi(t) = c_0, \quad \int_{\alpha}^{\beta} \frac{d\psi(t)}{(t - a_i)^j} = c_j^{(i)}, \quad i = 1, \dots, p, \quad j = 1, 2, \dots$$

Necessary and sufficient conditions for the existence of a unique solution of the problem are given. Orthogonal R -functions and Gaussian quadrature formulas play important roles in the proof.

1. Introduction. By a distribution function we shall mean a real-valued, bounded, nondecreasing function $\psi(t)$ with an infinite number of points of increase. By its Stieltjes transform we mean the function $\hat{\psi}(z) = \int_{-\infty}^{\infty} d\psi(t)/(t - z)$. If all the points of increase of ψ lie in an interval I , then $\hat{\psi}(z)$ is an analytic function in $C - \bar{I}$.

Let $a_1, a_2, \dots, a_p$ be distinct real numbers, ordered by size. We shall call the real interval $[\alpha, \beta]$ a *Stieltjes interval* for the point set $\{a_1, \dots, a_p\}$ if $(\alpha, \beta) \cap \{a_1, \dots, a_p\} = \emptyset$. (Intervals of the form $(-\infty, \beta]$ and $[\alpha, \infty)$ are allowed.)

Let $\{c_j^{(i)}: j = 1, 2, \dots\}$, $i = 1, \dots, p$, be given sequences of real numbers and c_0 be a real number. The *extended Stieltjes moment problem* (ESMP) is defined as follows: Find a distribution function $\psi(t)$ with all its points of increase in a given Stieltjes interval $[\alpha, \beta]$ such that

$$\int_{\alpha}^{\beta} d\psi(t) = c_0, \quad \int_{\alpha}^{\beta} \frac{d\psi(t)}{(t - a_i)^j} = c_j^{(i)}, \quad j = 1, 2, \dots, \quad i = 1, \dots, p.$$

In [3, 4] we studied the *extended Hamburger moment problem* (EHMP), which consists of finding a distribution function with the same properties as above, except the requirements that the points of increase lie in a given interval. Conditions for existence of a solution of the ESMP were briefly discussed in [5]. In this paper we give necessary and sufficient conditions for the existence of a unique solution of the problem. Since the tools used in proving the existence of solutions also are involved in the discussion of the uniqueness question, we give a very brief sketch of the existence proof here.

Received by the editors September 8, 1986.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 30E05; Secondary 42C05.

Key words and phrases. Moment problems, orthogonal R -functions.

©1988 American Mathematical Society
0002-9939/88 \$1.00 + \$.25 per page

The treatment of the uniqueness problem is related to the treatment of the uniqueness problem for the strong Stieltjes moment problem, given in terms of continued fractions in [1, 2]. For the classical Stieltjes moment problem, see e.g. [6, 7].

2. Orthogonal R -functions. Let $\mathcal{R}$ denote the linear space consisting of all functions of the form

$$(2.1) \quad R(z) = \alpha_0 + \sum_{i=1}^p \sum_{j=1}^{N_i} \frac{\alpha_{ij}}{(z - a_i)^j}, \quad \alpha_0, \alpha_{ij} \in \mathbb{C}.$$

Elements of $\mathcal{R}$ are called R -functions. We denote by $\mathcal{R}_{\mathbf{R}}$ the real space of all R -functions with real coefficients. A function R belongs to $\mathcal{R}$ iff it can be written in the form $R(z) = P(z)/Q(z)$, where Q is a polynomial with all its zeros among the points $a_1, \dots, a_p$, and where P is a polynomial with $\deg P \leq \deg Q$. The spaces $\mathcal{R}$ and $\mathcal{R}_{\mathbf{R}}$ are closed under multiplication.

Every natural number n has a unique decomposition $n = q_n p + r_n$, $1 \leq r_n \leq p$. We write $q = q_n$, $r = r_n$. We denote by $\mathcal{R}_n$ the space of R -functions of the form

$$(2.2) \quad R(z) = \alpha_0 + \sum_{i=1}^r \sum_{j=1}^{q+1} \frac{\alpha_{ij}}{(z - a_i)^j} + \sum_{i=r+1}^p \sum_{j=1}^q \frac{\alpha_{ij}}{(z - a_i)^j}.$$

Let Φ denote the linear functional defined on $\mathcal{R}$ by

$$(2.3) \quad \Phi \left(\alpha_0 + \sum_{i=1}^p \sum_{j=1}^{N_i} \frac{\alpha_{ij}}{(z - a_i)^j} \right) = \alpha_0 c_0 + \sum_{i=1}^p \sum_{j=1}^{N_i} \alpha_{ij} c_j^{(i)}.$$

We shall say that Φ is *positive on an interval* $[a, b]$ if $\Phi(R) > 0$ for every $R \in \mathcal{R}$ such that $R(t) \geq 0$ and $R(t) \not\equiv 0$ for $t \in (a, b)$. It follows immediately that a necessary condition for the ESMP for a given Stieltjes interval $[\alpha, \beta]$ to have a solution is that Φ is positive on $[\alpha, \beta]$. We shall in the following assume that this condition is satisfied.

If the functional Φ is positive on some interval $[\alpha, \beta]$, it is positive on $(-\infty, \infty)$, and therefore gives rise to an inner product $\langle \cdot, \cdot \rangle$ defined by $\langle R, S \rangle = \Phi(R \cdot S)$. By applying the Gram-Schmidt procedure to the sequence

$$\left\{ 1, \frac{1}{(z - a_1)}, \dots, \frac{1}{(z - a_p)}, \frac{1}{(z - a_1)^2}, \dots, \frac{1}{(z - a_p)^2}, \frac{1}{(z - a_1)^3}, \dots \right\}$$

we obtain an orthonormal sequence $\{Q_n\}$ of R -functions. We note that $\langle Q_n, R \rangle = 0$ for every $R \in \mathcal{R}_{n-1}$. Furthermore Q_n may be written as $Q_n(z) = V_n(z)/N_n(z)$, where

$$(2.4) \quad N_n(z) = (z - a_1)^{q+1} \dots (z - a_r)^{q+1} (z - a_{r+1})^q \dots (z - a_p)^q,$$

and $V_n(z)$ is a polynomial of degree at most n .

3. Existence of solutions. When Φ is positive on a Stieltjes interval $[\alpha, \beta]$, the polynomial $V_n(z)$ is of degree n , and has n real simple zeros in (α, β) (cf. [5], see also [3]). This can be seen as follows: Let $t_1, \dots, t_\lambda$ be all the zeros of odd order of $V_n(z)$ in (α, β) . If $\lambda < n$, then the function

$$T_n(z) = \frac{(z - t_1) \dots (z - t_\lambda)(z - a_r)}{N_n(z)}$$

belongs to $\mathcal{R}_{n-1}$, hence $\Phi(Q_n T_n) = 0$. On the other hand the function

$$T_n(z)Q_n(z) = \frac{(z - t_1) \cdots (z - t_\lambda)V_n(z)(z - a_r)}{N_n(z)^2}$$

has a fixed sign in (α, β) , since $(z - a_r)$ does not change sign in (α, β) . Thus $\Phi(Q_n T_n) \neq 0$, which is a contradiction. Consequently $\lambda = n$, and the result follows.

With the zeros $t_1^{(n)}, \dots, t_n^{(n)}$ of $V_n(z)$ (or equivalently of $Q_n(z)$) there is associated a Gaussian quadrature formula. There exist positive weights $\lambda_{n,1}, \dots, \lambda_{n,n}$ such that

$$(3.1) \quad \Phi(R) = \sum_{k=1}^n \lambda_{n,k} R(t_k^{(n)})$$

for every R of the form (2.1), where $N_i \leq 2q + 2$ for $i < r$, $N_i \leq 2q$ for $i > r$, and $N_r \leq 2q + 1$ (see [3, Theorem 4.2, Proposition 4.4; 4, Theorems 2.6, 2.7]). For given i, j we may then write

$$(3.2) \quad c_j^{(i)} = \Phi \left(\frac{1}{(z - a_i)^j} \right) = \sum_{k=1}^n \lambda_{n,k} \frac{1}{(t_k^{(n)} - a_i)^j}$$

for sufficiently large n . Thus we may also write $c_j^{(i)} = \int_\alpha^\beta d\phi_n(t)/(t - a_i)^j$, where the nondecreasing, uniformly bounded functions $\phi_n(t)$ are defined by $\phi_n(t) = \sum \{\lambda_{n,k} : t_k^{(n)} \leq t\}$. Application of Helly's selection and convergence theorems show that the sequence $\{\phi_n(t)\}$ contains at least one subsequence converging to a distribution function ϕ , and the limit function ϕ of every such convergent subsequence has all its points of increase in (α, β) and satisfies

$$\int_\alpha^\beta d\phi(t) = c_0, \quad \int_\alpha^\beta \frac{d\phi(t)}{(t - a_i)^j} = c_j^{(i)}, \quad j = 1, 2, \dots, i = 1, \dots, p.$$

(For more details, see [3].) Thus we have seen

THEOREM 1. *A necessary and sufficient condition for the ESMP on the given Stieltjes interval $[\alpha, \beta]$ to have a solution is that Φ is positive on $[\alpha, \beta]$.*

4. Uniqueness of solution. We define the R -function P_n associated with Q_n by

$$(4.1) \quad P_n(z) = \Phi_t \left(\frac{Q_n(t) - Q_n(z)}{t - z} \right).$$

(The subscript t indicates that the functional operates on its argument as a function of t .) The function $P_n(z)$ can be written in the form $P_n(z) = U_n(z)/N_n(z)$ where $U_n(z)$ is a polynomial of degree at most $n - 1$.

From the Gaussian quadrature formula it follows that we may write

$$(4.2) \quad P_n(z) = - \sum_{k=1}^n \lambda_{n,k} \frac{Q_n(z)}{t_k^{(n)} - z} = -Q_n(z) \int_\alpha^\beta \frac{d\phi_n(t)}{t - z},$$

which means $P_n(z)/Q_n(z) = -\hat{\phi}_n(z)$.

If the ESMP has a unique solution $\psi(t)$, then $\{\phi_n(t)\}$ converges to this solution, since otherwise convergent subsequences would produce distinct solutions. It follows that $\{\hat{\phi}_n(z)\}$ converges to $\hat{\psi}(z)$ for all $z \in \mathbf{C} - [\alpha, \beta]$. The sequence $\{P_n(z)/Q_n(z)\}$ is convergent for the same values of z .

We shall now show that if $\{P_n(z)/Q_n(z)\}$ converges on an interval of the form (a_ρ, a_σ) , where $(a_\rho, a_\sigma) \cap [\alpha, \beta] = \emptyset$, then the ESMP has a unique solution.

We note that the function

$$\begin{aligned} f(t) &= \frac{1}{(t-z)} \left[\frac{1}{N_{n-1}(t)} - \frac{1}{N_{n-1}(z)} \right] \\ &= \frac{N_{n-1}(z) - N_{n-1}(t)}{(t-z)N_{n-1}(t)N_{n-1}(z)} \end{aligned}$$

belongs to $\mathcal{R}_{n-1}$, and thus $\Phi(Q_n \cdot f) = 0$. This implies

$$\begin{aligned} &\Phi_t \left(\frac{1}{t-z} \left[\frac{Q_n(t)}{N_{n-1}(t)} - \frac{Q_n(z)}{N_{n-1}(z)} \right] \right) \\ &= \Phi_t \left(\frac{1}{t-z} \left[\frac{Q_n(t) - Q_n(z)}{N_{n-1}(z)} \right] \right) + \Phi_t(Q_n(t)f(t)) \\ &= \frac{1}{N_{n-1}(z)} P_n(z). \end{aligned}$$

Hence

$$(4.3) \quad P_n(z) = N_{n-1}(z) \Phi_t \left(\frac{1}{t-z} \left[\frac{Q_n(t)}{N_{n-1}(t)} - \frac{Q_n(z)}{N_{n-1}(z)} \right] \right).$$

We also note that the function

$$g(t) = \frac{V_n(t) - V_n(z)}{(t-z)N_{n-1}(t)}$$

belongs to $\mathcal{R}_{n-1}$, and thus

$$(4.4) \quad \Phi \left(Q_n(t) \frac{V_n(t) - V_n(z)}{(t-z)N_{n-1}(t)} \right) = 0.$$

Let ϕ be an arbitrary solution of the EHMP. From (4.3) we obtain

$$(4.5) \quad P_n(z) = N_{n-1}(z) \int_{-\infty}^{\infty} \frac{Q_n(t)}{(t-z)N_{n-1}(t)} d\phi(t) - Q_n(z)\hat{\phi}(z).$$

Similarly from (4.4) we get (noting that $N_n(t) = (t - a_r)N_{n-1}(t)$)

$$(4.6) \quad \int_{-\infty}^{\infty} \frac{Q_n(t)}{(t-z)N_{n-1}(t)} d\phi(t) = \frac{1}{V_n(z)} \int_{-\infty}^{\infty} \frac{Q_n(t)^2(t - a_r)}{t-z} d\phi(t).$$

Substitution of (4.6) into (4.5) and division by $Q_n(z)$ gives

$$(4.7) \quad \frac{P_n(z)}{Q_n(z)} = \frac{1}{(z - a_r)Q_n(z)^2} \int_{-\infty}^{\infty} \frac{(t - a_r)Q_n(t)^2}{t-z} d\phi(t) - \hat{\phi}(z).$$

Now assume that ϕ is a solution of the ESMP, so that we may write

$$(4.8) \quad -\hat{\phi}(z) = \frac{P_n(z)}{Q_n(z)} - \frac{1}{(z - a_r)Q_n(z)^2} \int_{\alpha}^{\beta} \frac{(t - a_r)Q_n(t)^2}{t-z} d\phi(t).$$

Let $z \in (a_\rho, a_\sigma)$ for some ρ, σ , where $(a_\rho, a_\sigma) \cap [\alpha, \beta] = \emptyset$. Then either $(t - a_\rho)$, $(t - a_\sigma)$, and $(t - z)$ are positive for all $t \in [\alpha, \beta]$, or $(t - a_\rho)$, $(t - a_\sigma)$, and $(t - z)$ are negative for all $t \in [\alpha, \beta]$. The numbers $(z - a_\rho)$ and $(z - a_\sigma)$ have opposite sign. Therefore $\hat{\phi}(z) + P_n(z)/Q_n(z)$ and $\hat{\phi}(z) + P_m(z)/Q_m(z)$ have opposite sign when n is of the form $n = pk + \rho$ and m is of the form $m = pk + \sigma$. It follows that if the sequences $\{P_{pk+\rho}(z)/Q_{pk+\rho}(z): k = 0, 1, 2, \dots\}$ and $\{P_{pk+\sigma}(z)/Q_{pk+\sigma}(z): k = 0, 1, 2, \dots\}$ are convergent and converge to the same value, then this value is $-\hat{\phi}(z)$. Thus if these two sequences converge to the same function on a subinterval of (a_ρ, a_σ) , then this function is $-\hat{\phi}(z)$. From the uniqueness of the Stieltjes transform of the solution follows uniqueness of the solution itself. (Recall that the transform is known on $\mathbf{C} - [\alpha, \beta]$ when it is known on an interval.)

We may sum up the results in this section as follows:

THEOREM 2. *The following statements are equivalent.*

- (A) *The ESMP has a unique solution.*
- (B) *The sequence $P_n(z)/Q_n(z)$ converges on $\mathbf{C} - [\alpha, \beta]$.*
- (C) *There exist an interval (a_ρ, a_σ) disjoint from $[\alpha, \beta]$ such that the sequences $\{P_{pk+\rho}(z)/Q_{pk+\rho}(z): k = 0, 1, 2, \dots\}$ and $\{P_{pk+\sigma}(z)/Q_{pk+\sigma}(z): k = 0, 1, 2, \dots\}$ converge to the same value (depending on z) for every z in some interval $(a, b) \subset (a_\rho, a_\sigma)$.*

REFERENCES

1. W. B. Jones and W. J. Thron, *Survey of continued fraction methods of solving moment problems and related topics*, Analytic Theory of Continued Fractions (Proceedings Loen, 1981), Lecture Notes in Math., vol. 932, Springer, Berlin, 1982, pp. 4–37.
2. W. B. Jones, W. J. Thron, and H. Waadeland, *A strong Stieltjes moment problem*, Trans. Amer. Math. Soc. **261** (1980), 503–528.
3. O. Njåstad, *An extended Hamburger moment problem*, Proc. Edinburgh Math. Soc. **28** (1985), 167–183.
4. —, *Unique solvability of an extended Hamburger moment problem*, J. Math. Anal. Appl. **124** (1987), 502–519.
5. —, *Orthogonal rational functions with poles in a finite subset of $\mathbf{R}$* . II. Symposium on Orthogonal Polynomials and Their Applications, Segovia, 1986.
6. O. Perron, *Die Lehre von den Kettenbrüchen*, 3. Auflage, Band 2, Teubner, Stuttgart, 1957.
7. T. J. Stieltjes, *Recherches sur les fractions continues*, Ann. Fac. Sci. Toulouse Math. **8** (1894), 1–122.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TRONDHEIM-NTH, N-7034, TRONDHEIM, NORWAY

Current address: Department of Mathematics, Campus Box 426, University of Colorado, Boulder, Colorado 80309