SMOOTH CONVEX t -NORMS DO NOT EXIST

C. ALSINA AND M. S. TOMAS

(Communicated by Richard R. Goldberg)

ABSTRACT. We show that smooth convex t -norms do not exist.

Let T be a t -norm, i.e., a binary operation on $[0, 1]$ which is associative, commutative, nondecreasing in each place, and has 1 as a unit element. A t -norm T is *convex* if, for all λ in $[0, 1]$ and for all a, b, c, d in $[0, 1]$, T satisfies:

$$T(\lambda a + (1 - \lambda)b, \lambda c + (1 - \lambda)d) \leq \lambda T(a, c) + (1 - \lambda)T(b, d).$$

Convex t -norms play a fundamental role in some studies of products of probabilistic metric spaces (see [2, 6]). An example of continuous convex t -norm is $W(x, y) = \text{Max}(x + y - 1, 0)$ and a discontinuous example is $Z(x, y) = 0$ whenever (x, y) is in $[0, 1) \times [0, 1)$ and $Z(x, y) = W(x, y)$ otherwise.

All other examples of continuous convex t -norms are closely related to W . Consequently, it is natural to ask whether smooth convex t -norms can exist. Our aim in this paper is to show that they cannot. To this end we begin with the following

LEMMA 1. *If T is a continuous convex t -norm, then: (i) $T \leq W$; (ii) T admits a representation of the form $T(x, y) = f^{(-1)}(f(x) + f(y))$, where f (the additive generator of T) is a continuous strictly decreasing function from $[0, 1]$ onto $[0, 1]$ such that $f(0) = 1$, $f(1) = 0$, and $f^{(-1)}$ is the pseudo-inverse of f , i.e., $f^{(-1)} = f^{-1}$ on $[0, 1]$ and $f^{(-1)}(x) = 0$, whenever $x \geq 1$; (iii) f is concave on $[f^{-1}(\frac{1}{2}), 1]$.*

PROOF. Since T is a convex t -norm, for any a, b and λ in $[0, 1]$ we have

$$(1) \quad T(\lambda a + (1 - \lambda)b, \lambda b + (1 - \lambda)a) \leq \lambda T(a, b) + (1 - \lambda)T(b, a) = T(a, b).$$

Suppose x, y in $[0, 1]$ are such that $x + y > 1$. Then substituting $a = x + y - 1$, $b = 1$, and $\lambda = (1 - x)/(2 - x - y)$ in (1) yields $T(x, y) \leq T(x + y - 1, 1) = x + y - 1$, from which (i) follows. The representation of T follows from the theorem of Ling [5]. Finally, the substitution $\lambda = \frac{1}{2}$ into (1) yields $T(\frac{1}{2}(a + b), \frac{1}{2}(a + b)) \leq T(a, b)$. Using the representation, for any a, b in $[f^{-1}(\frac{1}{2}), 1]$ we have $(a + b)/2 \geq f^{-1}(\frac{1}{2})$ and $2f((a + b)/2) < 1$. Therefore $f^{-1}(2f((a + b)/2)) > 0$ and we obtain

$$0 < f^{-1}\left(2f\left(\frac{a+b}{2}\right)\right) = T\left(\frac{a+b}{2}, \frac{a+b}{2}\right) \leq T(a, b) = f^{-1}(f(a) + f(b)),$$

whence $f((a + b)/2) \geq \frac{1}{2}(f(a) + f(b))$, i.e., f is concave on $[f^{-1}(\frac{1}{2}), 1]$.

DEFINITION 1. A t -norm T representable in the form

$$T(x, y) = f^{(-1)}(f(x) + f(y))$$

Received by the editors April 2, 1986 and, in revised form, September 15, 1986.
1980 *Mathematics Subject Classification* (1985 Revision). Primary 39C05, 26B25.

©1988 American Mathematical Society
0002-9939/88 \$1.00 + \$.25 per page

will be called *smooth* if f' and f'' exist on $(0, 1)$, $f'(x) \neq 0$ for all x in $(0, 1)$, $\lim_{x \rightarrow 0^+} x \cdot f'(x) = -\infty$, and $f''(x)$ is bounded on an arbitrary left-neighborhood of 1 of the form $(1 - \varepsilon, 1)$.

Thus for smooth t -norms T the first and second partial derivatives of T exist on $(0, 1) \times (0, 1)$.

LEMMA 2. If $T(x, y) = f^{(-1)}(f(x) + f(y))$ is a smooth convex t -norm, then

$$\lim_{k \rightarrow 1^-} \frac{f''(k)}{f'(k)^2} = 0.$$

PROOF. Since f'' is assumed to be bounded on a neighborhood of 1, we will prove that $\lim_{k \rightarrow 1^-} f'(k) = -\infty$. Choose a sequence $(y_n)_{n \in \mathbb{N}}$ converging to 0 so that $(1 - f(y_n))/y_n > n$. (This is possible because $\lim_{x \rightarrow 0^+} f'(x) = -\infty$.) Let $x_n = f^{-1}(1 - f(y_n))$. Since $f(x_n) + f(y_n) = 1$ and $T \leq W$ we have

$$0 = T(y_n, 1 - y_n) = f^{(-1)}(1 - f(x_n) + f(1 - y_n)),$$

i.e., $1 - f(x_n) + f(1 - y_n) \geq 1$. Hence $1 - x_n \leq y_n$ and we have

$$\frac{f(x_n)}{1 - x_n} \geq \frac{f(x_n)}{y_n} = \frac{1 - f(y_n)}{y_n} > n.$$

It follows that

$$\lim_{x \rightarrow 1^-} f'(x) = \lim_{n \rightarrow \infty} \frac{f(x_n)}{x_n - 1} = -\infty.$$

LEMMA 3. If g is a function from $(0, 1)$ into $\mathbb{R}$ satisfying

- (i) $g(u) \leq 0$ whenever u is in $(0, \frac{1}{2}]$;
- (ii) $g(u) \leq g(u + v)$ for all u, v in $(0, 1)$ with $u + v \leq 1$;
- (iii) $g(u) \cdot g(v) \geq (g(u) + g(v)) \cdot g(u + v)$ for all u, v in $(0, 1)$ with $u + v \leq 1$;
- (iv) $\lim_{h \rightarrow 0^+} g(h)$ exists

then $g(x) = 0$ for all x in $(0, 1)$.

PROOF. Given any x in $(0, \frac{1}{2}]$ the substitution $u = v = x$ into (iii) yields $g(x)^2 \geq 2g(x)g(2x)$. If $g(x) < 0$ then $g(x) \leq 2g(2x)$ and by induction

$$g(2x) \geq \frac{1}{2}g(x) \geq \frac{1}{2^{(n+1)}} \cdot g\left(\frac{x}{2^n}\right),$$

whence by (iv)

$$g(2x) \geq \frac{1}{2}g(x) \geq \lim_{n \rightarrow \infty} \frac{1}{2^{n+1}} \cdot g\left(\frac{x}{2^n}\right) = 0,$$

i.e., $g(x) \geq 0$ in contradiction with $g(x) < 0$. Therefore by (i) we have $g(x) = 0$ whenever x is in $(0, \frac{1}{2}]$. Now for any x in $(0, 1)$ we have by (ii), $g(x/2) = 0 \leq g(x)$. Consequently for any u, v in $(0, 1)$ with $u + v \leq 1$ we will have by (iii) and (ii):

$$g(u) \cdot g(v) \geq (g(u) + g(v)) \cdot g(u + v) \geq (g(u) + g(v))g(u).$$

Thus $0 \geq g(u)^2$ and we can conclude $g(u) = 0$ for all u in $(0, 1)$. We remark that condition (iv) is essential because there are nonzero functions such as $g(x) = -1/x$ that satisfy (i), (ii), and (iii).

THEOREM 1. *Smooth convex t -norms cannot exist.*

PROOF. Let T be a continuous convex t -norm. By Lemma 1, T can be represented in the form

$$(2) \quad T(x, y) = f^{(-1)}(f(x) + f(y)),$$

where f is concave on $[f^{-1}(\frac{1}{2}), 1]$. If T is smooth and convex then it is well known [4] that the following conditions hold:

$$(3) \quad \frac{\partial^2 T}{\partial x^2}(u, v) \geq 0$$

and

$$(4) \quad \frac{\partial^2 T}{\partial x^2}(u, v) \cdot \frac{\partial^2 T}{\partial y^2}(u, v) - \left[\frac{\partial^2 T}{\partial x \partial y}(u, v) \right]^2 \geq 0$$

for all u, v in $(0, 1)$. Using (2), after some computations we have that for any x, y in $(0, 1)$ with $f(x) + f(y) < 1$ the inequalities (3) and (4) become

$$(5) \quad \frac{f''(x)}{f'(x)^2} \leq \frac{f''(T(x, y))}{f'(T(x, y))^2},$$

and

$$(6) \quad \frac{f''(x)f''(y)}{f'(x)^2 f'(y)^2} \geq \frac{f''(x)f''(T(x, y))}{f'(x)^2 f'(T(x, y))^2} + \frac{f''(y)f''(T(x, y))}{f'(y)^2 f'(T(x, y))^2}.$$

Let $f(x) = u$ and $f(y) = v$. Then for u, v in $(0, 1)$ with $u + v < 1$, we have

$$(7) \quad \frac{f''(f^{-1}(u))}{f'(f^{-1}(u))^2} \leq \frac{f''(f^{-1}(u+v))}{f'(f^{-1}(u+v))^2}$$

and

$$(8) \quad \begin{aligned} & \frac{f''(f^{-1}(u))}{f'(f^{-1}(u))^2} \cdot \frac{f''(f^{-1}(v))}{f'(f^{-1}(v))^2} \\ & \geq \left[\frac{f''(f^{-1}(u))}{f'(f^{-1}(u))^2} + \frac{f''(f^{-1}(v))}{f'(f^{-1}(v))^2} \right] \cdot \frac{f''(f^{-1}(u+v))}{f'(f^{-1}(u+v))^2}. \end{aligned}$$

Define g from $(0, 1)$ into $\mathbf{R}$ by $g(z) = f''(f^{-1}(z))/f'(f^{-1}(z))^2$. Obviously g is well defined and by Lemma 2 we have

$$\lim_{k \rightarrow 1^-} \frac{f''(k)}{f'(k)^2} = \lim_{h \rightarrow 0^+} g(h) = 0.$$

Since f is concave on $[f^{-1}(\frac{1}{2}), 1]$ we have

$$(9) \quad g(u) \leq 0 \quad \text{whenever } u \text{ is in } (0, \tfrac{1}{2});$$

and by (7) and (8) for all u, v in $(0, 1)$ with $u + v < 1$ we obtain

$$(10) \quad g(u) \leq g(u + v)$$

and

$$(11) \quad g(u) \cdot g(v) \geq (g(u) + g(v)) \cdot g(u + v).$$

Now we can apply Lemma 3 and conclude that g must be the zero function, i.e., that $f''(f^{-1}(z)) = 0$ for all z in $(0, 1)$. This in turn implies that f must have the form $f(x) = ax + b$ for some constants a, b and contradicts the assumption that $\lim_{x \rightarrow 0^+} f'(x) = -\infty$.

ACKNOWLEDGMENTS. The authors thank Professor M. J. Frank (Illinois Institute of Technology, Chicago) for his interesting remarks concerning this paper and the referee for some helpful comments.

BIBLIOGRAPHY

1. J. Aczél, *Lectures on functional equations and their applications*, Academic Press, New York and London, 1966.
2. C. Alsina, *On countable products of probabilistic metric spaces and algebraic convexifications*, Pacific J. Math. **76** (1978), 291–300.
3. C. Alsina and B. Schweizer, *The countable product of probabilistic metric spaces*, Houston J. Math. **9** (1983), 303–310.
4. G. H. Hardy, J. E. Littlewood and G. Polya, *Inequalities*, Cambridge Univ. Press, Cambridge, 1934.
5. C. H. Ling, *Representation of associative functions*, Publ. Math. Debrecen **12** (1965), 189–212.
6. B. Schweizer and A. Sklar, *Probabilistic metric spaces*, Elsevier, New York, 1983.

DEPARTAMENT MATEMÀTIQUES I ESTADÍSTICA, ESCOLA TÉCNICA SUPERIOR D'ARQUITECTURA, UNIVERSITAT POLITÈCNICA DE CATALUNYA, AVDA DIAGONAL, 649, 08028 BARCELONA, SPAIN