ON A CONJECTURE OF GRAHAM

J. W. SANDER

(Communicated by Larry T. Goldstein)

ABSTRACT. Let $a_1 < a_2 < \cdots < a_n$ be a finite sequence of positive integers containing a prime power p^d with the property: $a_i \neq p^k a_j$ for all i, j and $k > 0$. Then $\max_{i,j} a_i/(a_i, a_j) \geq n$.

In [4] R. L. Graham asks if the following is true: Let $\mathcal{A}$ be a finite sequence of n positive integers $a_1 < a_2 < \cdots < a_n$. Then $\max_{i,j} a_i/(a_i, a_j) \geq n$.

Several partial results have been obtained (see the references), particularly for the case where one of the a_i 's is prime. The proofs for the latter follow similar lines, E. Z. Chein [2] and R. Klein [5] use G. Weinstein's improvement [9] upon R. Winterle's earlier result [10] while R. J. Simpson [7] can do without. In this paper we use refinements of the methods of Winterle [10] and Klein [5] to prove the following theorem. The sequence $\mathcal{A}$ is said to be p -simple for a prime p if $a_i \neq p^k a_j$ for $1 \leq i, j \leq n$, $k > 0$.

THEOREM. Let $\mathcal{A}$ be p -simple for a prime $p \neq 2$ and let $\mathcal{A}$ contain a prime power $a_k = p^d$ ($d \geq 0$). Then

$$\max_{1 \leq i, j \leq n} \frac{a_i}{(a_i, a_j)} \geq n.$$

In order to prove the theorem we first consider the case where a_1 is the required prime power. For a given sequence $\mathcal{A}$ let $A = \{a_1, \dots, a_n\}$.

LEMMA. Let $\mathcal{A}$ be p -simple with $a_1 = p^d$ ($d \geq 0$). Then

$$\max_{i,j} \frac{a_i}{(a_i, a_j)} \geq n.$$

PROOF. We may assume $d > 0$, since the lemma is trivial for $a_1 = 1$. We suppose

$$(1) \quad \max_{i,j} \frac{a_i}{(a_i, a_j)} \leq n-1.$$

This implies

$$(2) \quad a_n \leq (n-1)(a_n, a_1) \leq (n-1)p^d.$$

For $k = 0, 1, \dots, d-1$ define

$$B_k = \{a_i \in A : p^k \parallel a_i\}, \quad \text{and} \quad B = A \setminus \bigcup_{k=0}^{d-1} B_k.$$

Received by the editors July 18, 1986 and, in revised form, November 5, 1986.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 10A05; Secondary 02A05.

©1988 American Mathematical Society
0002-9939/88 \$1.00 + \$.25 per page

Furthermore, for $k = 0, 1, \dots, d-1$ and each $a_i \in B_k$ define

$$(3) \quad T_k(a_i) = p^{h_{i,k}}(a_i - p^{k+1})$$

such that $h_{i,k}$ is maximal with

$$(4) \quad T_k(a_i) \leq (n-1)p^d.$$

Obviously, $T_k(a_i) > 0$ and the maximality of $h_{i,k}$ gives

$$(5) \quad T_k(a_i) > (n-1)p^{d-1}.$$

We have for $a_i \in B_k$

$$(6) \quad a_i \leq (n-1)p^k;$$

otherwise $a_i = p^k a'_i$ with $a'_i \geq n$, $p \nmid a'_i$. This yields $a_i/(a_i, a_1) = p^k a'_i/p^k = a'_i \geq n$ which contradicts (1). By (3), (4) and (6) we have $h_{i,k} \geq d-k$ which implies

$$(7) \quad p^d | T_k(a_i).$$

Since $(a_i, a_i - p^{k+1}) = (a_i, p^{k+1}) = p^k$ for $a_i \in B_k$, we get

$$(8) \quad (a_i, T_k(a_i)) = p^k.$$

We claim for $a_i \in B_k$, $a_j \in B$,

$$(9) \quad T_k(a_i) \neq a_j;$$

otherwise by (5) and (8)

$$a_j/(a_j, a_i) = T_k(a_i)/(T_k(a_i), a_i) > n-1$$

which contradicts (1).

We claim for $a_i \in B_{k_1}$, $a_j \in B_{k_2}$, $a_i \neq a_j$,

$$(10) \quad T_{k_1}(a_i) \neq T_{k_2}(a_j);$$

otherwise

$$p^{h_{i,k_1}}(a_i - p^{k_1+1}) = p^{h_{j,k_2}}(a_j - p^{k_2+1}),$$

hence

$$a_i/p^{k_1} - p = a_j/p^{k_2} - p$$

which contradicts the p -simplicity of $\mathcal{A}$.

Now define the following function $F: A \rightarrow \mathbb{N}$ by

$$F(a_i) = \begin{cases} T_k(a_i) & \text{for } a_i \in B_k, \\ a_i & \text{for } a_i \in B. \end{cases}$$

By (9) and (10) F is 1-1. By (2), (4) and (7) we have for all $i = 1, \dots, n$

$$0 < F(a_i) \leq (n-1)p^d \quad \text{and} \quad p^d | F(a_i).$$

This contradiction proves the lemma.

PROOF OF THE THEOREM. We may assume that $\mathcal{A}$ is proper, i.e. the g.c.d. of the a_i 's is 1. We may also assume that p^d is the lowest p -power in A and $d > 0$. We suppose

$$(11) \quad \max_{i,j} \frac{a_i}{(a_i, a_j)} \leq n-1.$$

Define $B_0 = \{1, 2, \dots, n-1\} \setminus \{p, p^2, \dots, p^{d-1}\}$ and $B_d = \{lp^d: \lceil n/p^d \rceil \leq l \leq n-2\}$; for $h = 1, 2, \dots, d-1$ let

$$B_h = \{lp^h: \lceil n/p^h \rceil \leq l \leq n-1, (l, p) = 1\}.$$

Obviously, $B_0, \dots, B_d$ are pairwise disjoint.

We claim

$$(12) \quad A \subseteq \bigcup_{h=0}^d B_h;$$

therefore, let $a_i \in A$. If $a_i \leq n-1$ then $a_i \in B_0$ since p^d is the lowest p -power in A . If $a_i \geq n$ then $p|a_i$, for otherwise $a_i/(a_i, a_k) = a_i/(a_i, p^d) = a_i \geq n$, contradicting (11). Now we distinguish between two cases:

Case 1. $p^h|a_i$ for some h , $1 \leq h \leq d-1$. Then $a_i = p^h a'_i$, $(a'_i, p) = 1$, say. Hence $a'_i \geq \lceil n/p^h \rceil$ and by (11)

$$p^h a'_i = a_i \leq (n-1)(a_i, a_k) = (n-1)(a_i, p^d) = (n-1)p^h,$$

thus $a'_i \leq n-1$, so $a_i \in B_h$.

Case 2. $p^d|a_i$. Then $a_i = p^d a'_i$, say. Hence $a'_i \geq \lceil n/p^d \rceil$ and by the lemma $a_1 < a_k = p^d$, and by (11)

$$p^d a'_i = a_i \leq (n-1)(a_i, a_1) \leq (n-1)a_1 < (n-1)a_k = (n-1)p^d,$$

thus $a'_i < n-1$, so $a_i \in B_d$.

This proves the claim (12).

We define for $h = 1, \dots, d$

$$C_h = B_h \cap A, \quad C'_h = C_h \setminus \{(n-1)p^h\}, \quad C' = \bigcup_{h=1}^d C'_h.$$

Let $B'_0 = \{b \in B_0 : p \nmid b\}$. Define a function $f: C' \rightarrow B'_0$ in the following way:

$$(13) \quad x' \in C', \quad x = p^t b \quad (p \nmid b, \quad b \leq n-2) \text{ say; then } f(x) = b.$$

The function f is 1-1. Otherwise we have $b = b'$ for $x = p^t b$ and $x' = p^{t'} b'$ ($p \nmid bb'$). This implies $p^{t'} x = p^t x'$ which contradicts the p -simplicity of A .

For $b \in B'_0$ there is r with $1 \leq r \leq p-1$ and $b \equiv r \pmod{p}$. Define a function $g: f(C') \rightarrow B'_0$ by

$$g(b) = \begin{cases} b+1 & \text{for } 2 \nmid r, \\ b-1 & \text{for } 2 \mid r. \end{cases}$$

Since $2|(p-1)$ and $b \leq n-2$ in (13) we have indeed

$$(14) \quad g(f(C')) \subseteq B'_0.$$

It is easily seen that g is 1-1:

The only interesting case is $g(b) = b+1$, $g(b') = b'-1$, $g(b) = g(b')$. This implies $b = sp + r$, $b' = s'p + r'$ with $2 \nmid r$, $2|r'$, for some s, s' , and $b+2 = b'$. Hence $(s' - s)p = r - r' + 2$. Since $|r - r'| \leq p-2$ we can have $p|(r - r' + 2)$ only for $r - r' + 2 = 0$ or $r - r' + 2 = p$. The former gives $r \equiv r' \pmod{2}$, the latter $r = p-1 \equiv 0 \pmod{2}$, both contradicting the conditions on r and r' .

For $C = \bigcup_{h=1}^d C_h$ and $B = \{1, 2, \dots, n-1\}$ define the function $F : C \rightarrow B$ as follows:

$$F(x) = \begin{cases} g(f(x)) & \text{for } x \in C', \\ p^h & \text{for } x = (n-1)p^h \ (p \nmid (n-1), 1 \leq h \leq p-1). \end{cases}$$

Indeed, $p^h \in B$, because $p^d \in B$; otherwise, since the g.c.d. of the a_i 's is 1, there is a_m with $p \nmid a_m$; thus $a_k/(a_k, a_m) = p^d \geq n$ which contradicts (11). By (14) F is 1-1.

For $x \in C'$, $x = p^t b$ ($p \nmid b$), say, we have $p \nmid F(x)$, so $(x, F(x)) = (p^t b, b \pm 1) = 1$. Since $x \in A$, $F(x)$ cannot belong to A , because $x/(x, F(x)) \geq n$ again contradicts (11). Also, the numbers p^h ($1 \leq h \leq d-1$) do not belong to A , because p^d is the lowest p -power in A .

Thus we have found a 1-1 function $F : C \rightarrow B$ with $\text{card}(F(C) \cup (A \cap B)) = n$; but $F(C) \cup (A \cap B) \subseteq B$. This contradiction proves the theorem.

REFERENCES

1. R. D. Boyle, *On a problem of R. L. Graham*, Acta Arith. **34** (1978), 163-177.
2. E. Z. Chein, *On a conjecture of Graham concerning a sequence of integers*, Canad. Math. Bull. **21** (1978), 285-287.
3. P. Erdős, *Problems and results on combinatorial number theory*, A Survey on Combinatorial Theory, North-Holland, Amsterdam, 1973, Chapter 12.
4. R. L. Graham, *Unsolved problem 5749*, Amer. Math. Monthly **77** (1970), 775.
5. R. Klein, *The proof of a conjecture of Graham for sequences containing primes*, Proc. Amer. Math. Soc. **95** (1985), 189-190.
6. J. Marica and J. Schönheim, *Differences of sets and a problem of Graham*, Canad. Math. Bull. **12** (1969), 635-637.
7. R. J. Simpson, *On a conjecture of R. L. Graham*, Acta Arith. **40** (1982), 209-211.
8. W. Y. Vélez, *Some remarks on a number theoretic problem of Graham*, Acta Arith. **32** (1977), 233-238.
9. G. Weinstein, *On a conjecture of Graham concerning greatest common divisors*, Proc. Amer. Math. Soc. **63** (1977), 33-38.
10. R. Winterle, *A problem of R. L. Graham in combinatorial number theory*, Proc. Louisiana Conf. on Combinatorics, Graph Theory and Computing, Louisiana State Univ., Baton Rouge, La., 1970, pp. 357-361.

INSTITUT FÜR MATHEMATIK, UNIVERSITÄT HANNOVER, WELFENGARTEN 1, 3000 HANNOVER 1, FEDERAL REPUBLIC OF GERMANY