ON THE PARTIAL SUMS OF CONVEX FUNCTIONS OF ORDER $1/2$

RAM SINGH

(Communicated by Paul S. Muhly)

ABSTRACT. Let $f(z) = z + a_2z^2 + \cdots$ be regular and univalently convex of order $1/2$ in the unit disc U and let $s_n(z, f)$ denote its n th partial sum. In the present note we determine the radius of convexity of $s_n(z, f)$, depending on n , and generalize and sharpen a result of Ruscheweyh concerning the partial sums of convex functions. We also prove that for every $n \geq 1$, $\operatorname{Re}(s_n(z, f)/z) > 1/2$ in U .

1. Let S denote the class of functions

$$f(z) = z + \sum_{k=2}^{\infty} a_k z^k$$

which are regular and univalent in the unit disc $U = \{z \mid |z| < 1\}$. Denote by S_t and K the usual subclasses of S consisting of functions which map U onto starlike (with respect to the origin) and convex domains, respectively. Let $S_t(1/2) \subset S_t$ be the class of functions which are starlike of order $1/2$. Similarly, let $K(1/2) \subset K$ be the family of functions which are convex of order $1/2$. It is well known that $K \subset S_t(1/2)$. Let

$$s_n(z, f) = z + \sum_{k=2}^n a_k z^k$$

denote the n th partial sum of $f(z)$. Kobori [2] proved that if $f \in K$, then for each $n \geq 1$, $s_n(z, f)$ is starlike and univalent in $|z| < 1/2$. Using the theory of convolution, Ruscheweyh [5] in 1972 obtained the following result.

THEOREM A. *Let r_n denote the positive root of*

$$\varphi_n(r) = 1 - (n+1)r^n - nr^{n+1} \quad (n \in \mathbb{N}),$$

and let $f \in K$. Then $s_n(z, f)$ is univalent for $|z| < r_n$ and maps this disc onto a close-to-convex domain. For n even, r_n cannot be replaced by any larger number.

Robertson [3] in 1981 proved that if $f \in K(1/2)$, then each $s_n(z, f)$ is close-to-convex with respect to f and hence univalent in U .

Using the fact that the class K is closed with respect to Hadamard convolution, we can readily compute the radius of convexity of $s_n(z, f)$, $f \in K$, depending on n . In the present paper we determine the radius of convexity of $s_n(x, f)$, $f \in K(1/2)$, in terms of n . As a corollary to our result, we shall show that one can weaken the hypothesis of Ruscheweyh's theorem and still make a stronger assertion. We shall also prove that if $f \in K(1/2)$, then $\operatorname{Re}(s_n(z, f)/z) > 1/2$ ($z \in U$) for each $n \in \mathbb{N}$.

Received by the editors November 29, 1986 and, in revised form, January 26, 1987.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 30A26, 30A34.

Key words and phrases. Univalent, starlike and convex functions, convex null sequences.

2. We shall need the following definitions and results.

Let f and g be analytic in $|z| < R$ and $f(0) = g(0)$. In addition, suppose that g is univalent in $|z| < R$. Then we say that f is subordinate to g in $|z| < R$, in symbols, $f \prec g(|z| < R)$, if $f(|z| < R) \subset g(|z| < R)$.

LEMMA 1. *If f is analytic in U and $f(0) = f'(0) - 1 = 0$, then $f \in S_t(1/2)$ if and only if*

$$\operatorname{Re} \left(\frac{f(z_1) - f(z_2)}{z_1 - z_2} \frac{z_2}{f(z_2)} \right) > \frac{1}{2}$$

for $|z_1| < 1$, $|z_2| < 1$.

LEMMA 2. *For $0 \leq \theta \leq \pi$,*

$$\frac{1}{2} + \sum_{k=1}^n \frac{\cos k\theta}{k+1} \geq 0.$$

Lemma 1 is due to Ruscheweyh and Sheil-Small [6] and Lemma 2 is due to Rogosinski and Szegő [4].

3. We now prove the following:

THEOREM 1. *Let $f \in K(1/2)$ and let r_n be defined as in Theorem A. Then $s_n(z, f)$ maps the disc $|z| < r_n$ onto a convex domain. For even n , the number r_n cannot be replaced by any larger one.*

PROOF. Since $f \in K(1/2)$, $g(z) = zf'(z) \in S_t(1/2)$ and, therefore, in view of Lemma 1, it follows that for all z and ζ in U ,

$$\operatorname{Re} \left(\frac{g(z) - g(\zeta)}{z - \zeta} \frac{\zeta}{g(\zeta)} \right) > \frac{1}{2}.$$

Treating ζ as a constant, this leads to

$$\begin{aligned} \operatorname{Re} \left[1 + \frac{1}{\zeta} \left(1 - \frac{s_1(\zeta, g)}{g(\zeta)} \right) z + \frac{1}{\zeta^2} \left(1 - \frac{s_2(\zeta, g)}{g(\zeta)} \right) z^2 \right. \\ \left. + \cdots + \frac{1}{\zeta^n} \left(1 - \frac{s_n(\zeta, g)}{g(\zeta)} \right) z^n + \cdots \right] > \frac{1}{2} \quad (z \in U), \end{aligned}$$

and, consequently, for all $n \geq 1$, we get

$$\left| \frac{1}{\zeta^n} \left(1 - \frac{s_n(\zeta, g)}{g(\zeta)} \right) \right| \leq 1 \quad [1, \text{p. 41}].$$

Replacing ζ by z and writing g in terms of f , the above inequality provides

$$(1) \quad 1 - \frac{s'_n(z, f)}{f'(z)} = z^{n-1} \varphi(z),$$

where φ is analytic, $\varphi(0) = 0$ and $|\varphi(z)| \leq |z|$ in U . From (1), we get

$$\begin{aligned} \operatorname{Re} \left(1 + \frac{zs''(z, f)}{s'(z, f)} \right) &= \operatorname{Re} \left[\left(1 + \frac{zf''(z)}{f'(z)} \right) - z^{n-1} \frac{((n-1)\varphi(z) + z\varphi'(z))}{1 - z^{n-1}\varphi(z)} \right] \\ &\geq \frac{1}{1+r} - \frac{r^{n-1}}{(1-r^2)} \left[\frac{(n-1)(1-r^2)t + r(1-t^2)}{(1-r^{n-1}t)} \right], \end{aligned}$$

where $r = |z|$, $t = |\varphi(z)|$ and use has been made of the well-known inequalities:

$$|\varphi'(z)| \leq \frac{1 - |\varphi(z)|^2}{1 - |z|^2},$$

$$\operatorname{Re} \left(1 + \frac{zf''(z)}{f'(z)} \right) \geq \frac{1}{1 + |z|} \quad (f \in K(1/2)).$$

Letting

$$\psi(t) = \frac{(n-1)(1-r^2)t + r(1-t^2)}{1 - r^{n-1}t},$$

we readily see that

$$\begin{aligned} N(\psi'(t)) &= \text{the numerator of } \psi'(t) \\ &= (n-1)(1-r^2) + r^n - 2rt + r^n t^2, \end{aligned}$$

and that $(\partial/\partial t)N(\psi'(t)) \leq 0$.

$$\begin{aligned} \therefore \min N(\psi'(t)) &= N(\psi'(r)) = (n-1) - (n+1)r^2 + r^n + r^n + 2 \\ &= (r-1)^2[r^n + 2r^{n-1} + 4r^{n-2} + \cdots + 2(n-1)r + (n-1)] > 0. \end{aligned}$$

Thus we conclude that $\psi'(t) > 0$ for all admissible values of t and as such

$$\begin{aligned} \operatorname{Re} \left(1 + \frac{zs_n''(z, f)}{s_n'(z, f)} \right) &\geq \frac{1}{1+r} - \frac{nr^n}{1-r^n} \\ &= \frac{1 - (n+1)r^n - nr^{n+1}}{(1+r)(1-r^n)}. \end{aligned}$$

$s_n(z, f)$ will, therefore, map the disc $|z| < r_n$ onto a convex domain, where $0 < r_n < 1$ is the unique positive root of the polynomial $\varphi_n(r)$, defined in Theorem A. This proves the first assertion of our theorem. To prove the second assertion, let $n = 2m$ be any even positive integer, and consider the function $f_0(z) = -\log(1-z)$, which belongs to $K(1/2)$. Then

$$\begin{aligned} 1 + \frac{s_{2m}''(z, f_0)}{s_{2m}'(z, f_0)} &= \frac{1 + 2z + 3z^2 + \cdots + 2mz^{2m-1}}{1 + z + z^2 + \cdots + z^{2m-1}} \\ &= \frac{1 - (2m+1)z^{2m} + 2mz^{2m+1}}{(1-z)(1-z^{2m})}, \end{aligned}$$

and, therefore,

$$\begin{aligned} \left[1 + \frac{zs_n''(z, f_0)}{s_n'(z, f_0)} \right]_{z=-r} &= \frac{1 - (n+1)r^n - nr^{n+1}}{(1+r)(1-r^n)} \\ &= 0, \quad \text{when } r = r_n. \end{aligned}$$

The proof of Theorem 1, is, therefore, complete.

COROLLARY 1. *If $f \in S_t(1/2)$, then $s_n(z, f)$ maps the disc $|z| < r_n$ onto a domain which is starlike with respect to the origin. For even n , the number r_n cannot be replaced by any larger one.*

Since $K \subset S_t(1/2)$, Corollary 1 shows that one can weaken the hypothesis of Ruschewyh's theorem (K may be replaced by $S_t(1/2)$) and still make a stronger assertion ($|z| < r_n$ is mapped onto a starlike domain).

As Ruscheweyh [5] has shown, the number r_n satisfies

$$(2n)^{-1/n} \leq r_n \leq n^{-1/(n-1)}$$

and

$$r_n = 1 - \frac{\log(2n)}{n} + \frac{\log(n)\log(4en)}{2n^2} + o\left(\frac{\log(n)}{n^2}\right).$$

THEOREM 2. *Let $f \in K(1/2)$. Then for each $n \geq 1$,*

$$\operatorname{Re} \frac{s_n(z, f)}{z} > \frac{1}{2} \quad (z \in U).$$

The constant $1/2$ cannot be replaced by any larger one.

PROOF. Since $f \in K(1/2)$, the function $zf'(z) \in S_t(1/2)$ and, hence,

$$(2) \quad \operatorname{Re} f'(z) > \frac{1}{2} \quad (z \in U).$$

Now we can write

$$(3) \quad \begin{aligned} \frac{s_n(z, f)}{z} &= f'(z) * \left[1 + \sum_{k=2}^n \frac{z^{k-1}}{k} \right] \\ &= f'(z) * \left[1 + \sum_{k=1}^{n-1} \frac{z^k}{k+1} \right], \end{aligned}$$

where $*$ denotes Hadamard convolution.

Putting $z = re^{i\theta}$, $0 \leq r < 1$, $0 \leq \theta < 2\pi$, and making use of the minimum principle for harmonic functions along with Lemma 2, we obtain

$$(4) \quad \begin{aligned} \operatorname{Re} \left[1 + \sum_{k=1}^{n-1} \frac{z^k}{k+1} \right] &= \left[1 + \sum_{k=1}^{n-1} \frac{r^k \cos k\theta}{k+1} \right] \quad (0 \leq \theta \leq \pi) \\ &> 1 + \sum_{k=1}^{n-1} \frac{\cos k\theta}{k+1} \geq \frac{1}{2}. \end{aligned}$$

From (2), (4) and (3) we deduce that $\operatorname{Re}(s_n(z, f)/z) > 1/2$ in U . Since for the function $f_0(z) = -\log(1-z) \in K(1/2)$, we have $(s_2(z, f)/z) = 1 + (z/2)$, the sharpness of the number $1/2$ is obvious.

REMARK. It is clear from the proof of Theorem 2 that its assertion holds for the wider class of functions f which are regular in and satisfy the conditions $\operatorname{Re} f'(z) > 1/2$, $z \in U$.

It is well known that if $f \in K$, then $z/2 = s_1(z, f)/2 \prec f(z)$ in U . Since each $s_n(z, f)$, $f \in K(1/2)$, is univalent in U , it is natural to ask for the largest number λ_n , $0 < \lambda_n < 1$, such that $\lambda_n s_n(z, f) \prec s_{n+1}(z, f)$ in U . The following theorem, which we state without proof, provides a lower bound for λ_n .

THEOREM 3. *If $f \in K(1/2)$, then*

$$(i) \quad z/2 = \frac{1}{2}s_1(z, f) \prec s_2(z, f), \quad (z \in U),$$

and for $n \geq 2$,

$$(ii) \quad ((n-1)/(n+1))s_n(z, f) \prec s_{n+1}(z, f), \quad (z \in U).$$

The number $1/2$ in (i) cannot be replaced by any larger one.

REFERENCES

1. P. L. Duren, *Univalent functions*, Springer-Verlag, New York, 1983.
2. A. Kobori, *Zwei Satze über die abschnitte schlichter Potenzreihen*, Mem. Coll. Kyoto **17** (1934), 172–186.
3. M. S. Robertson, *Univalent functions starlike with respect to a boundary point*, J. Math. Anal. Appl. **81** (1981), 327–346.
4. W. Rogosinski and G. Szegő, *Über die Abschnitte von Potenzreihen die in einem Kreise beschränkt bleiben*, Math. Z. **28** (1928), 73–94.
5. St. Ruscheweyh, *On the radius of univalence of the partial sums of convex functions*, Bull. London Math. Soc. **4** (1972), 367–369.
6. St. Ruscheweyh and T. Sheil-Small, *Hadamard products of schlicht functions and the Pólya-Schoenberg conjecture*, Comment. Math. Helv. **48** (1973), 119–135.

DEPARTMENT OF MATHEMATICS, PUNJABI UNIVERSITY, PATIALA-147002, INDIA