ON THE CARDINALITY OF A TOPOLOGY

RUAN YONGBIN

(Communicated by Dennis Burke)

ABSTRACT. Let $o(X)$ denote the cardinality of topology of a space X . I. Juhasz proves that $o(X)^\omega = o(X)$ for regular hereditarily paracompact spaces. We prove it for more general classes of spaces.

If X is a topological space, let $o(X)$ denote the cardinality of its topology. A number of papers [1, 2, 3] have investigated classes for which $o(X) = o(X)^\omega$. We generalize several of these results by proving

THEOREM 1. *For every infinite hereditarily weakly θ -refinable T_3 space X , $o(X) = o(X)^\omega$.*

A space X is *weakly θ -refinable* [4] if and only if each open cover $\mathcal{U}$ of X has a refinement $\mathcal{V} = \bigcup_{n \in \omega} \mathcal{V}_n$ where each $\mathcal{V}_n$ is a discrete family of subsets of the space $\bigcup \mathcal{V}_n$.

If η and μ are infinite cardinals and $\mu' < \mu$ implies $\eta^{\mu'} < \eta^\mu$ and $\eta' < \eta$ implies $\eta'^\mu < \eta^\mu$, then η^μ is called a *jump*. By Theorem 0.2 of [2], if η^μ is a jump, μ is regular.

Since X is T_3 , by Theorem 4.3 of [2], if μ is an uncountable cardinal of cofinality ω and there is a discrete subset S' of X with $|S'| = \mu'$ for all $\mu' < \mu$, then there is a discrete subset S of X with $|S| = \mu$.

Before proving the theorem, we also need

LEMMA. *If $\langle \kappa_\alpha \mid \alpha \in \gamma \rangle$ is a sequence of infinite cardinals, then there is an $S \subset \gamma$ with $\prod_{\alpha \in \gamma} \kappa_\alpha = (\sup_{\alpha \in S} \kappa_\alpha)^{|S|}$.*

PROOF OF LEMMA. Enumerate the κ_α so that they form a nondecreasing sequence. Let $S \subset \gamma$ be of minimal order type so that $\prod_{\alpha \in \gamma} \kappa_\alpha = \prod_{\alpha \in S} \kappa_\alpha$. Without loss of generality, suppose S is infinite. Note that if $\beta < \sup S$, then $S - (\beta + 1)$ has the order type of S , in particular $|S - (\beta + 1)| = |S|$. So if $|S| = \lambda$ we have that $S = \lambda \circ \alpha$ for $\alpha = 1$ or for some limit ordinal $\alpha < \lambda^+$. Split S into λ many cofinal disjoint sets $\{A_\beta \mid \beta \in \lambda\}$. Note that every A_β is cofinal in S and so

$$\prod_{\delta \in A_\beta} \kappa_\delta \geq \sup_{\delta \in A_\beta} \kappa_\delta = \sup_{\alpha \in S} \kappa_\alpha = \kappa.$$

Received by the editors January 6, 1986 and, in revised form, April 18, 1986 and July 28, 1986.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 54A25, 04A30; Secondary 03E05.

Key words and phrases. Hereditary paracompactness, weak θ -refinability, weakly collectionwise normality.

Hence

$$\kappa^\lambda \leq \prod_{\beta \in \lambda} \prod_{\delta \in \mathcal{A}_\beta} \kappa_\delta = \prod_{\alpha \in S} \kappa_\alpha; \quad \text{so } \kappa^\lambda = \prod_{\alpha \in \gamma} \kappa_\alpha. \quad \square$$

PROOF OF THEOREM. Assume not and let X be a counterexample; so $o(X) = \kappa < \kappa^\omega$. Let $\lambda = \min\{\mu \mid \mu^\omega > \kappa\}$. Then $\text{cf}(\lambda) = \omega$ and $\lambda \leq \kappa$.

For $x \in X$, let $\sigma(x, X) = \min\{o(U) \mid x \in U \text{ and } U \text{ is open}\}$ and $\sigma = \sigma(X) = \sup\{\sigma(x, X) \mid x \in X\}$.

Note that there exists a finite subset $F \subset X$ such that $\sigma(X - F) < \lambda$. To see this assume the contrary and let $\langle \lambda_n \mid n \in \omega \rangle$ be a cofinal sequence in λ . At stage n pick $x_n \in X - \{x_k \mid k \in n\}$ with $\sigma(x_n, X) \geq \lambda_n$. Since X is regular there is a disjoint open family $\{U_n \mid n \in \omega\}$ with $x_n \in U_n$, hence $o(U_n) \geq \lambda_n$. So $o(X) \geq \prod_{n \in \omega} o(U_n) = \lambda^\omega > \kappa$, a contradiction.

Thus we can assume that $\sigma(X) < \lambda$ and that $o(X) = o(X - F)$ for all finite $F \subset X$ (recall that our original X was *hereditarily* weakly θ -refinable).

Let $\mathcal{U}$ be an open cover of X such that each U in $\mathcal{U}$ has $o(U) \leq \sigma$. Since X is weakly θ -refinable there is a refinement $\mathcal{V} = \bigcup_{n \in \omega} \mathcal{V}_n$ such that each $\mathcal{V}_n$ is a closed discrete family and $\bigcup \mathcal{V}_n = V_n$. Assume $\mathcal{V}_n = \{V_n^\alpha \mid \alpha \in \beta_n\}$ and β_n is infinite for each $n \in \omega$. Then $\kappa = o(X) \leq \prod_{n \in \omega} o(V_n)$. Now we derive a contradiction by showing that $\prod_{n \in \omega} o(V_n) < \kappa$.

Fixing $n \in \omega$, we have $o(V_n) = \prod_{\alpha \in \beta_n} o(V_n^\alpha) = \sigma_n^{\mu_n}$ for $\sigma_n = \sup\{o(V_n^\alpha) \mid \alpha \in \beta_n\}$ and $\mu_n = |\beta_n|$ by the lemma. Since $o(V_n^\alpha) \leq \sigma$ for every α , we have that $\sigma_n \leq \sigma$. Also $\sigma_n^{\mu_n} < \lambda$ since $o(V_n) \leq o(X) = \kappa$ and $\sigma_n^{\mu_n \cdot \omega} = \sigma_n^{\mu_n} \leq \kappa$ if $\mu_n \geq \omega$; otherwise $\sigma_n^{\mu_n} \leq \sigma < \lambda$. Note that $\prod\{\sigma_n^{\mu_n} \mid n \in \omega \text{ and } \sigma_n^{\mu_n} < \sigma\} \leq \sigma^\omega < \lambda$. So in order to show that $\prod_{n \in \omega} \sigma_n^{\mu_n} < \kappa$ we may assume that $\sigma \leq \sigma_n^{\mu_n}$ for all n . Then $\sigma_n^{\mu_n} = \sigma^{\mu_n} < \lambda$ (since $\sigma_n \leq \sigma \leq \sigma_n^{\mu_n} < \lambda$). Hence if $\mu = \sup_{n \in \omega} \mu_n$, $\prod_{n \in \omega} \sigma_n^{\mu_n} = \sigma^\mu$. If $\sigma^\mu < \lambda$, then $\prod_{n \in \omega} o(V_n) < \kappa$ as desired.

So assume that $\sigma^\mu \geq \lambda$. Since $\sigma^\omega < \lambda$, μ is uncountable and, since $\sigma^{\mu_n} < \lambda$ for all n , $\text{cf}(\mu) = \omega$. If $\eta = \min\{\alpha \mid \alpha^\mu \geq \lambda\}$, then $\eta' < \eta$ implies $\eta'^\mu < \eta^\mu$ and $\mu' < \mu$ implies $\eta^{\mu'} < \eta^\mu$. So if η is infinite, η^μ is a jump and μ is regular. This contradicts μ being uncountable and of cofinality ω .

So it remains to show that η is infinite if μ is. To show this, for each $n \in \omega$ and $\alpha \in \beta_n$ choose $x_\alpha \in V_n^\alpha$ and let $S_n = \{x_\alpha \mid \alpha \in \beta_n\}$. Since S_n is discrete and $|S_n| = \mu_n$, there is a discrete $S \subset X$ of cardinality μ . Thus $2^\mu \leq o(S) \leq o(X) = \kappa$. Since μ is infinite, $2^{\mu \cdot \omega} = 2^\mu \leq \kappa$ while $\lambda^\omega > \kappa$, so $2^\mu < \lambda$. Since $2^\mu = \eta^\mu$ for finite η and $\eta^\mu \geq \lambda$ by definition, η is infinite. $\square$

Our proof gives the stronger

THEOREM 2. *Let X be regular space. If for any finite subset F , $X - F$ is weakly θ -refinable, then $o(X) = o(X)^\omega$.*

Since weak θ -refinability is hereditary to F_σ -sets, $o(X) = o(X)^\omega$ also holds for 1st countable, T_3 , weakly θ -refinable spaces or 1st countable, T_3 paracompact spaces.

A similar proof also yields a theorem, independently proved by I. Juhasz, that $o(X) = o(X)^\omega$ for all hereditarily weakly collectionwise Hausdorff spaces.

In the end, I acknowledge my indebtedness to Zhou Hao-Xuan and Amer Beslagic for their help.

REFERENCES

1. I. Juhasz, *Two set-theoretic problems in topology*, Proc. 4th Prague Sympos. on General Topology (1976), Part A, Springer-Verlag, pp. 115–123.
2. ———, *Cardinal functions in topology—Ten years later*, MCT 123, Mathematisch Centrum, Amsterdam, 1980.
3. E. K. Van Douwen and Zhou Hao-Xuan, *The number of cozero-sets is an ω -power*, 1980.
4. H. R. Bennett and D. J. Lutzer, *A note on weak θ -refinability*, General Topology Appl. 2 (1972), 49–54.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MADISON, WISCONSIN 53706

Current address: Department of Mathematics, University of California-Berkeley, Berkeley, California 94720