SHORTER NOTES

The purpose of this department is to publish very short papers of unusually polished character, for which there is no other outlet.

ON AN INEQUALITY OF OSGOOD, PHILLIPS AND SARNAK

HAROLD WIDOM

(Communicated by Paul S. Muhly)

ABSTRACT. In their work [4, 5] on isospectral families for the Laplacian Osgood, Phillips and Sarnak needed and proved the following inequality: For any real-valued function ϕ belonging to the Sobolev space $H_{1/2}$ ($= W^{1/2,2}$) of the unit circle and satisfying the side condition $\int e^{\phi} e^{i\theta} d\theta = 0$ one has

$$\log \frac{1}{2\pi} \int e^{\phi} d\theta - \frac{1}{2\pi} \int \phi d\theta \leq \frac{1}{2} \sum_{k=1}^{\infty} k |\hat{\phi}(k)|^2$$

where $\hat{\phi}(k)$ is the k th Fourier coefficient of ϕ . Without the side condition the factor $\frac{1}{2}$ does not appear on the right side and the inequality is the first Lebedev-Milin inequality [1, §5.1]. The purpose of this note is to show that these are the first two of a series of inequalities which follow quickly from some theorems of G. Szegő on Toeplitz determinants.

Given a nonnegative integrable function f on the unit circle one defines the associated Toeplitz determinants by

$$D_n(f) = \det(\hat{f}(i-j))_{i,j=0,\dots,n}.$$

We have, first, the following family of inequalities:

For any $n = 0, 1, \dots$

$$(1) \quad \log D_n(e^{\phi}) - \frac{n+1}{2\pi} \int \phi d\theta \leq \sum_{k=1}^{\infty} k |\hat{\phi}(k)|^2.$$

Equality holds if $e^{-\phi}$ is a trigonometric polynomial of degree at most $n+1$.

To prove this we quote first the Szegő limit theorem [3, 5.5(a)] which states (after taking logarithms) that

$$(2) \quad \lim_{n \rightarrow \infty} \left\{ \log D_n(e^{\phi}) - \frac{n+1}{2\pi} \int \phi d\theta \right\} = \sum_{k=1}^{\infty} k |\hat{\phi}(k)|^2$$

Received by the editors September 14, 1987.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 42A05; Secondary 47B35.

Supported by a grant from the National Science Foundation.

©1988 American Mathematical Society
0002-9939/88 \$1.00 + \$.25 per page

and then the simpler fact [3, 5.5(5)] that the sequence

$$(3) \quad \left\{ \log D_n(e^\phi) - \frac{n+1}{2\pi} \int \phi d\theta \right\}$$

is nondecreasing. Thus each term of the sequence is less than or equal to its limit and (1) follows from (2). The assertion concerning equality in (1) follows from [3, 5.3(23)]: If $e^{-\phi}$ is a trigonometric polynomial of degree $n+1$ then the terms of the sequence (3) with n replaced by m are constant for $m \geq n$. Taking the limit as $m \rightarrow \infty$ and using (2) shows that we have equality in (1) in this case.

Suppose now that ϕ satisfies the side conditions

$$\int e^\phi e^{im\theta} d\theta = 0 \quad (m = 1, \dots, n).$$

Then $D_n(e^\phi)$ is the determinant of an $(n+1) \times (n+1)$ diagonal matrix each of whose diagonal entries equals $(2\pi)^{-1} \int e^\phi d\theta$. So

$$D_n(e^\phi) = \left\{ \frac{1}{2\pi} \int e^\phi d\theta \right\}^{n+1}$$

in this case and (1) becomes

$$\log \frac{1}{2\pi} \int e^\phi d\theta - \frac{1}{2\pi} \int \phi d\theta \leq \frac{1}{n+1} \sum_{k=1}^{\infty} k |\hat{\phi}(k)|^2$$

with equality if, for example, $e^{-\phi} = a + b \sin(n+1)\theta$ ($|b| < a$).

The knowledgeable reader might object that the proof of Szegő's limit theorem to which we referred requires a smoothness condition on ϕ . There are two reasons why this causes no problem. The first is that (2) holds without this extra condition [2]. The other is that, even if it did not, it is a simple matter to show that inequality (1) for smooth ϕ implies the inequality for general ϕ .

REFERENCES

1. P. L. Duren, *Univalent functions*, Springer-Verlag, 1983.
2. B. L. Golinskii and I. A. Ibragimov, *On Szegő's limit theorem*, Math. USSR Izv. **5** (1971), 421-446.
3. U. Grenander and G. Szegő, *Toeplitz forms and their application*, Univ. of California Press, 1958.
4. B. Osgood, R. Phillips and P. Sarnak, *Extremal determinants of Laplacians*, J. Funct. Anal. (to appear).
5. ———, *Compact isospectral sets of surfaces*, J. Funct. Anal. (to appear).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, SANTA CRUZ, CALIFORNIA 95064