PLURISUBHARMONIC FUNCTIONS OUTSIDE COMPACT SETS

URBAN CEGRELL

(Communicated by Irwin Kra)

ABSTRACT. We study the Monge-Ampère mass of plurisubharmonic functions defined outside a compact set.

1. Introduction. Let Ω be an open and bounded pseudoconvex set in $\mathbb{C}^n$, $n \geq 2$, and K a compact subset of Ω so that $\Omega \setminus K$ is connected. By the Hartogs extension theorem, every analytic function on $\Omega \setminus K$ extends to Ω . When it comes to plurisubharmonic functions the situation is different.

The purpose of this note is to study the Monge-Ampère mass $\int_{\Omega' \setminus K} (dd^c u)^n$, where $u \in \text{PSH} \cap L_{\text{loc}}^\infty(\Omega \setminus K)$ and $K \Subset \Omega' \Subset \Omega$.

The case when $u \in \text{PSH} \cap C^\infty(\Omega \setminus K)$ has already been studied by Fornaess and Sibony [3] and Griffith [4].

2. Estimates.

PROPOSITION. Assume that $\varphi \in \text{PSH} \cap L_{\text{loc}}^\infty(\Omega \setminus K)$, where K is a removable singularity set for the plurisubharmonic functions (cf. [2]). Then $\int_{\Omega' \setminus K} (dd^c \varphi)^n < +\infty$.

PROOF. By Bedford and Taylor [1], $(dd^c \varphi)^n$ is a well-defined positive measure on $\Omega \setminus K$. Since φ extends to a plurisubharmonic function on Ω , we can choose a sequence $\varphi_j \in \text{PSH} \cap C^\infty$ near $\overline{\Omega'}$ such that $\varphi_j \searrow \varphi$, $j \rightarrow +\infty$.

Let $W = \{z \in \Omega', d(z, K) > \frac{1}{2} d(\partial \Omega', K)\}$. Then $\delta = \inf_{z \in W} \varphi(z) > -\infty$, and if we put $\psi_j = \sup(\varphi_j, \delta)$, then $\psi_j = \varphi_j$ on W . So $\int_{\Omega'} (dd^c \psi_j)^n = \int_{\Omega'} (dd^c \varphi_j)^n$ by Stokes' theorem. Therefore,

$$\begin{aligned} \int_{\Omega' \setminus K} (dd^c \varphi)^n &\leq \lim_{j \rightarrow +\infty} \int_{\Omega'} (dd^c \varphi_j) \\ &= \lim_{j \rightarrow +\infty} \int_{\Omega'} (dd^c \psi_j)^n \\ &= \int_{\Omega'} (dd^c \sup(\varphi, \delta))^n < +\infty \end{aligned}$$

since $\sup(\varphi, \delta) \in \text{PSH} \cap L_{\text{loc}}^\infty(\Omega)$.

The next theorem is a generalization of Theorem 2.2 in Fornaess and Sibony [3].

THEOREM. Let $\psi \in \text{PSH} \cap C^2(\Omega)$ and assume that $\Omega' = \{\psi < 1\}$ and that $K = \{\psi \leq s\}$ for an s , $0 < s < 1$. If $\varphi \in \text{PSH} \cap L_{\text{loc}}^\infty(\Omega \setminus K)$, then

$$\int_{\{s < \psi < 1\}} |\varphi| (dd^c \varphi)^{n-1} \wedge dd^c \psi < +\infty$$

Received by the editors January 22, 1987.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 32F05; Secondary 31C10.

©1988 American Mathematical Society
0002-9939/88 \$1.00 + \$.25 per page

and

$$\int_{\{s < \psi < 1\}} (\psi - s)(dd^c \varphi)^n < +\infty.$$

PROOF. Since Ω is pseudoconvex and ψ plurisubharmonic, there is, to every $z_0 \in \{s < \psi < 1\}$, an analytic function such that $f(z) \neq f(z_0) \forall z \in K$. If we restrict φ to $\{f(z) = f(z_0)\}$ and apply the maximum principle, we conclude that φ is uniformly bounded above on $\overline{\Omega'} \setminus K$. It is therefore no restriction to assume that $\varphi \leq 0$.

Let $s < r < 1$. Then, by Stokes' theorem we have

$$\begin{aligned} & \int_{\{r < \psi < 1\}} -\varphi(dd^c \psi) \wedge (dd^c \varphi)^{n-1} + \int_{\{r < \psi < 1\}} (\psi - r)(dd^c \varphi)^n \\ & \leq \lim_{t \searrow r} \int_{\{r < \psi < 1\}} -\varphi dd^c(\max \psi, t) \wedge (dd^c \varphi)^{n-1} + \int_{\{r < \psi < 1\}} (\psi - r)(dd^c \varphi)^n \\ & = \lim_{t \searrow r} \left[\int_{\{\psi=1\}} -\varphi d^c(\max \psi, t) \wedge (dd^c \varphi)^{n-1} \right. \\ & \quad \left. + \int_{\{r < \psi < 1\}} d \max(\psi, t) \wedge d^c \varphi \wedge (dd^c \varphi)^{n-1} + \int_{\{r < \psi < 1\}} (\psi - r)(dd^c \varphi)^n \right] \\ & = \lim_{t \searrow r} \left[\int_{\{\psi=1\}} -\varphi d^c(\psi, t) \wedge (dd^c \varphi)^{n-1} \right. \\ & \quad \left. + \int_{\{\psi=1\}} (\max(\psi, t) - r) d^c \varphi \wedge (dd^c \varphi)^{n-1} \right. \\ & \quad \left. - \int_{\{\psi=r\}} (\max(\psi, t) - r) d^c \varphi \wedge (dd^c \varphi)^{n-1} \right. \\ & \quad \left. + \int_{\{r < \psi < 1\}} (\psi - \max(\psi, t))(dd^c \varphi)^n \right] \\ & = \int_{\{\psi=1\}} -\varphi d^c \psi \wedge (dd^c \varphi)^{n-1} + \int_{\{\psi=1\}} (\psi - r) d^c \varphi \wedge (dd^c \varphi)^{n-1}. \end{aligned}$$

Since the right-hand side is uniformly bounded in r , $s < r < 1$, and since $\varphi \leq 0$, we get

$$\int_{\{s < \psi < 1\}} |\varphi| dd^c \psi \wedge (dd^c \varphi)^{n-1} + \int_{\{s < \psi < 1\}} (\psi - s)(dd^c \varphi)^n < +\infty,$$

where each member is nonnegative.

3. An example. The following example shows that the convergence factor $\psi - s$ really is needed in the Theorem. There are functions φ and ψ such that $\int_{\{s < \psi < 1\}} (dd^c \varphi)^n = +\infty$, where φ can be taken to be plurisubharmonic and bounded.

Define $u(z) = -\sqrt{|z|^2 - \frac{1}{2}}$; then u is subharmonic on $\{z \in \mathbb{C}; \frac{1}{2} < |z|^2 < 1\}$, $-1/\sqrt{2} < u < 0$, and $\Delta u = (1 - |z|^2)/(|z|^2 - \frac{1}{2})^{3/2}$.

Define $\psi(z, w) = |z|^2 + \frac{2}{3}|w|^8$, and let $0 < \eta < \frac{1}{20}$, and put

$$\varphi_\eta(z, w) = \max \left[-\frac{1}{2} \sqrt{(1+\eta)^2 |z|^2 - \frac{1}{2} + |w|^2}, |w|^2 + (1+\eta)|w|^4 - \sqrt{\frac{3}{2}(\eta + \eta^2/2)} \right].$$

Then φ_η is plurisubharmonic on $\{\frac{1}{2} < \psi < (20/21)^2\}$ for, if $(1+\eta)^2 |z|^2 \leq \frac{1}{2}$, we have, when $\psi(z, w) > \frac{1}{2}$,

$$\frac{2}{3}|w|^8 > \frac{1}{2} - |z|^2 > \frac{1}{2}(1 - (1+\eta)^{-2}) = \frac{1}{2}(2\eta + \eta^2)/(1+\eta)^2 = (\eta + \eta^2/2)(1+\eta)^2$$

so

$$(1+\eta)|w|^4 > \sqrt{\frac{3}{2}(\eta + \eta^2/2)}.$$

Assume now that $\frac{1}{2} < |z|^2 + \frac{2}{3}|w|^8 < (20/21)^2$ and that

$$\sqrt{\frac{3}{5}} \frac{\sqrt{\eta + \eta^2/2}}{1+\eta} < |w|^4 < \sqrt{\frac{2}{3}} \frac{39}{40} \frac{\sqrt{\eta + \eta^2/2}}{1+\eta}.$$

Then

$$\varphi_\eta = -\frac{1}{2} \sqrt{(1+\eta)^2 |z|^2 - \frac{1}{2} + |w|^2}$$

if

$$\begin{aligned} & -\frac{1}{2} \sqrt{(1+\eta)^2 |z|^2 - \frac{1}{2}} > (1+\eta)|w|^4 - \sqrt{\frac{3}{2}(\eta + \eta^2/2)} \\ \Leftrightarrow & (1+\eta)^2 |z|^2 < \frac{1}{2} + 4 \left(\sqrt{\frac{3}{2}(\eta + \eta^2/2)} - (1+\eta)|w|^4 \right)^2, \end{aligned}$$

which holds if

$$\left(\frac{1}{2} - \frac{2}{3}|w|^8 \right) (1+\eta)^2 < (1+\eta)^2 |z|^2 < \frac{1}{2} + 4 \left(\sqrt{\frac{3}{2}(\eta + \eta^2/2)} - (1+\eta)|w|^4 \right)^2$$

which is a well-defined domain if

$$\begin{aligned} & \eta + \eta^2/2 - \frac{2}{3}|w|^8 (1+\eta)^2 \\ & < 6(\eta + \eta^2/2) - 8\sqrt{\frac{3}{2}(\eta + \eta^2/2)}(1+\eta)|w|^4 + 4(1+\eta)^2 |w|^8 \\ \Leftrightarrow & 0 < 5(\eta + \eta^2/2) + \frac{14}{3}(1+\eta)^2 |w|^8 - 8\sqrt{\frac{3}{2}(\eta + \eta^2/2)}(1+\eta)|w|^4, \end{aligned}$$

which holds with $|w|^4$ in the interval above—the right-hand side is then strictly larger than

$$5 \left(\eta + \frac{\eta^2}{2} \right) + \frac{14}{3}(1+\eta)^2 \frac{3}{5} \frac{\eta + \eta^2/2}{(1+\eta)^2} - 8 \cdot \frac{39}{40} \left(\eta + \frac{\eta^2}{2} \right) = 0.$$

Therefore,

$$\int_{\frac{1}{2} < |z|^2 + \frac{2}{3}|w|^8 < (20/21)^2} (dd^c \varphi_\eta)^2 \geq c \int_{A_\eta} \frac{dz dw}{((1+\eta)^2 |z|^2 - \frac{1}{2})^{3/2}},$$

where

$$A_\eta = \left\{ \begin{aligned} & \left(\frac{1}{2} - \frac{2}{3}|w|^8 \right) (1+\eta)^2 < (1+\eta)^2 |z|^2 < \frac{1}{2} + 4 \left(\sqrt{\frac{3}{2}(\eta + \eta^2/2)} - (1+\eta)|w|^4 \right)^2, \\ & \frac{\sqrt{\eta + \eta^2/2}}{1+\eta} \alpha < |w|^4 < \beta \frac{\sqrt{\eta + \eta^2/2}}{1+\eta}, \end{aligned} \right.$$

and where we have chosen $\sqrt{\frac{3}{5}} < \alpha < \beta < \frac{39}{40} \sqrt{\frac{2}{3}}$ and where c is a strictly positive constant. The right-hand side is not smaller than

$$\begin{aligned} c \left\{ \left(\alpha^2 - \frac{3}{5} \right) + \frac{39}{40} \sqrt{\frac{2}{3}} - \beta \right\} \int \frac{(\eta + \eta^2/2) dw}{2^3 (\sqrt{\frac{3}{2}}(\eta + \eta^2/2) - (1 + \eta)|w|^4)^3} \\ \frac{\sqrt{\frac{3}{5}(\eta + \eta^2/2)}}{1 + \eta} < |w|^4 < \sqrt{\frac{2}{3}} \frac{39}{40} \frac{\sqrt{\eta + \eta^2/2}}{1 + \eta} \\ &\geq d(\eta + \eta^2/2)^{1-3/2+1/4} \\ &= \frac{d}{(\eta + \eta^2/2)^{1/4}} \rightarrow +\infty, \quad \eta \rightarrow 0 \quad (d \text{ pos. const.}). \end{aligned}$$

Furthermore, $-\frac{1}{2} < \varphi_\eta \leq 2$ so if we put

$$\varphi = \sum_{j=20}^{\infty} \frac{1}{j^2} \varphi_{1/j^4},$$

then

$$\varphi \in \text{PSH} \cap L^\infty \left(\frac{1}{2} < |z|^2 + \frac{2}{3}|w|^8 < (20^4/(20^4 + 1))^2 \right)$$

and

$$\int_{\frac{1}{2} < |z|^2 + \frac{2}{3}|w|^8 < (20/21)^2} (dd^c \varphi)^2 = +\infty.$$

REMARK. It is possible to modify the φ_η 's to get the function $\varphi \in C^\infty$ on $\Omega \setminus \{w = 0\}$. We do not know if one can modify φ to be C^∞ on Ω but still have $\int_{\Omega \setminus K} (dd^c \varphi)^2 = +\infty$.

REFERENCES

1. E. Bedford and B. A. Taylor, *A new capacity for plurisubharmonic functions*, Acta Math. **199** (1982), 1–40.
2. U. Cegrell, *Removable singularity sets for plurisubharmonic functions and related problems*, Proc. London Math. Soc. (3) **36** (1978), 310–336.
3. J. E. Fornæss and N. Sibony, *Plurisubharmonic functions on ring domains*, Complex Analysis (S. G. Krantz, Ed.), Lecture Notes in Math., vol. 1268, Springer, Berlin and New York.
4. P. Griffiths, *Two theorems on extension of holomorphic mappings*, Invent. Math. **14** (1971), 27–62.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF UMEA, 901 97 UMEA, SWEDEN