THE VALIDITY OF BEURLING THEOREMS IN POLYDISCS

V. MANDREKAR

(Communicated by Paul S. Muhly)

ABSTRACT. This paper gives necessary and sufficient conditions for an invariant subspace $\mathcal{M}$ of $H^2(T^2)$ to be of the form $qH^2(T^2)$ (q inner) in terms of double commutativity of the shifts. Recent results in [8] follow directly from our work. Relation to the work in [1] is also discussed.

In [5], Rudin gave an example of a shift-invariant subspace of $H^2(T^2)$ which is not of the form $qH^2(T^2)$, where q is an inner function. In addition, [5] gives an example of an "outer function" f for which $\mathcal{M}_f$, the smallest invariant subspace generated by f is not equal to $H^2(T^2)$. Motivated from the prediction theory for random fields, Soltani [8] gave complicated necessary and sufficient conditions for an "outer function" in the sense of [5] to satisfy $\mathcal{M}_f = H^2(T^2)$. Our purpose here is to use the Theorem of Halmos for two commuting isometries due to Słocinski [7] to obtain necessary and sufficient conditions on the invariant subspace $\mathcal{M}$ of $H^2(T^2)$ to be of the form $qH^2(T^2)$ (q inner). As a consequence of this result we obtain necessary and sufficient conditions on $\mathcal{M}_f$ for an outer function f , to be $H^2(T^2)$. We use results of [3] to relate our conditions to those in [8]. As a by-product of our main theorem we show that all invariant subspaces unitarily equivalent to $qH^2(T^2)$ are exactly of the same form. The last result relates to some recent work of Agrawal, Clark, and Douglas [1], where the problem of unitary equivalence of invariant subspaces of $H^2(T^n)$ was solved under some sufficient conditions. Their case does not include subspaces of the type considered here. We thank the referee and the editor for bringing the work in [1] to our attention. We begin with some notation.

Let $\mathbf{Z}$ be the set of integers. We denote by m, n etc. the elements of $\mathbf{Z}$. Let U be the open unit disc and T the boundary of U in the complex plane $\mathbf{C}$. Let $\mathbf{Z}^2, \mathbf{C}^2, U^2$ and T^2 be the respective cartesian products and σ_2 the normalized Lebesgue measure on T^2 . For $p > 0$, we denote by $L^p(T^2, \sigma_2)$ the usual Lebesgue space of the equivalence class of p -integrable functions and

$$H^p(U^2) = \left\{ f : f : U^2 \rightarrow \mathbf{C} \text{ analytic and } \sup_{0 \leq r \leq 1} \int_T |f_r(\mathbf{t})|^p d\sigma_2 < \infty \right\}.$$

Here $f_r(\mathbf{t}) = f(z)$ with $z = r\mathbf{t}$. Let $\mathbf{z} = (z_1, z_2) = (r_1 e^{i\theta_1}, r_2 e^{i\theta_2})$ and $\mathbf{t} = (e^{i\theta_1}, e^{i\theta_2})$, then $P(\mathbf{z}, \mathbf{t}) = P_{r_1}(\theta_1 - \theta_2)$. $P_{r_1}(\theta_1 - \theta_2)$ is called Poisson kernel with

---

Received by the editors March 30, 1987.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 32A30, 60G10.

Key words and phrases. Polydisc, Beurling Theorem.

Supported in part by ONR N00014-85-K-0150 and the Air Force Office of Scientific Research Contract No. F49620 85 C 0144.

$P_r(\theta) = (1-r^2)/(1-2r \cos \theta + r^2)$. It is known that for $f \in H^p(U^2)$, $\lim_{r \rightarrow 1} f_r(\mathbf{t}) = f^*(\mathbf{t})$ exists and is in $L^p(T^2, \sigma_2)$. For $f \in L^p(T^2, \sigma_2)$, let $f^e(z) = \int_{T^2} P(\mathbf{z}, \mathbf{t}) f(\mathbf{t}) d\sigma_2$, then $f^e \in H^p(U^2)$. In case $p = 2$, $f^e \in H^2(U^2)$ if $f \in L^2(T^2, \sigma_2)$ and $f(\mathbf{t}) = \sum_{m=0}^\infty \sum_{n=0}^\infty a_{mn} t_1^m t_2^n$ and $f^e = \sum_{m=0}^\infty \sum_{n=0}^\infty a_{mn} z_1^m z_2^n$. Conversely, every $f \in H^2(U^2)$ has this form and $f^*(\mathbf{t}) = \sum_{m=0}^\infty \sum_{n=0}^\infty a_{mn} t_1^m t_2^n$. For further information see [5].

In order to characterize invariant subspaces of the form qH^2 in terms of the section of the shifts on it, we note that subspaces of the form qH^2 can be represented as

$$(1) \quad qH^2 = \sum_{m=0}^\infty \sum_{n=0}^\infty \bigoplus V_1^m V_2^n(M)$$

where M equals the span of q in $H^2(T^2)$ and V_i is the multiplication by t_i on $H^2(T^2)$ with $\mathbf{t} = (t_1, t_2) \in T^2$. As V_1 commutes with V_2 (in short, $V_1 \sim V_2$), we get from (1) and Theorem 1 (i) $\Rightarrow$ (ii) of [7] that V_1 and V_2 are doubly commuting (i.e. $V_1 \sim V_2, V_1 \sim V_2^*$). In fact, we have

2. THEOREM. *An invariant subspace $\mathcal{M} \neq \{0\}$ of $H^2(T^2)$ is of the form $q \cdot H^2$ with q inner function if and only if V_1 and V_2 are doubly commuting on $\mathcal{M}$.*

PROOF. Necessity was proved above. To prove the sufficiency we get, in view of Theorem 1 ((ii) $\Rightarrow$ (iv)) [7],

$$(3) \quad \mathcal{M} = \sum_{m=0}^\infty \sum_{n=0}^\infty \bigoplus V_1^m V_2^n (R_1^\perp \cap R_2^\perp),$$

where $R_i^\perp = \mathcal{M} \ominus V_i \mathcal{M}$. From Theorem 1 (i) $\Leftrightarrow$ (iv) and Corollary 1 of [7, p. 256] we get that $R_1^\perp \cap R_2^\perp = \{0\}$ implies $\mathcal{M} = \{0\}$ giving contradiction. Hence $R_1^\perp \cap R_2^\perp \neq \{0\}$. We shall now prove that $R_1^\perp \cap R_2^\perp$ is one-dimensional. The above argument shows that dimension of $R_1^\perp \cap R_2^\perp$ is at least one. It remains to prove that it is not greater than one. Let $q_1, q_2 \in R_1^\perp \cap R_2^\perp$, then $\int_{T^2} t_1^m t_2^n q_1 q_2 d\sigma_2 = 0$ for $m, n > 0$. As $V_2(R_1^\perp) \subseteq (R_1^\perp)$ [7, Theorem 1 (iii)], we get $\int_{T^2} t_2^n q_1 \overline{t_1^m q_2} d\sigma_2 = 0$ for all $n > 0$ and $m > 0$. Since $t_1^m = \overline{t_1^{-m}}$, by the symmetry of the problem, we get for $(m, n) \neq (0, 0)$ $\int t_1^m t_2^n q_1 \overline{q_2} d\sigma_2 = 0$. Since $q_1 \overline{q_2} \in L^1(T^2, \sigma_2)$ we get $q_1 \overline{q_2} = c$ a.e. σ_2 . Suppose now that $q_1, q_2 \in R_1^\perp \cap R_2^\perp$, $q_1 \neq 0$, $q_2 \neq 0$ and $q_1 \perp q_2$. This implies that $|q_1|^2 = c_1 \neq 0$ a.e. σ_2 , $|q_2|^2 = c_2 \neq 0$ a.e. σ_2 but $q_1 \overline{q_2} = 0$. The last statement follows from the fact $q_1 \perp q_2$ and a.e. σ_2 , $q_1 \overline{q_2} = c$ giving $c = 0$ (integrating with respect to σ_2). This is impossible. Hence $R_1^\perp \cap R_2^\perp$ is one dimensional. Also q generating $R_1^\perp \cap R_2^\perp$ is an inner function. Assume $|q| = 1$ a.e. choosing q of norm 1. Now (3) gives the result.

In a recent paper [1], Agrawal, Clark and Douglas study the question of unitary equivalence of invariant subspaces of H^2 of polydiscs. Under the assumption of full range of an invariant subspace $\mathcal{M}$ (for definition see [1, (2), p. 5]), they show that all invariant subspaces $\mathcal{N}$ of H^2 unitarily equivalent to $\mathcal{M}$ are of the form $\varphi \mathcal{M}$ (φ an inner function). Unfortunately, subspaces of the form qH^2 do not have full range [1, Remark 3, p. 6] unless q is constant. We use our result to study all invariant subspaces unitarily equivalent to qH^2 . We consider the case of subspaces of $H^2(T^2)$.

If $\mathcal{M}$ is an invariant subspace unitarily equivalent to qH^2 for some inner function, then in view of Theorem 2, we get V_1 and V_2 are doubly commuting on $\mathcal{M}$. Hence $\mathcal{M} = q'H^2$ for some inner function q' by Theorem 2 with $|q'(t)| = 1$ a.e. σ_2 . We thus have the following.

4. THEOREM. (a) *The class of all invariant subspaces of the form qH^2 , q inner function with $|q(t)| = 1$ a.e. σ_2 are unitarily equivalent.*

(b) *Any invariant subspace $\mathcal{M}$ of $H^2(T^2)$ unitarily equivalent to $\mathcal{N} = qH^2$ of the above form is of the same form.*

REMARK. Note that in (b) $\mathcal{M} = q'/q\mathcal{N}$. Comparing this with Lemma 1 of [1] we get that the function $\varphi = q'/q$ with q' and q relatively prime in the sense of [1, p. 4].

In [6], Rudin gives an example of the subspace $\mathcal{M} = q'/q\mathcal{N}$ with q', q inner but not relatively prime. We should also note that there exists subspaces of the form qH^2 which do not satisfy analytic condition used in §3 of [1].

The example of Rudin [5] mentioned in the introduction shows that there exists $f \in H^2$ such that

$$\mathcal{M}_f = \overline{sp}\{V_1^m V_2^n f; m, n \geq 0\}$$

is not of the form qH^2 . The following Corollary gives conditions on $\mathcal{M}_f$ for this to hold.

5. COROLLARY. $\mathcal{M}_f = qH^2(T^2)$ if and only if V_1 and V_2 are doubly commuting on $\mathcal{M}_f$.

Following Helson [2], we say that a function g is H -outer if $\mathcal{M}_g = H^2(T^2)$.

6. COROLLARY. A function $f \in H^2(T^2)$ has the property $f = q \cdot g$ with q inner and g H -outer if and only if V_1 and V_2 doubly commute on $\mathcal{M}_f$.

PROOF. By Corollary 5, only if part follows as $\mathcal{M}_f = qH^2(T^2)$. To prove the converse, we note that by Corollary 5, $\mathcal{M}_f = qH^2(T^2)$ given $f = q \cdot g$, $g \in H^2(T^2)$. Hence $\mathcal{M}_f = q \cdot \mathcal{M}_g$ giving g is H -outer.

In [5, p. 73], a function $f \in H^2(U^2)$ is called outer (we call it R -outer) if $\log |f(z)| = \int_{T^2} P(z, t) \log |f^*| d\sigma_2$. Given a function $f \in H^2(T^2)$, we denote by $f^e \in H^2(U^2)$ the function given by $\int_{T^2} P(z, t) f(t) d\sigma_2$. In this case we note that $(f^e)^* = f$. It is already known [5, Theorem 4.4.6] that f is H -outer then f^e is R -outer. From this we get in view of Corollary 5 the following.

7. COROLLARY. Let g be H -outer then g^e is R -outer and V_1 and V_2 doubly commute on $\mathcal{M}_g$.

We now prove the converse of Corollary 7. Assume now that V_1, V_2 doubly commute on $\mathcal{M}_f$ and f^e is R -outer then by Corollaries 6 and 7, the definition of f^e, g^e and the fact that $|q| = 1$ we get $f^e = pg^e$ with $|p| = 1$. Thus we get that the slice function $f_w^e(\lambda) = p_w(\lambda)g_w^e(\lambda)$. Using Lemma 4.4.4(a) of [5] and the uniqueness of outer function [2, p. 13] we get $p_w(\lambda) = 1$ for all w and λ giving $p = 1$ i.e., $f = g$.

Combining this with Corollary 7 gives

8. COROLLARY. *Let $f \in H^2(T^2)$ then $\mathcal{M}_f = H^2(T^2)$ if and only if f^e is R -outer and V_1 and V_2 doubly commute.*

In view of Theorem 4.2 of [3], we get that Corollary 7 includes Beurling Theorem proved in [8, Theorem 1.5]. Now using essentially classical techniques [4, 3] one can derive associated results in prediction theory given in [8].

REFERENCES

1. O. P. Agrawal, D. N. Clark and R. G. Douglas, *Invariant subspaces in the polydisk*, Pacific J. Math. **121** (1986), 1–11.
2. H. Helson, *Lectures on invariant subspaces*, Academic Press, New York, 1964.
3. G. Kallianpur and V. Mandrekar, *Non-deterministic random fields and Wold and Halmos decompositions for commuting isometries*, Prediction Theory and Harmonic Analysis (V. Mandrekar and H. Salehi, eds.), North-Holland, Amsterdam, 1983.
4. P. Masani, *Shift-invariant spaces and prediction theory*, Acta Math. **107** (1962), 275–290.
5. W. Rudin, *Function theory in polydiscs*, Benjamin, New York, 1979.
6. —, *Invariant subspaces of H^2 on a torus*, J. Functional Analysis **61** (1985), 378–384.
7. M. Słociński, *On Wold-type decomposition of a pair of commuting isometries*, Ann. Polon. Math. **37** (1980), 255–262.
8. A. R. Soltani, *Extrapolation and moving average representation for stationary random fields and Beurling's Theorem*, Ann. Probab. **12** (1984), 120–132.

DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, EAST LANSING,
MICHIGAN 48824