THE CARDINALITY OF REDUCED POWER SET ALGEBRAS

ALAN D. TAYLOR

(Communicated by Thomas J. Jech)

ABSTRACT. We prove a general result on the cardinality of reduced powers of structures via filters that has several consequences including the following: if I is a uniform, countably complete ideal on the real line $\mathcal{R}$ and $\mathcal{B}$ is the Boolean algebra of subsets of $\mathcal{R}$ modulo I , then $|\mathcal{B}| > 2^{\aleph_0}$ and if $2^\nu \leq 2^{\aleph_0}$ for all $\nu < 2^{\aleph_0}$ then $|\mathcal{B}| = 2^{2^{\aleph_0}}$. This strengthens some results of Kunen and Pelc [7] and Prikry [8] obtained by Boolean ultrapower techniques. Our arguments are all combinatorial and some applications are included.

1. Introduction. Our notation and terminology are fairly standard. If A and B are sets then ${}^A B$ denotes the set of all functions mapping A into B and $\mathcal{P}(A)$ denotes the power set of A . Throughout, κ and λ will denote infinite cardinals. An ideal I on κ is a collection of subsets of κ that is closed under subset formation and finite unions. All the ideals we will consider are assumed to be proper ($I \neq \mathcal{P}(\kappa)$) but not necessarily κ -complete (i.e. closed under unions of size less than κ) or even uniform (i.e. containing all subsets of κ of size less than κ). If I is an ideal on κ then I^* denotes the filter on κ that is dual to I ; that is, $X \in I^*$ iff $\kappa - X \in I$, and I^+ denotes $\mathcal{P}(\kappa) - I$.

If I is an ideal on κ and B is an arbitrary set then the reduced power of B (via I^*), denoted $\prod_{\kappa} B/I^*$, is the set of all equivalence classes of functions in ${}^{\kappa} B$ where we identify two such functions f and g if $\{\alpha < \kappa: f(\alpha) \neq g(\alpha)\} \in I$. The Boolean algebra $\mathcal{P}(\kappa)/I$ is the set of all equivalence classes of subsets of κ with the induced ordering from $\subseteq$, where we identify two such subsets X and Y if their symmetric difference $X \Delta Y$ is in I .

Since we will be comparing $\mathcal{P}(\kappa)/I$ and $\prod_{\kappa} \kappa/I^*$ it is worth pointing out the following (see [3 or 1, p. 338]). The function which associates to each $X \subseteq \kappa$ its characteristic function induces a natural bijection between $\mathcal{P}(\kappa)/I$ and $\prod_{\kappa} 2/I^*$ where, of course, $2 = \{0, 1\}$. Hence $|\mathcal{P}(\kappa)/I| = |\prod_{\kappa} 2/I^*| \leq |\prod_{\kappa} \kappa/I^*|$. In general, the inequality cannot be replaced by equality since trivially $|\prod_{\kappa} \kappa/I^*| \geq \kappa$ (consider the constant functions from κ to κ) but $|\mathcal{P}(\kappa)/I| = 2$ if I is a prime (i.e. maximal) ideal on κ .

2. The main results. At the heart of our first theorem is the following trivial observation: there is a natural bijection between κ -sequences of subsets of λ and λ -sequences of subsets of κ . This correspondence is given by $(X_\alpha: \alpha < \kappa) \leftrightarrow (Y_\beta: \beta < \lambda)$ where $\alpha \in Y_\beta$ iff $\beta \in X_\alpha$. The proof of the following result consists of simply verifying that if I is a λ^+ -complete ideal on κ then this correspondence induces a (well-defined) bijection between $\prod_{\kappa} \mathcal{P}(\lambda)/I^*$ and ${}^\lambda(\mathcal{P}(\kappa)/I)$.

Received by the editors March 20, 1987.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 03E05. Secondary 06E05.

©1988 American Mathematical Society
0002-9939/88 \$1.00 + \$.25 per page

THEOREM 2.1. *If I is a λ^+ -complete ideal on κ then*

$$\left| \prod_{\kappa} {}^{\lambda}2/I^* \right| = \left| \prod_{\kappa} 2/I^* \right|^{\lambda}.$$

PROOF. Define $\phi: \prod_{\kappa} {}^{\lambda}2/I^* \rightarrow {}^{\lambda}(\prod_{\kappa} 2/I^*)$ as follows: if $[f] \in \prod_{\kappa} {}^{\lambda}2/I^*$, then $f: \kappa \rightarrow {}^{\lambda}2$ and we set

$$\phi([f]) = \langle [f_{\beta}]: \beta < \lambda \rangle$$

where $f_{\beta}: \kappa \rightarrow 2$ is given by $f_{\beta}(\alpha) = f(\alpha)(\beta)$. Note that $[f_{\beta}] \in \prod_{\kappa} 2/I^*$.

To complete the proof of the theorem it suffices to show that ϕ is a well-defined bijection.

CLAIM 1. ϕ is well defined.

PROOF. Suppose that $f, g: \kappa \rightarrow {}^{\lambda}2$ and $[f] = [g]$. Let $X = \{\alpha < \kappa: f(\alpha) = g(\alpha)\}$. Then $X \in I^*$. Fix $\beta < \lambda$ and let $Y_{\beta} = \{\alpha < \kappa: f_{\beta}(\alpha) = g_{\beta}(\alpha)\}$. We claim that $X \subseteq Y_{\beta}$. That is, if $\alpha \in X$ then $f(\alpha) = g(\alpha)$ so $f(\alpha)(\beta) = g(\alpha)(\beta)$. But then $f_{\beta}(\alpha) = g_{\beta}(\alpha)$ so $\alpha \in Y_{\beta}$ as desired. But now since $X \in I^*$ and $X \subseteq Y_{\beta}$ we have $Y_{\beta} \in I^*$ and so $[f_{\beta}] = [g_{\beta}]$. Since β was arbitrary this shows that $\langle [f_{\beta}]: \beta < \lambda \rangle = \langle [g_{\beta}]: \beta < \lambda \rangle$. Thus $\phi([f]) = \phi([g])$ and so ϕ is well defined as claimed.

CLAIM. ϕ is one-to-one.

PROOF. Suppose that $f, g: \kappa \rightarrow {}^{\lambda}2$ and assume that $\phi([f]) = \phi([g])$. Then for every $\beta < \lambda$ we have $[f_{\beta}] = [g_{\beta}]$ and so $Y_{\beta} \in I^*$ where $Y_{\beta} = \{\alpha < \kappa: f_{\beta}(\alpha) = g_{\beta}(\alpha)\}$. Let $Y = \bigcap_{\beta < \lambda} Y_{\beta}$. Since I is λ^+ -complete we have $Y \in I^*$. Now let $X = \{\alpha < \kappa: f(\alpha) = g(\alpha)\}$. We claim that $Y \subseteq X$. That is, if $\alpha \in Y$ then for every $\beta < \lambda$ we have $\alpha \in Y_{\beta}$, and so $f_{\beta}(\alpha) = g_{\beta}(\alpha)$. But then for every $\beta < \lambda$ we have $f(\alpha)(\beta) = g(\alpha)(\beta)$ and so the functions $f(\alpha)$ and $g(\alpha)$ are equal. Hence $\alpha \in X$. Now, since $Y \in I^*$ and $Y \subseteq X$ we have $X \in I^*$ and so $[f] = [g]$ as desired.

CLAIM 3. ϕ is onto.

PROOF. Suppose $\langle [f_{\beta}]: \beta < \lambda \rangle \in {}^{\lambda} \prod_{\kappa} 2/I^*$. Define $f: \kappa \rightarrow {}^{\lambda}2$ by $f(\alpha)(\beta) = f_{\beta}(\alpha)$. Then $[f] \in \prod_{\kappa} {}^{\lambda}2/I^*$ and clearly $\phi([f]) = \langle [f_{\beta}]: \beta < \lambda \rangle$. $\square$

Specializing Theorem 2.1 to the case where $\kappa = 2^{\lambda}$ and making use of some known results yields the following.

THEOREM 2.2. *Suppose I is a λ^+ -complete ideal on $\kappa = 2^{\lambda}$. Then*

- (i) $|\mathcal{P}(\kappa)/I|^{\lambda} = |\prod_{\kappa} \kappa/I^*| \geq \kappa$, and
- (ii) if I is also uniform (i.e. $[\kappa]^{<\kappa} \subseteq I$), then $|\mathcal{P}(\kappa)/I| > \kappa$.

PROOF. Part (i) follows immediately from Theorem 2.1 and our comments regarding $\mathcal{P}(\kappa)/I$ and $\prod_{\kappa} 2/I^*$. For part (ii) we need the following well-known result (see [3, p. 204]): for every infinite cardinal γ there exists a family $\mathcal{F} \subseteq {}^{\gamma}\gamma$ with $|\mathcal{F}| = \gamma^+$ and $|\{\alpha < \gamma: f(\alpha) = g(\alpha)\}| < \gamma$ whenever $f, g \in \mathcal{F}$ and $f \neq g$. An immediate consequence of this is the fact that if I is a uniform ideal on κ then $|\prod_{\kappa} \kappa/I^*| \geq \kappa^+$. To complete the proof of (ii), let $\mu = |\mathcal{P}(\kappa)/I|$ and suppose for contradiction that $\mu \leq \kappa$. Then

$$\kappa^+ \leq \left| \prod_{\kappa} \kappa/I^* \right| = |\mathcal{P}(\kappa)/I|^{\lambda} = \mu^{\lambda} \leq \kappa^{\lambda} = (2^{\lambda})^{\lambda} = 2^{\lambda} = \kappa. \quad \square$$

Finally, if we specialize still further and again use some known results we obtain the following.

THEOREM 2.3. *Suppose I is an ω_1 -complete ideal on the real line $\mathcal{R}$. Then*

- (i) $|\mathcal{P}(\mathcal{R})/I| = |\prod_{\mathcal{R}} \mathcal{R}/I^*| = |\prod_{\mathcal{R}} \mathcal{R}/I^*|^\omega$,
- (ii) *if I is uniform then $|\mathcal{P}(\mathcal{R})/I| > 2^{\aleph_0}$, and*
- (iii) *if I is uniform and $2^\nu \leq 2^{\aleph_0}$ for all $\nu < 2^{\aleph_0}$ then $|\mathcal{P}(\mathcal{R})/I| = 2^{2^{\aleph_0}}$.*

PROOF. Part (i) follows immediately from Theorem 2.2(i) and the following result of Comfort-Hager [2] and Monk-Sparks [6]: if B is an infinite ω_1 -complete Boolean algebra, then $|B|^\omega = |B|$. Since $\mathcal{P}(\mathcal{R})/I$ is ω_1 -complete and infinite, this gives us that $|\mathcal{P}(\mathcal{R})/I| = |\mathcal{P}(\mathcal{R})/I|^\omega$.

Part (ii) is an immediate consequence of Theorem 2.2(ii).

Part (iii) requires the following well-known result of Tarski (see [3, p. 204]): if γ is an infinite cardinal and $2\gamma = \gamma$ then there exists a family $\mathcal{F} \subseteq {}^\gamma\gamma$ with $|\mathcal{F}| = 2^\gamma$ and $|\{\alpha < \gamma: f(\alpha) = g(\alpha)\}| < \gamma$ whenever $f, g \in \mathcal{F}$ and $f \neq g$. (To see this, let $\langle X_\beta^\alpha: \beta < \gamma_\alpha \rangle$ be an enumeration of $\mathcal{P}(\alpha)$ in order-type $\gamma_\alpha \leq \gamma$ for each $\alpha < \gamma$ and for each $X \subseteq \gamma$ define $f_X: \gamma \rightarrow \gamma$ by $f_X(\alpha) = \beta$ iff $X \cap \alpha = X_\beta^\alpha$.) In the context of (iii), with $\gamma = 2^{\aleph_0}$ ($= |\mathcal{R}|$), this gives us that $|\prod_{\mathcal{R}} \mathcal{R}/I^*| \geq 2^{2^{\aleph_0}}$, and so (iii) now follows from (i). $\square$

The second equality in part (i) of Theorem 2.3 should be compared with Keisler's result [5] that $|\prod \mathcal{R}/D| = |\prod \mathcal{R}/D|^\omega$ for any countably incomplete ultrafilter D on ω . Part (ii) of Theorem 2.3 eliminates some saturation requirements from an old (unpublished) result of Kunen (recently rediscovered by Pelc [7]) and part (iii) does the same for a result of Prikry [8]. The proofs of Kunen, Pelc and Prikry all employed Boolean ultrapowers.

3. Some applications. We conclude with a few applications of the results in §2.

APPLICATION 1. We begin with an application that is due as much to Andrzej Pelc (see [7]) as to the present author. It answers the following question that we asked in [9, Problem R, p. 51]: Suppose we have countably many ω_1 -complete uniform ideals on the real line $\mathcal{R}$. Can we then find a subset X of $\mathcal{R}$ such that neither X nor $\mathcal{R} - X$ belongs to any of the ideals in our collection? An affirmative answer to this can be derived as follows. By Theorem 5.4.2 of [9], it suffices to show that if I is an ω_1 -complete uniform ideal on $\mathcal{R}$ then $\mathcal{P}(\mathcal{R})/I$ has no countable dense set (in the forcing theoretic sense). Assume we have such an I and $\{X_n: n \in \omega\} \subseteq I^+$. To each distinct $[Y] \in \mathcal{P}(\mathcal{R})/I$ we associate $\{n \in \omega: [X_n] \leq [Y]\}$. Since $|\mathcal{P}(\mathcal{R})/I| > 2^{\aleph_0}$ by Theorem 2.3(ii), we get $[Y_1], [Y_2] \in \mathcal{P}(\mathcal{R})/I$ with $[Y_1] = [Y_2]$ and $[X_n] \leq [Y_1]$ iff $[X_n] \leq [Y_2]$ for every $n \in \omega$. But this shows that $\{X_n: n \in \omega\}$ is not a dense set since either $Y_1 - Y_2 \in I^+$ or $Y_2 - Y_1 \in I^+$ and neither contains X_n (modulo I) for any $n \in \omega$.

APPLICATION 2. Theorem 2.3(ii) also yields yet another proof that not every set of reals is Lebesgue measurable. To see this, assume the contrary and let $I = \{X \subseteq \mathcal{R}: \mu(X) = 0\}$. Then I is clearly ω_1 -complete and it is uniform since every set of reals of cardinality less than the continuum has outer measure zero. Theorem 2.3(ii) now implies that $|\mathcal{P}(\mathcal{R})/I| > 2^{\aleph_0}$, and this contradicts the well-known fact that every measurable set differs from a Borel set by a set of measure zero (since there are only $2^{\aleph_0}$ Borel sets).

APPLICATION 3. We showed above that a consequence of Theorem 2.3 is the fact that no ω_1 -complete uniform ideal I on $\mathcal{R}$ can yield a Boolean algebra $\mathcal{P}(\mathcal{R})/I$

with a countable dense set. A natural question (again related to versions of Ulam's problem in [9 and 10]) is whether a dense set of size $\aleph_1$ is possible. Woodin [11] has shown that relative to large cardinals an affirmative answer is consistent. In his model CH holds. As a final application of Theorem 2.2 we consider the question of whether CH is necessary and prove the following: if $\mathcal{R}$ carries an ω_1 -complete uniform ideal I so that $\mathcal{P}(\mathcal{R})/I$ has a dense set of size $\aleph_1$ then either $2^{\aleph_0} = \aleph_1$ or $2^{\aleph_0} > \aleph_{\omega_1}$. To see this we argue as follows. Suppose I is an ω_1 -complete, uniform ideal on $\mathcal{R}$ and D is a dense set of size $\aleph_1$ for $\mathcal{P}(\mathcal{R})/I$. Define $\phi: \mathcal{P}(\mathcal{R})/I \rightarrow \mathcal{P}(D)$ by $\phi([A]) = \{d \in D: d \leq [A]\}$. It is easy to see that ϕ is one-to-one and so $|\mathcal{P}(\mathcal{R})/I| \leq 2^{\aleph_1}$. On the other hand, Theorem 2.2(ii) guarantees that $|\mathcal{P}(\mathcal{R})/I| > 2^{\aleph_0}$, and so $2^{\aleph_0} < 2^{\aleph_1}$. Now, if κ is the exact completeness of I it is easy to see that I projects to a κ -complete ideal on κ , and since I has a dense set of size of ω_1 it is clearly ω_2 -saturated. Thus, if $2^{\aleph_0} < \aleph_{\omega_1}$ we must have $\kappa = \omega_1$ and so we arrive at a situation where we have an ω_2 -saturated ω_1 -complete ideal on ω_1 with $2^{\aleph_0} < 2^{\aleph_1}$. A well-known result of Jech-Prikry [4, p. 3] now yields $2^{\aleph_0} = \aleph_1$ as desired.

REFERENCES

1. C. C. Chang and H. J. Keisler, *Model theory*, North-Holland, Amsterdam, 1973.
2. W. W. Comfort and A. W. Hager, *Cardinality of λ -complete Boolean algebras*, Pacific J. Math. **40** (1972), 541–545.
3. T. Frayne, A. C. Morel and D. S. Scott, *Reduced direct products*, Fund. Math. **51** (1962), 195–228.
4. T. Jech and K. Prikry, *Ideals over uncountable sets: applications of almost disjoint functions and generic ultrapowers*, Mem. Amer. Math. Soc. no. 214 (1979).
5. H. J. Keisler, *On cardinalities of ultrapowers*, Bull. Amer. Math. Soc. **70** (1964), 644–647.
6. J. D. Monk and P. R. Sparks, *Counting Boolean algebras*, Notices Amer. Math. Soc. **18** (1971), 551.
7. A. Pelc, *Ideals on the real line and Ulam's problem*, Fund. Math. **109** (1980).
8. K. Prikry, *Ideals and powers of cardinals*, Bull. Amer. Math. Soc. **81** (1975), 907–909.
9. A. D. Taylor, *On saturated sets of ideals and Ulam's problem*, Fund. Math. **109** (1980), 37–53.
10. —, *Regularity properties of ideals and ultrafilters*, Ann. Math Logic **16** (1979), 33–55.
11. H. Woodin, *Some consistency results in ZF using AD*, Cabal Seminar, Lecture Notes in Math., vol. 1019, Springer, pp. 172–199.

DEPARTMENT OF MATHEMATICS, UNION COLLEGE, SCHENECTADY, NEW YORK 12308