SHORTER NOTES

The purpose of this department is to publish very short papers of unusually polished character, for which there is no other outlet.

REMARK ON A GENERALIZED QUASI-VARIATIONAL INEQUALITY

WON KYU KIM

(Communicated by William J. Davis)

ABSTRACT. In this note, we prove a generalization of a generalized quasi-variational inequality of Shih and Tan in locally convex Hausdorff topological vector spaces.

In a recent paper [1], Shih and Tan proved some generalized quasi-variational inequalities in locally convex Hausdorff topological vector spaces (Theorems 1, 2 and 3 in [1]) or in normed spaces (Theorem 4 in [1]). In this note, we prove the following generalization of their Theorem 4 to locally convex Hausdorff topological vector spaces as follows:

THEOREM. *Let E be a locally convex Hausdorff topological vector space, and X be a nonempty compact convex subset of E . Let $S: X \rightarrow 2^X$ be continuous such that for each $x \in X$, $S(x)$ is a nonempty closed convex subset of X , and let $T: X \rightarrow 2^{E^*}$ be upper semicontinuous from the relative topology of X to the strong topology of E^* such that for each $x \in X$, $T(x)$ is a nonempty compact convex subset of E^* . Then there exists a point $y \in X$ such that*

- (1) $y \in S(y)$,
- (2) $\inf_{u \in T(y)} \operatorname{Re}(u, y - x) \leq 0$ for all $x \in S(y)$.

PROOF. By Theorem 3 of [1], it suffices to show that the set

$$\Sigma_2 := \left\{ y \in X \mid \sup_{x \in S(y)} \inf_{u \in T(y)} \operatorname{Re}(u, y - x) > 0 \right\}$$

is open in X . Let $y_0 \in \Sigma_2$. Then there exists $x_0 \in S(y_0)$ such that

$$t = \inf_{u \in T(y_0)} \operatorname{Re}(u, y_0 - x_0) > 0.$$

Received by the editors May 1, 1987 and, in revised form, November 20, 1987.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 49A29, 49A40; Secondary 52A07.

Key words and phrases. Locally convex Hausdorff topological vector space, generalized quasi-variational inequality, compact convex set, strong topology, weak*-topology.

Let $U_0 := \{p \in E^* \mid \operatorname{Re}(p, y_0 - x_0) > t/2\}$. Then U_0 is a weak*-open neighborhood of $T(y_0)$. Let $W := \{p \in E^* \mid \sup_{x, y \in X} |(p, x - y)| < t/6\}$. Then W is a strongly open neighborhood of $0 \in E^*$, so that $U_1 = T(y_0) + W$ is a strongly open neighborhood of $T(y_0)$. Now let $U := U_0 \cap U_1$; then U is a strongly open neighborhood of $T(y_0)$. Since T is upper semicontinuous at y_0 , there exists an open neighborhood V_1 of y_0 such that $T(y) \subset U$ for all $y \in V_1$. Furthermore, since $x \rightarrow \inf_{u \in T(y_0)} \operatorname{Re}(u, y_0 - x)$ and $y \rightarrow \inf_{u \in T(y_0)} \operatorname{Re}(u, y - y_0)$ are continuous, there exist an open neighborhood V of x_0 such that $\inf_{u \in T(y_0)} \operatorname{Re}(u, y_0 - x) > t/2$ for all $x \in V$, and an open neighborhood V_2 of y_0 such that $|\inf_{u \in T(y_0)} \operatorname{Re}(u, y - y_0)| < t/6$ for all $y \in V_2$. Since $x_0 \in S(y_0) \cap V$ and S is lower semicontinuous at y_0 , there exists an open neighborhood V_3 of y_0 such that $S(y) \cap V \neq \emptyset$ for all $y \in V_3$. Now let $\tilde{V} := (y_0 - x_0 + V) \cap (V_1 \cap V_2 \cap V_3)$; then $\tilde{V}$ is an open neighborhood of y_0 . For each $y \in \tilde{V}$, there exists $x \in V$ such that $y = y_0 - x_0 + x \in V_1 \cap V_2 \cap V_3$. Since $y \in V_1 \cap V_2 \cap V_3$, there exists $x_1 \in S(y) \cap V$, and $T(y) \subset U \subset T(y_0) + W$. It follows that

$$\begin{aligned} \inf_{u \in T(y)} \operatorname{Re}(u, y - x_1) &\geq \inf_{u \in T(y_0)} \operatorname{Re}(u, y - x_1) + \inf_{u \in W} \operatorname{Re}(u, y - x_1) \\ &\geq \inf_{u \in T(y_0)} \operatorname{Re}(u, y - x_1) - \frac{t}{6} \\ &\geq \inf_{u \in T(y_0)} \operatorname{Re}(u, y - y_0) + \inf_{u \in T(y_0)} \operatorname{Re}(u, y_0 - x_1) - \frac{t}{6} \\ &> -\frac{t}{6} + \frac{t}{2} - \frac{t}{6} = \frac{t}{6} > 0. \end{aligned}$$

Therefore we have $\sup_{x \in S(y)} \inf_{u \in T(y)} \operatorname{Re}(u, y - x) > 0$, so that $y \in \Sigma_2$. This shows that Σ_2 is open in X , and it completes the proof.

REFERENCES

1. M. H. Shih and K. K. Tan, *Generalized quasi-variational inequalities in locally convex topological vector spaces*, J. Math. Anal. Appl. **108** (1985), 333–343.

DEPARTMENT OF MATHEMATICS EDUCATION, CHUNGBUK NATIONAL UNIVERSITY,
CHEONGJU 360-763, KOREA