GENERALIZED OPEN MAPPING THEOREMS FOR BILINEAR MAPS, WITH AN APPLICATION TO OPERATOR ALGEBRAS

P. G. DIXON

(Communicated by John B. Conway)

ABSTRACT. Cohen [4] gave an example of a surjective bilinear mapping between Banach spaces which was not open, and Horowitz [8] gave a much simpler example. We build on Horowitz' example to produce a similar result for bilinear mappings such that every element of the target space is a linear combination of n elements of the range. An immediate application is that Bercovici's construction [1] of an operator algebra with property (A_1) but not $(A_1(r))$ can be extended to achieve property $(A_{1/n})$ without $(A_{1/n}(r))$.

Let $T: X \times Y \rightarrow Z$ be a continuous bilinear mapping between Banach spaces. We shall be concerned with the following two properties of such a mapping.

DEFINITION. For $n = 1, 2, 3, \dots$ we say T is $(1/n)$ -surjective if for every $z \in Z$ there exist $x_1, x_2, \dots, x_n \in X$ and $y_1, y_2, \dots, y_n \in Y$ with

$$(1) \quad z = T(x_1, y_1) + T(x_2, y_2) + \dots + T(x_n, y_n).$$

If, further, there exists a constant $K > 0$ such that for every z the $x_1, \dots, x_n$ and $y_1, \dots, y_n$ may be found, satisfying (1), with

$$\sum_{i=1}^n \|x_i\| \|y_i\| \leq K \|z\|,$$

then we say that T is $(1/n)$ -open.

REMARKS. (a) These properties are generalizations of the properties $(A_{1/n})$ and " $(A_{1/n}(r))$ for some $r > 0$ " of operator algebras studied by Bercovici, Foiaş and Pearcy [2]: hence the " $1/n$ " in our notation.

(b) If $n = 1$, " $(1/n)$ -surjective" is "surjective" and " $(1/n)$ -open" is "open".

(c) Notice that

$$\sum_{i=1}^n T(x_i, y_i) = \sum_{i=1}^n T(\lambda_i x_i, \lambda_i^{-1} y_i)$$

for any nonzero scalars $\lambda_1, \dots, \lambda_n$. We may therefore arrange that $\|x_i\| = \|y_i\|$ ($1 \leq i \leq n$). Thus, T is $(1/n)$ -open if and only if there is a constant $L > 0$ such that every $z \in Z$ with $\|z\| \leq 1$ may be expressed in the form (1) with $\|x_i\| \leq L$, $\|y_i\| \leq L$ ($1 \leq i \leq n$).

(d) We may work over the real field or the complex field and, in the latter case, we may study either bilinear or sesquilinear maps. The definitions above and Theorems 1 and 2 below apply to all three cases.

Received by the editors September 1, 1987.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 46A30; Secondary 47D35.

Key words and phrases. Open mapping theorem, bilinear map, dual algebra.

©1988 American Mathematical Society
0002-9939/88 \$1.00 + \$.25 per page

The best positive result known is the following, which is present, at least implicitly, in [9, Theorem 1.3, 7, Lemma 3.5, 3 and 5].

THEOREM 1. *Suppose $T: X \times Y \rightarrow Z$ is a continuous bilinear (or sesquilinear) map between separable Banach spaces X, Y, Z . Then*

(a) *if, for every $z \in Z$, there exists a positive integer n and $x_1, \dots, x_n \in X, y_1, \dots, y_n \in Y$ satisfying (1), then T is $(1/n)$ -surjective for some n (i.e. the dependence of n on z can be removed);*

(b) *if T is $(1/n)$ -surjective, then T is $(1/2n)$ -open.*

It is known [6] that the separability condition cannot be removed from (a). We conjecture that it cannot be removed from (b) either. Horowitz [8] showed that, even for finite-dimensional X, Y, Z , (b) cannot be improved to “surjective implies open” in the $n = 1$ case. Our main result here is the extension of his example to show that, generally, $(1/n)$ -surjective does not imply $(1/n)$ -open. It remains an interesting open question whether $(1/n)$ -surjective implies $(1/m)$ -open for $n < m < 2n$.

THEOREM 2. *For each positive integer n , there exists a $(1/n)$ -surjective sesquilinear (or bilinear) map $T: \mathbb{C}^{n+2} \times \mathbb{C}^{n+2} \rightarrow \mathbb{C}^N$ (where $N = n^2 + n + 2$), which is not $(1/n)$ -open.*

PROOF. We show the existence of a sesquilinear map T : a complex bilinear map $\tilde{T}$ with the same properties is then obtained by:

$$\tilde{T}(x, y) := T(x, \bar{y}) \quad (x, y \in \mathbb{C}^{n+2}),$$

and a real bilinear map is obtained just by replacing $\mathbb{C}$ by $\mathbb{R}$.

We observe that a sesquilinear map $T: X \times Y \rightarrow Z$ is $(1/n)$ -surjective (respectively, $(1/n)$ -open), if and only if the map $T^{(n)}: X^n \times Y^n \rightarrow Z$ is surjective (respectively, open), where

$$T^{(n)}((x_i)_{i=1}^n, (y_i)_{i=1}^n) := \sum_{i=1}^n T(x_i, y_i).$$

Our map $T: \mathbb{C}^{n+2} \times \mathbb{C}^{n+2} \rightarrow \mathbb{C}^N$ is defined by

$$\begin{aligned} T((u_1, \dots, u_n, v, w), (x_1, \dots, x_n, y, z)) \\ = ((u_i \bar{x}_j)_{i,j=1}^n, (u_i \bar{y})_{i=1}^n, u_1 \bar{z} + v \bar{x}_1 + w \bar{y}, w \bar{x}_1 + v \bar{y}). \end{aligned}$$

The reader may verify that in the case $n = 1$ this reduces to Horowitz' example, except for the complex conjugates and order of components. To construct $T^{(n)}$ we simply replace the numbers $u_1, \dots, z \in \mathbb{C}$ by vectors $\mathbf{u}_1, \dots, \mathbf{z} \in \mathbb{C}^n$ and the products $u_i \bar{x}_j$, etc., by scalar products $\mathbf{u}_i \cdot \mathbf{x}_j$, etc.:

$$\begin{aligned} T^{(n)}((\mathbf{u}_1, \dots, \mathbf{u}_n, \mathbf{v}, \mathbf{w}), (\mathbf{x}_1, \dots, \mathbf{x}_n, \mathbf{y}, \mathbf{z})) \\ = ((\mathbf{u}_i \cdot \mathbf{x}_j)_{i,j=1}^n, (\mathbf{u}_i \cdot \mathbf{y})_{i=1}^n, \mathbf{u}_1 \cdot \mathbf{z} + \mathbf{v} \cdot \mathbf{x}_1 + \mathbf{w} \cdot \mathbf{y}, \mathbf{w} \cdot \mathbf{x}_1 + \mathbf{v} \cdot \mathbf{y}). \end{aligned}$$

To show that $T^{(n)}$ is surjective, take any vector

$$\mathbf{h} = ((\xi_{ij})_{i,j=1}^n, (\eta_i)_{i=1}^n, \zeta, \theta) \in \mathbb{C}^N.$$

If $\eta_i = 0$ for all i , then let

$$\mathbf{x}_j = \mathbf{e}_j \quad (1 \leq j \leq n),$$

where $(\mathbf{e}_j)_i = 1$ if $i = j$, 0 if $i \neq j$,

$$\begin{aligned} \mathbf{y} &= \mathbf{0}, \\ \mathbf{z} &= \mathbf{0}, \\ \mathbf{u}_i &= (\xi_{i1}, \xi_{i2}, \dots, \xi_{in}) \quad (1 \leq i \leq n), \\ \mathbf{v} &= (\zeta, 0, 0, \dots, 0), \\ \mathbf{w} &= (\theta, 0, 0, \dots, 0); \end{aligned}$$

otherwise, let

$$\begin{aligned} \mathbf{u}_i &= \mathbf{e}_i \quad (1 \leq i \leq n) \\ \mathbf{x}_j &= (\bar{\xi}_{1j}, \bar{\xi}_{2j}, \dots, \bar{\xi}_{nj}) \quad (1 \leq j \leq n), \\ \mathbf{y} &= (\bar{\eta}_1, \bar{\eta}_2, \dots, \bar{\eta}_n), \\ \mathbf{v} &= \theta(\mathbf{y} \cdot \mathbf{y})^{-1} \mathbf{y}, \\ \mathbf{w} &= \mathbf{0}, \\ \mathbf{z} &= (\zeta - (\mathbf{v} \cdot \mathbf{x}_1))^{-1} \mathbf{u}_1. \end{aligned}$$

In both cases, $T^{(n)}((\mathbf{u}_1, \dots, \mathbf{w}), (\mathbf{x}_1, \dots, \mathbf{z})) = \mathbf{h}$. This proves that $T^{(n)}$ is surjective, so T is $(1/n)$ -surjective.

To show that T is not $(1/n)$ -open, suppose otherwise. Then, by Remark (c) and the equivalence of norms on finite-dimensional spaces, there exists a number $M \geq 1$ such that, whenever $\mathbf{h} = ((\xi_{ij}), (\eta_i), \zeta, \theta) \in \mathbb{C}^N$ with $|\xi_{ij}|, |\eta_i|, |\zeta|, |\theta| \leq 1$ for all i, j , there exist $\mathbf{u}_i, \mathbf{v}, \mathbf{w}, \mathbf{x}_i, \mathbf{y}, \mathbf{z} \in \mathbb{C}^n$ ($1 \leq i \leq n$) such that

$$(2) \quad T^{(n)}((\mathbf{u}_1, \dots, \mathbf{u}_n, \mathbf{v}, \mathbf{w}), (\mathbf{x}_1, \dots, \mathbf{x}_n, \mathbf{y}, \mathbf{z})) = \mathbf{h}$$

and

$$(3) \quad \|\mathbf{u}_i\|, \|\mathbf{v}\|, \|\mathbf{w}\|, \|\mathbf{x}_i\|, \|\mathbf{y}\|, \|\mathbf{z}\| \leq M \quad (1 \leq i \leq n),$$

where $\|\cdot\|$ denotes the Euclidean norm: $\|\mathbf{x}\| := (\mathbf{x} \cdot \mathbf{x})^{1/2}$ ($\mathbf{x} \in \mathbb{C}^n$).

Extending Horowitz' argument, we let $\mathbf{h}$ be defined by:

$$\begin{aligned} \xi_{11} &= a, & \xi_{ii} &= 1 \quad (2 \leq i \leq n), & \xi_{ij} &= 0 \quad (i \neq j), \\ \eta_1 &= a, & \eta_i &= 0 \quad (2 \leq i \leq n), \\ \zeta &= a, & \theta &= 1; \end{aligned}$$

where $0 < a < 1$ and we shall eventually make a tend to zero. Suppose $\mathbf{u}_i, \mathbf{v}, \mathbf{w}, \mathbf{x}_i, \mathbf{y}, \mathbf{z}$ satisfy (2) and (3) above. We wish to show that $\|\mathbf{x}_1\| \|\mathbf{u}_1\|$ must be small. This is most conveniently done by the following matrix argument.

Define the $n \times n$ matrices $\mathbf{A} = (\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n)$, $\mathbf{B} = (\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n)$ from the column vectors

$$\begin{aligned} \mathbf{a}_1 &= a^{-1/2} \mathbf{u}_1, & \mathbf{a}_i &= \mathbf{u}_i \quad (2 \leq i \leq n), \\ \mathbf{b}_1 &= a^{-1/2} \mathbf{x}_1, & \mathbf{b}_j &= \mathbf{x}_j \quad (2 \leq j \leq n). \end{aligned}$$

Then the equations $\mathbf{u}_i \cdot \mathbf{x}_j = \xi_{ij}$ ($1 \leq i, j \leq n$) become $\mathbf{A}^* \mathbf{B} = \mathbf{I}$. It follows that $\mathbf{A}$ and $\mathbf{B}$ are invertible and $\mathbf{B}\mathbf{A}^* = \mathbf{I} = \mathbf{B}^* \mathbf{A}$, so $\mathbf{B}^* \mathbf{B} \mathbf{A}^* \mathbf{A} = \mathbf{I}$. Hence

$$\begin{aligned} 1 &= (\mathbf{B}^* \mathbf{B} \mathbf{A}^* \mathbf{A})_{11} \\ &= \sum_{i=1}^n (\mathbf{b}_i \cdot \mathbf{b}_1)(\mathbf{a}_1 \cdot \mathbf{a}_i) \\ &= a^{-2} \|\mathbf{x}_1\|^2 \|\mathbf{u}_1\|^2 + \sum_{i=2}^n a^{-1} (\mathbf{x}_i \cdot \mathbf{x}_1)(\mathbf{u}_1 \cdot \mathbf{u}_i). \end{aligned}$$

Since $\|\mathbf{x}_i\|, \|\mathbf{u}_i\| \leq M$ ($1 \leq i \leq n$), this yields

$$a^{-2} \|\mathbf{x}_1\|^2 \|\mathbf{u}_1\|^2 \leq 1 + (n-1)a^{-1} \|\mathbf{x}_1\| \|\mathbf{u}_1\| M^2.$$

Hence

$$(4) \quad \|\mathbf{x}_1\| \|\mathbf{u}_1\| \leq a(1 + (n-1)M^2) \leq anM^2.$$

Next, since $\mathbf{A}$ is invertible, $(\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n)$ is a basis for $\mathbb{C}^n$. Now $\mathbf{u}_i \cdot (\mathbf{x}_1 - \mathbf{y}) = 0$ for all i , so it follows that $\mathbf{x}_1 = \mathbf{y}$. Then the equation $\mathbf{w} \cdot \mathbf{x}_1 + \mathbf{v} \cdot \mathbf{y} = \theta = 1$ gives

$$(5) \quad (\mathbf{w} + \mathbf{v}) \cdot \mathbf{x}_1 = 1,$$

and so

$$(6) \quad \|\mathbf{w} + \mathbf{v}\| \|\mathbf{x}_1\| \geq 1.$$

We also have

$$\mathbf{u}_1 \cdot \mathbf{z} + \mathbf{v} \cdot \mathbf{x}_1 + \mathbf{w} \cdot \mathbf{y} = \zeta = a,$$

so

$$\mathbf{u}_1 \cdot \mathbf{z} = a - (\mathbf{v} + \mathbf{w}) \cdot \mathbf{x}_1 = -(1-a),$$

by (5), and so

$$(7) \quad \|\mathbf{u}_1\| \geq (1-a)/M.$$

Combining (4) and (7),

$$(8) \quad \|\mathbf{x}_1\| \leq anM^3/(1-a).$$

Finally,

$$\begin{aligned} 2M &\geq \|\mathbf{w} + \mathbf{v}\| \\ &\geq \|\mathbf{x}_1\|^{-1}, \quad \text{by (6),} \\ &\geq (1-a)/anM^3, \quad \text{by (8).} \end{aligned}$$

Now M is a constant, independent of a , so letting a tend to zero gives the required contradiction, completing the proof of Theorem 2.

COROLLARY. *For every positive integer n , there is a commutative $(n^2 + n + 3)$ -dimensional algebra of operators on the Hilbert space $\mathbb{C}^{2n+5}$ which has property $(\mathbf{A}_{1/n})$, but which, for all $r > 0$, does not have property $(\mathbf{A}_{1/n}(r))$.*

The proof of this is the same as that of [1], where Bercovici derives this result for $n = 1$ from Horowitz' example. We therefore omit further discussion and refer the reader to [1] for an explanation of the terminology as well as for the proof.

REFERENCES

1. H. Bercovici, *Note on property (A_1)*, Linear Algebra Appl. **91** (1987), 213–216.
2. H. Bercovici, C. Foias and C. Pearcy, *Dual algebras with applications to invariant subspaces and dilation theory*, CBMS Regional Conf. Ser. in Math., no. 56, Amer. Math. Soc., Providence, R. I., 1985.
3. B. Chevreau and J. Esterle, *Pettis' lemma and topological properties of dual algebras*, Michigan Math. J. **34** (1987), 143–146.
4. P. J. Cohen, *A counterexample to the closed graph theorem for bilinear maps*, J. Funct. Anal. **16** (1974), 235–239.
5. H. G. Dales, *Automatic continuity theory* (in preparation).
6. P. G. Dixon, *Non-separable Banach algebras whose squares are pathological*, J. Funct. Anal. **26** (1977), 190–200.
7. ———, *Automatic continuity of positive functionals on topological involution algebras*, Bull. Austral. Math. Soc. **23** (1981), 265–281.
8. C. Horowitz, *An elementary counterexample to the open mapping principle for bilinear maps*, Proc. Amer. Math. Soc. **53** (1975), 293–294.
9. R. J. Loy, *Multilinear mappings and Banach algebras*, J. London Math. Soc. (2) **14** (1976), 423–429.

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF SHEFFIELD, SHEFFIELD,
S3 7RH, ENGLAND