A NONLINEAR ERGODIC THEOREM FOR A REVERSIBLE SEMIGROUP OF LIPSCHITZIAN MAPPINGS IN A HILBERT SPACE

HAJIME ISHIHARA AND WATARU TAKAHASHI

(Communicated by John B. Conway)

ABSTRACT. Let C be a nonempty closed convex subset of a Hilbert space, S a right reversible semitopological semigroup, $\mathcal{S} = \{T_t : t \in S\}$ a continuous representation of S as Lipschitzian mappings on a closed convex subset C into C , and $F(\mathcal{S})$ the set of common fixed points of mappings T_t , $t \in S$. Then we deal with the existence of a nonexpansive retraction P of C onto $F(\mathcal{S})$ such that $PT_t = T_tP = P$ for each $t \in S$ and Px is contained in the closure of the convex hull of $\{T_tx : t \in S\}$ for each $x \in C$.

1. Introduction. Let C be a closed convex subset of a real Hilbert space H and T be a mapping of C into itself. T is said to be a Lipschitzian mapping if for each $n \geq 1$ there exists a positive real number k_n such that

$$\|T^n x - T^n y\| \leq k_n \|x - y\|$$

for all $x, y \in C$. A Lipschitzian mapping is said to be nonexpansive if $k_n = 1$ for all $n \geq 1$ and asymptotically nonexpansive if $\lim_n k_n = 1$, respectively. We denote by $F(T)$ the set of fixed points of T . The first nonlinear ergodic theorem for nonexpansive mappings was established in 1975 by Baillon [1]: Let C be a closed convex subset of a Hilbert space and T be a nonexpansive mapping of C into itself. If the set $F(T)$ is nonempty, then for each $x \in C$, the Cesàro means

$$S_n(x) = \frac{1}{n} \sum_{k=0}^{n-1} T^k x$$

converge weakly to some $y \in F(T)$. In this case, putting $y = Px$ for each $x \in C$, P is a nonexpansive retraction of C onto $F(T)$ such that $PT = TP = P$ and $Px \in \overline{\text{co}}\{T^n x : n = 0, 1, 2, \dots\}$ for each $x \in C$, where $\overline{\text{co}}A$ is the closure of the convex hull of A . In [11 and 12], Takahashi proved the existence of such a retraction—"ergodic retraction"—for an amenable and a right reversible semigroup of nonexpansive mappings in a Hilbert space. On the other hand, Hirano and Takahashi [4] proved the Cesàro means for asymptotically nonexpansive mappings converge weakly to a fixed point.

In this paper, we deal with the existence of "ergodic retraction" for a right reversible semigroup of Lipschitzian mappings; that is, we prove a nonlinear ergodic theorem for such a semigroup in a Hilbert space. This theorem is a generalization

Received by the editors June 20, 1987.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 47A35; Secondary 47H09.

Key words and phrases. Ergodic theorem, reversible semigroup, asymptotically nonexpansive mapping, fixed point.

of the result [4, 12]. Furthermore, we prove a weak convergence theorem which is similar to that of [7, 9].

2. Nonlinear ergodic theorem. Let S be a semitopological semigroup, i.e., S is a semigroup with a Hausdorff topology such that for each $a \in S$ the mappings $s \rightarrow a \cdot s$ and $s \rightarrow s \cdot a$ from S to S are continuous. Let C be a nonempty closed convex subset of a Hilbert space H . Then a family $\mathcal{S} = \{T_t: t \in S\}$ of mappings from C into itself is said to be a Lipschitzian semigroup on C if $\mathcal{S}$ satisfies the following:

- (1) $T_{ts}(x) = T_t T_s(x)$ for $t, s \in S$ and $x \in C$;
- (2) the mapping $(s, x) \rightarrow T_s(x)$ from $S \times C$ into C is continuous when $S \times C$ has the product topology;
- (3) for each $s \in S$, there exists $k_s > 0$ such that $\|T_s(x) - T_s(y)\| \leq k_s \|x - y\|$ for $x, y \in C$.

A semitopological semigroup S is right reversible if any two closed left ideals of S have nonvoid intersection. In this case, $(S, \leq)$ is a directed system when the binary relation " $\leq$ " on S is defined by $a \leq b$ if and only if $\{a\} \cup \overline{Sa} \supseteq \{b\} \cup \overline{Sb}$. Let $F(\mathcal{S})$ denote the set $\{x \in C: T_s x = x \text{ for all } s \in S\}$ of common fixed points of mappings T_s , $s \in S$ in C (fixed point theorems for Lipschitzian semigroups were proved in [6, 3]). Then we have the following:

THEOREM 1. *Let C be a closed convex subset of a real Hilbert space H and let S be a right reversible semitopological semigroup. Let $\mathcal{S} = \{T_t: t \in S\}$ be a Lipschitzian semigroup on C with $\limsup_s k_s \leq 1$. Then $F(\mathcal{S})$ is a closed convex subset of C .*

PROOF. Closedness of $F(\mathcal{S})$ is obvious. To show convexity it is sufficient to prove that $z = (x + y)/2 \in F(\mathcal{S})$ for all $x, y \in F(\mathcal{S})$. We have

$$\|T_t z - x\|^2 = \|T_t z - T_t x\|^2 \leq k_t^2 \|z - x\|^2 = \frac{1}{4} k_t^2 \|x - y\|^2$$

and

$$\|T_t z - y\|^2 = \|T_t z - T_t y\|^2 \leq k_t^2 \|z - y\|^2 = \frac{1}{4} k_t^2 \|x - y\|^2.$$

Thus

$$\|T_t z - z\|^2 = \frac{1}{2} \|T_t z - x\|^2 + \frac{1}{2} \|T_t z - y\|^2 - \frac{1}{4} \|x - y\|^2 \leq \frac{1}{4} (k_t^2 - 1) \|x - y\|^2$$

and hence $\lim_t \|T_t z - z\| = 0$. Therefore we obtain

$$z = \lim_t T_t z = \lim_t T_{st} z = \lim_t T_s T_t z = T_s \lim_t T_t z = T_s z$$

for all $s \in S$.

Let $\{x_\alpha: \alpha \in A\}$ be a bounded net of a Hilbert space H and let C be a closed convex subset of H . Then we define

$$r(x) = \limsup_{\beta \leq \alpha} \|x_\alpha - x\| \quad \text{and} \quad r = \inf\{r(x): x \in C\}.$$

It is well known that there exists a unique element $a \in C$ with $r(a) = r$, called the asymptotic center of $\{x_\alpha: \alpha \in A\}$ in C . A useful lemma is a result proved in [5], which we state here as:

LEMMA 1. $r^2 + \|a - x\|^2 \leq r(x)^2$ for every $x \in C$.

As a consequence, we have the following.

LEMMA 2 (LIM [8]). Let $\{y_\beta\}$ be a net of C such that $\limsup_\beta r(y_\beta) \leq r$. Then $y_\beta \rightarrow a$.

PROOF. By Lemma 1, we have $r^2 + \|a - y_\beta\|^2 \leq r(y_\beta)^2$ and hence

$$r^2 + \limsup_\beta \|a - y_\beta\|^2 \leq \limsup_\beta r(y_\beta)^2 \leq r^2.$$

Therefore we have $\limsup_\beta \|a - y_\beta\|^2 = 0$ and this implies $y_\beta \rightarrow a$.

We also know that if $\{x_\alpha\} \subseteq C$ and if $\{x_\alpha\}$ converges weakly to $y \in C$ then $y = a$ [2, Theorem 4.2]. Let Q be the metric projection of H onto $F(\mathcal{S})$. Then, by Phelps [10], Q is nonexpansive. Before proving a nonlinear ergodic theorem, we prove the following crucial lemma.

LEMMA 3. Let C be a closed convex subset of a real Hilbert space H and let S be a right reversible semitopological semigroup. Let $\mathcal{S} = \{T_t: t \in S\}$ be a Lipschitzian semigroup on C with $\limsup_s k_s \leq 1$. Suppose that $F(\mathcal{S}) \neq \emptyset$. Then for each $x \in C$, $\{QT_sx: s \in S\}$ converges to the asymptotic center of $\{T_sx: s \in S\}$ in $F(\mathcal{S})$.

PROOF. Let z be the asymptotic center of $\{T_sx: s \in S\}$ in $F(\mathcal{S})$. Then, for all $s, t \in S$ we obtain

$$\begin{aligned} r(QT_sx) &\leq \sup_{a \geq ts} \|T_ax - QT_sx\| = \sup_{a \geq t} \|T_ax - QT_sx\| \\ &= \sup_{a \geq t} \|T_aT_sx - T_aQT_sx\| \leq \left(\sup_{a \geq t} k_a \right) \|T_sx - QT_sx\| \\ &\leq \left(\sup_{a \geq t} k_a \right) \|T_sx - z\| \end{aligned}$$

and hence

$$r(QT_sx) \leq \limsup_t k_t \|T_sx - z\| \leq \|T_sx - z\|$$

for all $s \in S$. Therefore we have $\limsup_s r(QT_sx) \leq \limsup_s \|T_sx - z\| \leq r$. By Lemma 2, we obtain $QT_sx \rightarrow z$.

THEOREM 2. Let C be a closed convex subset of a Hilbert space H and let S be a right reversible semitopological semigroup. Let $\mathcal{S} = \{T_t: t \in S\}$ be a Lipschitzian semigroup on C with $\limsup_s k_s \leq 1$. Suppose that

$$F(\mathcal{S}) = \bigcap \{F(T_s): s \in S\} \neq \emptyset.$$

Then the following are equivalent:

- (a) $\bigcap_{s \in S} \overline{\text{co}}\{T_tx: t \geq s\} \cap F(\mathcal{S}) \neq \emptyset$ for each $x \in C$.
- (b) There is a nonexpansive retraction P of C onto $F(\mathcal{S})$ such that $PT_t = T_tP = P$ for every $t \in S$ and $Px \in \overline{\text{co}}\{T_tx: t \in S\}$ for every $x \in C$.

PROOF. (b) $\Rightarrow$ (a). Let $x \in C$. Then $Px \in F(\mathcal{S})$. Also

$$Px \in \bigcap_{s \in S} \overline{\text{co}}\{T_tx: t \geq s\}.$$

In fact,

$$Px = PT_sx \in \overline{\text{co}}\{T_tT_sx: t \in S\} \subset \overline{\text{co}}\{T_tx: t \geq s\}$$

for every $s \in S$.

(a) $\Rightarrow$ (b). Let $x \in C$ and $f \in F(\mathcal{S})$. Then for each $s, t \in S$, we have

$$\begin{aligned} \limsup_a \|T_a x - f\| &\leq \sup_{a \geq ts} \|T_a x - f\| = \sup_{a \geq t} \|T_{as} x - f\| \\ &= \sup_{a \geq t} \|T_a T_s x - T_a f\| \leq \left(\sup_{a \geq t} k_a \right) \|T_s x - f\| \end{aligned}$$

and hence

$$\limsup_a \|T_a x - f\| \leq \left(\limsup_t k_t \right) \|T_s x - f\| \leq \|T_s x - f\|$$

for every $s \in S$. So $\limsup_s \|T_s x - f\| \leq \liminf_s \|T_s x - f\|$ and hence the $\lim_s \|T_s x - f\|$ exists. Let Q be the metric projection of H onto $F(\mathcal{S})$. Then by Lemma 3, $\{QT_s x\}$ converges to the asymptotic center z of $\{T_s x: s \in S\}$ in $F(\mathcal{S})$. Let $u \in \bigcap_{s \in S} \overline{\text{co}}\{T_t x: t \geq s\} \cap F(\mathcal{S})$. Then, since

$$\|z - u\|^2 = \|T_s x - u\|^2 - \|T_s x - z\|^2 - 2\langle z - u, T_s x - z \rangle$$

for every $s \in S$, we have

$$\begin{aligned} \|z - u\|^2 + 2 \lim_s \langle z - u, T_s x - z \rangle &= \lim_s \|T_s x - u\|^2 - \lim_s \|T_s x - z\|^2 \\ &= \limsup_s \|T_s x - u\|^2 - \limsup_s \|T_s x - z\|^2 \\ &= r(u)^2 - r^2 \geq 0. \end{aligned}$$

Let $\varepsilon > 0$. Then we have

$$2 \lim_s \langle z - u, T_s x - z \rangle > -\|z - u\|^2 - \varepsilon.$$

Hence there exists $s_0 \in S$ such that

$$2\langle z - u, T_s x - z \rangle > -\|z - u\|^2 - \varepsilon$$

for every $s \geq s_0$. Since $u \in \overline{\text{co}}\{T_t x: t \geq s_0\}$, we have

$$2\langle z - u, u - z \rangle \geq -\|z - u\|^2 - \varepsilon.$$

This inequality implies $\|z - u\|^2 \leq \varepsilon$. Since ε is arbitrary, we have $z = u$. Therefore

$$\bigcap_{s \in S} \overline{\text{co}}\{T_t x: t \geq s\} \cap F(\mathcal{S}) = \{z\}.$$

Set $Px = \lim_t QT_t x$ for every $x \in C$. Then we have $T_s Px = Px$ and

$$PT_s x = \lim_t QT_t T_s x = \lim_t QT_{ts} x = Px$$

for every $s \in S$ and $x \in C$. From $\{Px\} = \bigcap_{s \in S} \overline{\text{co}}\{T_t x: t \geq s\} \cap F(\mathcal{S})$, it is obvious that $Px \in \overline{\text{co}}\{T_s x: s \in S\}$ for each $x \in C$. Since

$$\begin{aligned} \|Px - Py\| &= \lim_t \|QT_t x - QT_t y\| \leq \limsup_t \|T_t x - T_t y\| \\ &\leq \left(\limsup_t k_t \right) \|x - y\| \leq \|x - y\| \end{aligned}$$

for every $x, y \in C$, it follows that P is nonexpansive.

We now turn to consider the weak convergence of $\{T_s x: s \in S\}$ and obtain the results similar to these of Lau [7] and Passty [9]. We denote by $\omega(x)$ the set of all weak limit points of subnets of the net $\{T_s x: s \in S\}$. Note that if $F(\mathcal{S})$ is nonempty then $\{T_s x: s \in S\}$ is bounded and hence $\omega(x)$ is nonempty. We start with proving the following lemma.

LEMMA 4. Let C be a closed convex subset of a Hilbert space H and let S be a right reversible semitopological semigroup. Let $\mathcal{S} = \{T_t: t \in S\}$ be a Lipschitzian semigroup on C with $\limsup_s k_s \leq 1$. Suppose that $F(\mathcal{S}) \neq \emptyset$. Let $x \in C$. If $\omega(x) \subseteq F(\mathcal{S})$, then the net $\{T_s x: s \in S\}$ converges weakly to some $y \in F(\mathcal{S})$.

PROOF. By Lemma 3, the net $\{QT_s x: s \in S\}$ converges strongly to some $y \in F(\mathcal{S})$. Since $\omega(x) \subseteq \bigcap_{s \in S} \overline{\text{co}}\{T_t x: t \geq s\}$, we have

$$\bigcap_{s \in S} \overline{\text{co}}\{T_t x: t \geq s\} \cap F(\mathcal{S}) \supseteq \omega(x) \neq \emptyset.$$

Hence, as in the proof of Theorem 2, we obtain

$$\{y\} = \bigcap_{s \in S} \overline{\text{co}}\{T_t x: t \geq s\} \cap F(\mathcal{S}).$$

Therefore we have $\{y\} = \omega(x)$.

Let C be a closed convex subset of a Hilbert space H and let $\mathcal{S} = \{T_t: t \in S\}$ be a Lipschitzian semigroup on C . A subset G of S is called a uniformly generating set of $\mathcal{S}$ if for each $s \in S$ and $\varepsilon > 0$, there exist $g_1, g_2, \dots, g_m \in G$ such that

$$\|T_s x - T_{g_1 g_2 \dots g_m} x\| < \varepsilon$$

for every $x \in C$.

THEOREM 3. Let C be a closed convex subset of a Hilbert space H and let S be a right reversible semitopological semigroup. Let $\mathcal{S} = \{T_t: t \in S\}$ be a Lipschitzian semigroup on C with $\limsup_s k_s \leq 1$. Suppose that $F(\mathcal{S}) \neq \emptyset$. Let $x \in C$. If $\lim_s \|T_{gs} x - T_s x\| = 0$ for all g in a uniformly generating set G of $\mathcal{S}$, then the net $\{T_s x: s \in S\}$ converges weakly to some $y \in F(\mathcal{S})$.

PROOF. By Lemma 4, it suffices to show that $\omega(x) \subseteq F(\mathcal{S})$. Let $\{T_{s_\alpha} x\}$ be a subnet of $\{T_s x: s \in S\}$ converging weakly to some $y \in C$. Let $\varepsilon > 0$. Then for each $t \in S$, there exist $g_1, g_2, \dots, g_m \in G$ such that

$$\|T_t T_{s_\alpha} x - T_{g_1 g_2 \dots g_m} T_{s_\alpha} x\| < \varepsilon$$

for every α . Hence we have

$$\begin{aligned} \|T_{s_\alpha} x - T_t y\| &\leq \|T_{s_\alpha} x - T_{g_1 g_2 \dots g_m s_\alpha} x\| + \|T_{g_1 g_2 \dots g_m s_\alpha} x - T_{t s_\alpha} x\| + \|T_{t s_\alpha} x - T_t y\| \\ &\leq \|T_{s_\alpha} x - T_{g_1 s_\alpha} x\| + \varepsilon + k_t \|T_{s_\alpha} x - y\| \\ &\leq \|T_{s_\alpha} x - T_{g_1 s_\alpha} x\| + \|T_{g_1 s_\alpha} x - T_{g_1 g_2 \dots g_m s_\alpha} x\| + \varepsilon + k_t \|T_{s_\alpha} x - y\| \\ &\leq \|T_{s_\alpha} x - T_{g_1 s_\alpha} x\| + k_{g_1} \|T_{s_\alpha} x - T_{g_2 \dots g_m s_\alpha} x\| + \varepsilon + k_t \|T_{s_\alpha} x - y\| \\ &\leq \|T_{s_\alpha} x - T_{g_1 s_\alpha} x\| + k_{g_1} \|T_{s_\alpha} x - T_{g_2 s_\alpha} x\| + k_{g_1} k_{g_2} \|T_{s_\alpha} x - T_{g_3 s_\alpha} x\| \\ &\quad + \dots + k_{g_1} k_{g_2} \dots k_{g_{m-1}} \|T_{s_\alpha} x - T_{g_m s_\alpha} x\| + \varepsilon + k_t \|T_{s_\alpha} x - y\| \end{aligned}$$

and

$$\limsup_\alpha \|T_{s_\alpha} x - T_t y\| \leq \varepsilon + k_t \limsup_\alpha \|T_{s_\alpha} x - y\|.$$

Since ε is arbitrary, we have

$$\limsup_\alpha \|T_{s_\alpha} x - T_t y\| \leq k_t \limsup_\alpha \|T_{s_\alpha} x - y\|.$$

On the other hand, since y is the asymptotic center of $\{T_{s_\alpha}x\}$ in C we obtain

$$\limsup_t r(T_t y) \leq \left(\limsup_t k_t \right) r(y) \leq r$$

and hence $T_t y \rightarrow y$. Therefore we have $y \in F(\mathcal{S})$.

REMARK. Let γ be a positive real number and let $\mathcal{S} = \{T_t : t \in S\}$ be a Lipschitzian semigroup with $\limsup_s k_s \leq \gamma$. Then, putting $k'_s = \sup_{t \geq s} k_t$, we have

$$\|T_s x - T_s y\| \leq k_s \|x - y\| \leq \sup_{t \geq s} k_t \|x - y\| = k'_s \|x - y\|$$

and $\lim_s k'_s = \limsup_s k_s$. Hence $\mathcal{S}$ is a Lipschitzian semigroup with $\lim_s k'_s \leq \gamma$.

REFERENCES

1. J. B. Baillon, *Un théorème de type ergodique pour les contractions non linéaires dans un espace de Hilbert*, C. R. Acad. Sci. Paris. Sér. A-B **280** (1975), 1511–1514.
2. K. Goebel and S. Reich, *Uniform convexity, hyperbolic geometry, and nonexpansive mappings*, Marcel Dekker, New York, 1984.
3. K. Goebel, W. A. Kirk and R. L. Thele, *Uniformly lipschitzian families of transformations in Banach space*, Canad. J. Math. **26** (1974), 1245–1256.
4. N. Hirano and W. Takahashi, *Nonlinear ergodic theorems for nonexpansive mappings in Hilbert space*, Kodai Math. J. **2** (1979), 11–25.
5. H. Ishihara and W. Takahashi, *Fixed point theorems for uniformly lipschitzian semigroups in Hilbert spaces*, J. Math. Anal. Appl. **127** (1987), 206–210.
6. H. Ishihara, *Fixed point theorems for lipschitzian semigroups* (to appear).
7. A. T. Lau, *Semigroup of nonexpansive mappings on a Hilbert space*, J. Math. Anal. Appl. **105** (1985), 514–522.
8. T. C. Lim, *On asymptotic centers and fixed points of nonexpansive mappings*, Canad. J. Math. **32** (1980), 421–430.
9. G. B. Passty, *Construction of fixed points for asymptotically nonexpansive mappings*, Proc. Amer. Math. Soc. **84** (1982), 212–216.
10. R. R. Phelps, *Convex sets and nearest points*, Proc. Amer. Math. Soc. **8** (1957), 790–797.
11. W. Takahashi, *A nonlinear ergodic theorem for an amenable semigroup of nonexpansive mappings in a Hilbert space*, Proc. Amer. Math. Soc. **81** (1981), 253–256.
12. —, *A nonlinear ergodic theorem for a reversible semigroup of nonexpansive mappings in a Hilbert space*, Proc. Amer. Math. Soc. **97** (1986), 55–58.

DEPARTMENT OF INFORMATION SCIENCE, TOKYO INSTITUTE OF TECHNOLOGY, OH-KAYAMA, MEGURO-KU, TOKYO 152, JAPAN