ON STABILITY OF ENDOMORPHISMS

HIROSHI IKEDA

(Communicated by George R. Sell)

ABSTRACT. In this note we prove a generalization of R. Mañé's theorem. R. Mañé proved that C^r absolutely stable endomorphisms satisfy Axiom A. We prove that if an endomorphism f is both C^r structurally and infinitesimally stable, then f satisfies Axiom A.

1. Introduction. This note is a generalization of R. Mañé's theorem on absolute stability [4]. We prove this theorem under weaker hypotheses. Our Theorem is the following

THEOREM. *If $f \in \text{End}^r(M)$ is both C^r structurally and infinitesimally stable, then f satisfies Axiom A.*

First we establish some background. Let M be a compact connected smooth manifold without boundary and let $\text{End}^r(M)$, $r \geq 1$, be the space of C^r endomorphisms of M endowed with the C^r topology. We say that f is C^r structurally stable if there exists a neighborhood $\mathcal{U}$ of f in $\text{End}^r(M)$ such that for every $g \in \mathcal{U}$ there exists a homeomorphism h of M satisfying $fh = hg$. If $f \in \text{End}^r(M)$, let $\text{Per}(f)$ be the set of all periodic points of f and let $\Omega(f) = \{x \in M \mid \text{for every neighborhood } U \text{ of } x, \text{ there exists } n > 0 \text{ with } f^n(U) \cap U \neq \emptyset\}$. We denote by $S(f)$ the set of singularities of f , i.e. those points x of M where $Tf|_{T_x M}$ is not injective.

We give two definitions following R. Mañé [4].

DEFINITION. We say that $f \in \text{End}^r(M)$ is C^r absolutely stable if there exist a neighborhood $\mathcal{U}$ of f in $\text{End}^r(M)$ and a constant $K > 0$ such that for all $g \in \mathcal{U}$ there exists a homeomorphism h of M satisfying $gh = hf$ and $d(h, I) \leq Kd(f, g)$ where $d(\cdot, \cdot)$ is defined by $d(f_1, f_2) = \sup\{\rho(f_1(x), f_2(x)) \mid x \in M\}$, $\rho(\cdot, \cdot)$ being a metric on M and I is the identity map of M .

DEFINITION. We say that $f \in \text{End}^r(M)$ satisfies Axiom A if there exist a continuous splitting $TM|_{\Omega(f)} = E^s \oplus E^u$, and a Riemannian metric $|\cdot|$ on M , and constants $K > 0$, $0 < \lambda < 1$ satisfying:

- (a) $(Tf)E^s \subset E^s$, $(Tf)E^u = E^u$;
- (b) $|(Tf)^n|_{E_x^s}| \leq K\lambda^n$ for $x \in \Omega(f)$, $n > 0$,
 $|(Tf)^n v| \geq K\lambda^{-n}|v|$ for $x \in \Omega(f)$, $v \in E_x^u$, $n > 0$;
- (c) if $x_1 \neq x_2 \in \Omega(f)$ and $f(x_1) = f(x_2) = x$, then $E_x^s = \{0\}$;
- (d) $\text{Per}(f)$ is dense in $\Omega(f)$;
- (e) $S(f) \cap \Omega(f) = \emptyset$.

In [8] the following conjecture was stated.

Received by the editors September 29, 1986.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 58F10; Secondary 58F15.

Key words and phrases. Infinitesimal stability, Axiom A, prehyperbolic set, $\bar{P}$ -stability.

CONJECTURE. If $f \in \text{End}^r(M)$ is C^r structurally stable, then $\Omega(f) \cap S(f) = \emptyset$.

In fact, $\text{Per}(f) \cap S(f) = \emptyset$ holds if f is C^r structurally stable, and $\Omega(f) \cap S(f) = \emptyset$ holds if f is C^r absolutely stable. The latter property is a part of the theorem by R. Mañé [4].

THEOREM M. C^r absolutely stable endomorphisms satisfy Axiom A.

On the other hand, F. Przytycki defined Axiom A somewhat different from Mañé's Axiom A, and found sufficient conditions for Ω -stability of an endomorphism [6, 7].

Our Theorem is a generalization of Theorem M.

2. Proof of the Theorem. In this section we will prove our Theorem. To prove the Theorem we need some definitions and lemmas. If $\Lambda \subset M$ is a compact subset of M let $\Gamma^b(\Lambda)$ be the space of bounded sections of $TM|_\Lambda$ with the norm $\|\eta\| = \sup\{\|\eta(x)\| \mid x \in \Lambda\}$ and let $\Gamma^0(\Lambda)$ be the closed subspace of continuous sections. If $f \in \text{End}^r(M)$ and $f(\Lambda) \subset \Lambda$ let $T_f M|_\Lambda$ be the vector bundle on Λ consisting of couples (p, v) with $p \in \Lambda$, $v \in T_{f(p)}M$. Let $\Gamma_f^b(\Lambda)$, $\Gamma_f^0(\Lambda)$ be the corresponding spaces of bounded and continuous sections of $T_f M|_\Lambda$.

We define the linear operator $L_f: \Gamma^b(\Lambda) \rightarrow \Gamma_f^b(\Lambda)$ by

$$L_f(\eta) = (Tf) \circ \eta - \eta \circ f \quad \text{for } \eta \in \Gamma^b(\Lambda).$$

DEFINITION. Let $f \in \text{End}^r(M)$ and let $\Lambda \subset M$ be a compact subset with $f(\Lambda) = \Lambda$. We say that Λ is a *prehyperbolic set* for f if there exist a continuous splitting $TM|_\Lambda = E^s \oplus E^u$, and a Riemannian metric $|\cdot|$ on M and constants $K > 0$, $0 < \lambda < 1$ satisfying:

$$(a) \quad (Tf)E^s \subset E^s, (Tf)E^u = E^u;$$

$$(b) \quad \begin{aligned} |(Tf)^n|E_x^s| &\leq K\lambda^n \text{ for } x \in \Lambda, n > 0, \\ |(Tf)^n v| &\geq K\lambda^{-n}|v| \text{ for } x \in \Lambda, v \in E_x^u, n > 0; \end{aligned}$$

$$(c) \text{ if } x_1 \neq x_2 \in \Lambda \text{ and } f(x_1) = f(x_2) = x, \text{ then } E_x^s = \{0\}.$$

We say that $f \in \text{End}^r(M)$ is *infinitesimally stable* if the linear operator $L_f: \Gamma^0(M) \rightarrow \Gamma_f^0(M)$ is surjective. It is evident that the absolute stability of f implies the structural stability of f . Also it is easy to prove that if $f \in \text{End}^r(M)$ is absolutely stable, then f is infinitesimally stable by the similar argument for the case of diffeomorphisms in [1]. If $x \in M$ let $\omega(x) = \omega(x, f)$ be the set of ω -limit points of x for f and let $\bar{L}^+(f)$ be the closure of $L^+(f) = \{\omega(x) \mid x \in M\}$.

In the proof of the theorem, we shall use the following two lemmas of [4].

LEMMA 1. If $f \in \text{End}^r(M)$ is infinitesimally stable, then $\bar{L}^+(f)$ is prehyperbolic.

LEMMA 2. Suppose that $f \in \text{End}^r(M)$ and $\bar{L}^+(f)$ is prehyperbolic. Then given $x \in \bar{L}^+(f)$ and a neighborhood $\mathcal{U}$ of f there exist a neighborhood U of x and $g \in \mathcal{U}$ such that:

- (a) $g(y) = f(y)$ for all $y \in U$;
- (b) $x \in \text{Per}(g)$.

From the argument of R. Mañé [4], it clearly follows that we only need to prove the following proposition in order to prove the Theorem.

PROPOSITION. *If $f \in \text{End}^r(M)$ is both C^r structurally and infinitesimally stable, then $\overline{L}^+(f) = \overline{\text{Per}(f)}$.*

To prove the Proposition, we introduce the concept of $\overline{P}$ -stability. $\overline{P} = \overline{\text{Per}(f)}$ will denote the closure of the set of periodic points of f . We say that $f \in \text{End}^r(M)$ is $\overline{P}$ -stable if there exists a neighborhood $\mathcal{U}$ of f in $\text{End}^r(M)$ such that for each $g \in \mathcal{U}$ there exists a homeomorphism $h: \overline{\text{Per}(f)} \rightarrow \overline{\text{Per}(g)}$ satisfying $hf = gh$ [5]. It is obvious that the structural stability implies the $\overline{P}$ -stability.

PROOF OF PROPOSITION. We first suppose that $\overline{\text{Per}(f)} \subsetneq \overline{L}^+(f)$. We show that $\overline{L}^+(f) \cap S(f) = \emptyset$. Suppose that $x \in \overline{L}^+(f) \cap S(f)$. By Lemmas 1 and 2, we can take $g \in \text{End}^r(M)$ nearby f coinciding with f in a neighborhood of x (therefore $x \in S(g)$) and such that $x \in \text{Per}(g)$. Hence $x \in \text{Per}(g) \cap S(g)$. This contradicts the stability of g . Hence $\overline{L}^+(f) \cap S(f) = \emptyset$. Also it is obvious that $\overline{\text{Per}(f)}$ is prehyperbolic. Therefore $\overline{\text{Per}(f)}$ has a decomposition, $\Lambda_1 \cup \Lambda_2 \cup \cdots \cup \Lambda_n$ into disjoint prehyperbolic sets, following the method of Przytycki [7] and Newhouse [5]. Here each Λ_i is a prehyperbolic set such that the periodic points are dense in Λ_i . Let U_i be a compact neighborhood of Λ_i for $1 \leq i \leq n$. If we choose U_i small enough, we have that $U_i \cap U_j = \emptyset$ if $i \neq j$ and $\overline{L}^+(f) \setminus \bigcup_{1 \leq i \leq n} U_i \neq \emptyset$.

Now suppose that $f|_{\Lambda_i}$ is injective. Then shrinking U_i if necessary, $f|_{U_i}$ is a diffeomorphism onto its image. By Theorem 7.3 of [2], there is a neighborhood $\mathcal{N}_i$ of f in $\text{End}^r(M)$ such that if $g \in \mathcal{N}_i$ then there exists a homeomorphism h close to identity which conjugates Λ_i with a g -invariant subset of U_i .

Next suppose that $f|_{\Lambda_i}$ is not injective. Then it is easy to show that $f|_{\Lambda_i}$ is a (quasi-) expanding map, i.e. $\dim E_x^u = \dim M$ for any $x \in \Lambda_i$. By the similar argument to Przytycki [7], there exists a neighborhood $\mathcal{N}_i$ of f in $\text{End}^r(M)$ such that if $g \in \mathcal{N}_i$ then there exists a homeomorphism h from Λ_i onto its image satisfying $gh = hf$ and $h(\Lambda_i) \subset U_i$.

From above arguments and $\overline{P}$ -stability of f , it follows that there exists a neighborhood $\mathcal{N}$ of f in $\text{End}^r(M)$ such that $\overline{\text{Per}(g)} \subset \bigcup_{1 \leq i \leq n} U_i$ for any $g \in \mathcal{N}$. Let $x \in \overline{L}^+(f) \setminus \bigcup_{1 \leq i \leq n} U_i$. By Lemma 2, there exists $g \in \mathcal{N}$ such that x is a periodic point of g . This is a contradiction.

Subsequently we obtain the Corollary of the Theorem:

COROLLARY. *If $f \in \text{End}^r(M)$ is both C^r structurally and infinitesimally stable, then f satisfies Axiom A and no cycle condition.*

REFERENCES

1. J. Guckenheimer, *Absolutely Ω -stable diffeomorphisms*, Topology **11** (1972), 195–197.
2. M. Hirsch and C. Pugh, *Stable manifolds and hyperbolic sets*, Proc. Sympos. Pure Math., vol. 14, Global Analysis, Amer. Math. Soc., Providence, R.I., 1970, pp. 133–163.
3. R. Mañé, *On infinitesimal and absolute stability of diffeomorphisms*, Dynamical Systems (Warwick, 1974), Lecture Notes in Math., vol. 468, Springer-Verlag, Berlin, Heidelberg and New York, pp. 151–161.
4. —, *Axiom A for endomorphisms*, Lecture Notes in Math., vol. 597, Springer-Verlag, Berlin, Heidelberg and New York, pp. 379–388.
5. S. Newhouse, *Hyperbolic limit sets*, Trans. Amer. Math. Soc. **67** (1972), 125–150.
6. F. Przytycki, *Anosov endomorphisms*, Studia Math. **58** (1976), 249–285.

7. ———, *On Ω -stability and structural stability of endomorphisms satisfying Axiom A*, *Studia Math.* **60** (1977), 61–77.
8. C. Pugh and R. Mañé, *Stability of endomorphisms*, *Lecture Notes in Math.*, vol. 468, Springer-Verlag, Berlin, Heidelberg and New York, 1975, pp. 175–184.

DEPARTMENT OF MATHEMATICS, SCHOOL OF SCIENCE AND ENGINEERING, WASEDA
UNIVERSITY, SHINJUKU, TOKYO, JAPAN